

CIGARETTE SMOKING AND  
CARBON MONOXIDE  
D. M. AVIADO, M.D.

1005051060

**CONFIDENTIAL**

*Background document, not to be  
cited as such*

**CIGARETTE SMOKING AND CARBON MONOXIDE**

**Prepared for the Council for Tobacco Research**

By

DOMINGO M. AVIADO, M. D.  
Professor of Pharmacology  
University of Pennsylvania School of Medicine  
Philadelphia, Pennsylvania 19104

1005051061

January 31, 1973

## CONTENTS

SUMMARY .....	1
I. INTRODUCTORY REMARKS .....	6
Bibliography (Review articles on carbon monoxide and tobacco) .....	9
II. CARBOXYHEMOGLOBIN BLOOD LEVELS AND CIGARETTE SMOKING .....	11
A. Habitual Smokers .....	11
Table 1. Carboxyhemoglobin levels in the blood of habitual smokers .....	14
Bibliography .....	17
B. Influence of smoking on alveolar air and blood levels of carboxyhemoglobin .....	19
Table 2. Acute effects of cigarette smoking on carboxyhemoglobin levels. ....	21
Bibliography .....	22
C. Passive Smoking .....	24
Bibliography .....	26
D. Carbon monoxide content of cigarette smoke .....	27
Table 3. Measurement of carbon monoxide liberated from tobacco .....	29
Bibliography .....	30
III. SOURCES OF CARBON MONOXIDE .....	32
A. Carboxyhemoglobin levels in the blood of nonsmokers .....	32
Table 4. Carboxyhemoglobin levels in the blood of nonsmokers .....	33
Bibliography .....	34
B. Vehicular traffic and smoking .....	36
Table 5. Carboxyhemoglobin blood levels of nonsmokers following exposure	
to vehicular traffic .....	38
Table 6. Carboxyhemoglobin blood levels of smokers following exposure	
to vehicular traffic .....	39
Bibliography .....	40
IV. RESPIRATORY SYSTEM .....	42
A. Acute respiratory effects in humans .....	42
Bibliography .....	44
B. Chronic respiratory effects in humans .....	45
Bibliography .....	46
C. Acute and chronic respiratory effects in animals .....	47
Bibliography .....	49
V. CIRCULATORY SYSTEM .....	51
A. Heart Rate .....	51
Bibliography .....	53
B. Myocardium .....	56
Bibliography .....	59
C. Coronary Circulation .....	61
Bibliography .....	64
D. Systemic Circulation .....	66
Bibliography .....	67
E. Arteries .....	69
Bibliography .....	72
F. Blood Cells and Plasma .....	74
Bibliography .....	76
VI. NERVOUS SYSTEM .....	79
A. Cerebral Blood Vessels .....	79
Bibliography .....	80
B. Eye and Visual Pathways .....	81
Bibliography .....	83
C. Hearing and Auditory Pathways .....	85
Bibliography .....	86
D. Behavior of Man .....	88
Bibliography .....	91
E. Behavior of Animals .....	93
Bibliography .....	94
F. Electroencephalogram in Man and Animals .....	96
Bibliography .....	97

1005051062

#### IV B. Chronic Respiratory Effects in Humans

The medical examination of traffic officers stationed at the Holland Tunnell in New York has provided an opportunity to determine the effects of chronic exposure to 70 ppm carbon monoxide. Sievers et al. (1942) examined 156 such officers and failed to find any evidence of injury to health that was attributable to carbon monoxide exposure. There were no signs or symptoms of respiratory abnormalities. Other reports of elevated carbon monoxide levels in highway tunnels have appeared, but the clinical examination of the traffic officers has not been included (Braja and Trompeo, 1964; D'Arca et al. 1964; Miranda et al., 1967; Yamate and Matsumura, 1968).

Astrup et al. (1968) and Klausen et al. (1968) exposed 8 male subjects to inhalation of 0.5% carboxyhemoglobin, resulting in a blood level of 10%. There were no changes in ventilation, circulation or metabolism.

1005051109



## SUMMARY

I. Introductory remarks. The literature on carbon monoxide has been reviewed to determine its role in cigarette smoking. A total of 3,500 articles dealing with carbon monoxide were examined, half of them appearing between 1966 and 1972, and bibliographies of these are appended to this report. The present status of the interrelationships between cigarette smoking and carbon monoxide is summarized in the following paragraphs by listing a number of generalizations that are grouped into 3 categories: statements that are generally agreed upon or accepted, those that are unsettled or controversial, and those on which no information is available.

II. Carboxyhemoglobin blood levels and cigarette smoking. There is general agreement that cigarette smoke contains carbon monoxide and that a detectable amount of carboxyhemoglobin is present in the blood of habitual smokers. However, the amount of the carboxyhemoglobin content is unsettled and misquoted. The overall mean level for 30 investigations comprising 2,054 subjects is 3.76 % saturation in the blood collected 4 to 12 hours after the last cigarette. In 21 of these investigations nonsmokers were compared with habitual smokers; the net difference was a mean of + 2.19 %, indicating that the blood of habitual smokers contained this amount more than that of nonsmokers. Immediately after smoking one or more cigarettes, the mean peak level is 5.26 %. The phenomenon of "passive smoking" has been examined. The highest level reported for passive smokers was 3.3 % for nonsmokers in a room (170 cu m) with a carbon monoxide concentration of

1005051064

30 ppm from the consumption of 9.5 cigarettes for each of 11 smokers during a period of 2 hours.

There is no information on the carboxyhemoglobin that smokers sustain throughout a 24-hour period. The level before smoking and the peak level after smoking give the range, but what is needed is an integrated level for the 24-hour period.

III. Sources of carbon monoxide. There is agreement that the blood of nonsmokers contains carboxyhemoglobin derived from endogenous and exogenous sources. The latter include vehicular exhaust, which is a major source of carbon monoxide. Of the 26 investigations of blood levels in nonsmokers, the overall mean for 1,662 subjects was 1.45% carboxyhemoglobin and this may increase two- or fourfold, depending on the level of air pollution. The combination of exposure to vehicular traffic and tobacco smoking causes a further elevation of carboxyhemoglobin approximately equal to the sum of exposure to each. The peak blood levels for 446 subjects had an overall mean of 6.8%. The consequence of this combined exposure to carbon monoxide is unsettled. The effect on the subjects may also be the outcome of exposure to lead, ozone, hydrocarbons and nitrogen oxides from vehicular exhaust.

**1005051065**

IV. Respiratory system. There is general agreement that acute exposure producing a blood level of to carbon monoxide/up to 10% carboxyhemoglobin does not influence respiratory minute volume, diffusing capacity or mechanical properties of the lung in healthy subjects. Chronic exposure of tunnel traffic officers to 70 ppm for several years does not influence respiration. The controversial areas are as follows: The

effect of cigarette smoking on patients with chronic lung disease includes an elevated carboxyhemoglobin level and reduction in pulmonary diffusing capacity.

That carbon monoxide per se causes disturbance of lung function has not been proven in animals. Exposure of dogs to a mixture of 8 to 14 % carbon monoxide air did not produce in/ ultrastructural changes, but exposure of rats produced swelling of alveolar epithelial mitochondria and nucleoplasm. There is no information on ultrastructural changes in primates which will help resolve differences in species.

V. Circulatory system. It is generally agreed that acute carbon monoxide poisoning produces abnormalities in the electrocardiogram and myocardial lesions. Acute exposure to carbon monoxide, resulting in 10 % carboxyhemoglobin saturation, does not influence heart or cardiac output. There are several unsettled claims. That the carbon monoxide content of cigarette smoke causes coronary heart disease has not been proven in animals. On the contrary, exposure of dogs to 100 ppm carbon monoxide for 14 weeks did not produce any exaggeration of the myocardial infarction that was experimentally induced by coronary embolization. A retrospective study of patients in Los Angeles did not show a relationship between carbon monoxide levels and the occurrence of acute myocardial infarction. There is no information on the role of carbon monoxide contained in cigarette smoke in provoking an anginal attack. The investigation comparing cigarette smoking with exposure to vehicular traffic included several constituents. A more direct approach is to compare cigarette smoking with inhalation/monoxide to obtain equivalent levels of carboxyhemoglobin

1005051066

in the same subject. The increase in capillary permeability induced in human subjects exposed to carbon monoxide alone needs to be compared with exposure to cigarette smoking.

The development of arteriosclerosis by carbon monoxide is another controversial area. The positive evidence is based on animals fed with cholesterol that develop arteriosclerosis following chronic exposure to carbon monoxide. However, examination of individuals who have been exposed to an environment of up to 1,000 ppm with blood levels of 2 to 26 % carboxyhemoglobin for an average duration of 10.5 years did not reveal any early onset of arteriosclerosis. One possible explanation of the difference between short-term exposure of animals and long-term exposure of humans is the development of tolerance to carbon monoxide in the latter. No information is available on development of tolerance to carbon monoxide associated with cigarette smoking.

VI. Nervous system. There is general agreement that acute carbon monoxide poisoning causes lesions of cerebral blood vessels, eye and visual pathways, auditory system and brain. There is also agreement that the following effects appear with carboxyhemoglobin levels of 5 to 10 %: impairment of critical flicker fusion frequency and alteration of psychomotor abilities. No information is available as to the effect of inhalation of carbon monoxide at the same carboxyhemoglobin levels as that associated with cigarette smoking.

VII. Other organ systems. There is general agreement that acute carbon monoxide poisoning causes disturbance of renal function, the endocrine system and the musculo-skeletal-dermal system. In pregnancy, carbon monoxide is

1005051067

- RISPLER and ROSS 1965 (A 303) 195  
 RITTER 1956 (300) 60, 57  
 RITUCCI and LUVONI 1965 (A 694) 220  
 RIZZI 1968 (A 772) 227  
 ROBBINS, BORG and ROBINSON 1968 (A 345) 198  
 ROBIN, RAVENS and BING 1969 (327) 65, 61  
 ROBINSON and ROBBINS 1970 (A 196) 189  
 ROCHE, BERTOYE, VINCENT, MOTIN, GARIN, BOLOT and CHADENSON 1968 (A 833) 231  
 RODKEY 1970 (A 346) 198  
 RODKEY and COLLISON 1970 (A 42) 178  
 RODKEY, COLLISON and ENGEL 1969 (A 711) 223  
 RODKEY, COLLISON and O'NEAL 1971 (A 43) 178; (A 182) 187  
 RONDIA 1970 (A 578) 213  
 RONDIA, GUYAUX and HEUSGHEM 1966 (A 304) 195  
 ROOT 1962 (29) 10, 6  
 ROPSHITZ and OVENSTONE 1968 (A 640) 217  
 ROSE, JONES, JENKINS and SIEGEL 1970 (522) 95, 93  
 ROSE 1969 (A 695) 220  
 ROSE and ROSE 1971 (668) 115, 172  
 ROSENBERG 1968 (86) 23, 20  
 ROSENBERG 1971 (87) 23, 20  
 ROSENBERG 1972 (88) 23, 20  
 ROSENBLUTH 1968 (A 524) 209  
 ROSENTHAL 1968 (A 696) 220  
 ROSKAMM 1964 (301) 60, 56  
 ROSSI-FANELLI and ANTONINI 1958 (631) 112, 111  
 ROSSIN and ROBERTS 1972 (30) 10, 7  
 ROSSO and DUGHIERA 1964 (A 712) 223  
 ROUCH, RIOUFOL and BOURBON 1971 (54) 18, 13, 15; (160) 35, 33a  
 ROUGHTON 1970 (A 141) 184  
 RUBINO 1964 (226) 50, 47  
 RUDOLPH, BOYLE, DRESDEN and GILL 1972 (632) 112, 11  
 RUEL and BARTHE 1954 (55) 18, 14; (161) 35, 33  
 RUHL and LIN 1936 (56) 18, 14; (89) 23, 20, 21  
 RUMEN and CHANCE 1970 (A 142) 184  
 SADOKIERSKI 1965 (A 713) 223  
 SAITA and LUSSANA 1971 (435) 80, 79  
 SALNIS and HACHATUROV 1970 (A 773) 227  
 SANDERS and WARRINGTON 1971 (A 525) 209  
 SANNA-RANDACCIO and NISSARDI 1969 (A 442) 205  
 SANZHIEVA 1970 (A 608) 215  
 SANZHIEVA and ZAVARZIN 1971 (A 609) 215  
 SARACOGU 1951 (262) 55, 51  
 SARTORELLI 1967 (A 403) 205  
 SARUTA 1937 (136) 31, 29  
 SASAKI, HIRANO, NAGAHAMA and USUI 1966 (544) 98, 96  
 SATAKE, IIDA, TATSUBANA, YAMAZAKI and MATSUOKA 1968 (A 444) 205  
 SATO 1966 (471) 86, 85  
 SATOH, KIYOTANI, MINAFI and KONDO 1966 (A 697) 220  
 SAVATEEV, TONKOPIJ and FROLOV 1970 (A 834) 231  
 SAYERS and DAVENPORT 1930 (31) 10, 6  
 SAYERS, YANT, LEVY and FULTON 1929 (495) 91, 88  
 SCHAEFER 1964 (A 183) 187  
 SCHAFFERNICHT, AIEGLER and REINHARD 1970 (A 714) 223  
 SCHIECHE, KEBLER and KOBER 1970 (A 715) 223  
 SCHIEVELBEIN 1968 (32) 10, 7  
 SCHIEVELBEIN and EBERHARDT 1972 (328) 65, 61  
 SCHLECHT 1971 (A 610) 215  
 SCHMELZER, STEINER, MAYER, NEDETZKA and FASOLD 1972 (A 143) 184  
 SCHMIDT 1970 (454) 83, 82  
 SCHMIDT 1971 (107) 26, 25  
 SCHMIDT 1939 (57) 18, 13, 14; (90) 23, 21; (162) 35, 33  
 SCHMIDT 1940 (58) 18, 14; (91) 23, 21  
 SCHOTT, TOMMASI, BOURRAT and MICHEL 1967 (A 526) 209  
 SCHRAUZER and LEE 1970 (A 611) 215  
 SCHRENK 1942 (59) 18, 13, 14  
 SCHULTE 1963 (496) 92, 88  
 SCHULTE 1965 (A 835) 231  
 SCHUTTMANN 1968 (A 716) 223  
 SCOPPETTA 1968 (592) 106  
 SCORER 1971 (A 259) 192  
 SEGAL 1970 (523) 95, 93  
 SELING 1966 (263) 55, 52  
 SELTZER 1970 (569) 115, 174  
 SEMAR, TRESER and LANGE 1967 (A 348) 198  
 SESSA and SANNA 1966 (A 527) 209  
 SCASSELLATI SFORZOLINI and SAVINO 1968 (137) 31, 29  
 SFORZOLINI and SAVINO 1970 (A 305) 196  
 SHAFER, SMILAY and MacMILLAN 1965 (264) 55, 52  
 SHAW, CINKOTAI and THOMSON 1966 (A 445) 205  
 SHIDA and KUROIWA 1969 (A 528) 209  
 SHIELDS 1971 (92) 23, 20  
 SHIMOJIMA 1970 (A 529) 209  
 SHINTANI 1968 (227) 50, 48  
 SHIRABE, MAWATARI and KUROIWA 1970 (A 717) 223  
 SHIRAKI 1969 (A 530) 209  
 SHIRUKI and TATETSU 1967 (A 531) 209  
 SHOJI, YAMAMOTO, NISHIDA, ISHIKAWA, TAKADA and INOUE 1967 (A 306) 196  
 SIASEV 1966 (A 698) 220  
 SIEGEL and MOHLER 1969 (A 184) 188  
 SIEGENTHALER 1965 (A 232) 191  
 SIEGRIST 1966 (A 718) 224  
 SIEVERS, EDWARDS, MURRAY and SCHRENK 1942 (60) 18, 14  
 SIGGAARD-ANDERSEN, KJELDSSEN, PETERSEN and ASTRUP 1967 (387) 73, 70

1005051310

## I. INTRODUCTORY REMARKS

In recent years there has been increasing concern as to the harmful effects of carbon monoxide released as an air pollutant. The importance of cigarette smoking as a source of carbon monoxide has been recently stressed. This review is an attempt to clarify the role of carbon monoxide in cigarette smoking. The relationship will be analyzed in terms of carboxyhemoglobin blood levels and their influence on the respiratory, circulatory, nervous, renal, reproductive, endocrine and musculoskeletal systems.

At the outset it is pertinent to summarize the present state of knowledge relating to carbon monoxide in general and to carbon monoxide in cigarette smoke in particular. The information is summarized in the following publications:

a. The toxicity of carbon monoxide has been reviewed by Sayers and Davenport (1930), Killick (1940), Lilienthal (1950), Root (1962) and Theodore et al. (1971). These review articles have appeared at intervals of a decade and do not include the cigarette smoke as a source of carbon monoxide.

b. The importance of carbon monoxide as an air pollutant has been reviewed by Goldsmith (1964), Kaye (1965), Finck (1966), Giever (1967), Goldsmith and Landaw (1968), Beard (1969), Leclercq and Proteau (1970), Casarett (1971) and Jech (1972). These reviews appearing at yearly intervals emphasize the origin of carbon monoxide poisoning from sources other than cigarette smoking.

1005051069



c. A symposium entitled "Biological Effects of Carbon Monoxide," edited by R. F. Coburn, appeared in 1970. Most investigators active in carbon monoxide research participated in this meeting. There is one article relating to cigarette smoking as a source of carbon monoxide (Lawther and Cummins, 1970).

d. Government agencies have released three documents concerning regulation of carbon monoxide levels. The National Academy of Science and the National Academy of Engineering have jointly released "Effects of Chronic Exposure to Low Levels of Carbon Monoxide on Human Health, Behavior, and Performance" in 1969; the Environmental Health Service has released "Air Quality Criteria for Carbon Monoxide" in 1970; and the National Institute for Occupational Safety and Health has published "Criteria for a Recommended Standard: Occupational Exposure to Carbon Monoxide" in 1972. The major change was the reduction in the threshold limit value from 100 ppm for carbon monoxide to 50 ppm. The following authors have written review articles on regulation of control of carbon monoxide levels: Dinman (1968); Goldsmith and Cohen (1969), DuBois (1970), Grut et al. (1970) and Rossin and Roberts (1972).

e. The carbon monoxide liberated during the use of tobacco is reviewed in the following monographs: "Tobacco: Experimental and Clinical Studies," by Larson et al. (1961), and supplements by Larson and Silvette (1968, 1971); "Tobacco and Tobacco Smoke," by Wynder and Hoffmann (1967); and "Nikotin: Pharmacologie und Toxikologie des Tabakrauches," edited by Schievelbein

1005051070

(1968). In the publications prepared by the National Clearinghouse for Smoking and Health, there has been an increasing amount of space devoted to carbon monoxide ("Smoking and Health," 1964; "The Health Consequences of Smoking," 1967, 1968, 1969, 1971 and 1972). In the 1972 supplement, the danger to the nonsmoker who inhales the carbon monoxide of the smoke liberated by a smoker has been emphasized. The contents of these publications relating to carbon monoxide in cigarette smoke are discussed in section IX below.

f. The bibliography of carbon monoxide was compiled in 1966 by Cooper. Among the 983 articles abstracted, 22 were devoted to carbon monoxide in smoking.

The present review is written primarily to discuss the relationship between cigarette smoking and carbon monoxide. Articles which directly relate to carbon monoxide and cigarette smoking are reviewed in this document. The investigations dealing with objective measurements of carbon monoxide exposure are emphasized, particularly those which relate to the effects of cigarette smoking on body systems (sections II to VIII). The articles which discuss the significance of carbon monoxide in cigarette smoking have been scrutinized for validity of statements. Included in this commentary (section IX) are the statements which appeared in all the volumes of "Smoking and Health" (1967 to 1972). The bibliography for 1966 to 1972 on carbon monoxide in general is appended to complete the reference list started by Cooper for articles appearing up to 1966 (section XI).

1005051071

## BIBLIOGRAPHY

I. INTRODUCTORY REMARKS  
(Review articles on carbon monoxide and tobacco)

- |  | Reprint |
|--|---------|
| BEARD R R : Toxicological appraisal of carbon monoxide. <u>J Air Pollut Contr Ass</u> 19: 722-7, 1969.   | 1       |
| CASARETT L J : Toxicology: the respiratory tract. <u>Ann Rev Pharm</u> 11: 425-446, 1971 (Ed)  | 2       |
| COBURN R F : Biological effects of carbon monoxide. <u>Ann NY Acad Sci</u> 174: 1-430, 1970.   | 3       |
| COOPER A G : Carbon monoxide. A bibliography with abstracts. <u>Pub Health Service</u> Washington, 1-440, 1966.  | 4       |
| DINMAN B D : Pathophysiologic determinants of community air quality standards for carbon monoxide. <u>J Occup Med</u> 10: 446-63, 1968.  | 5       |
| DuBOIS A B : Establishment of "threshold" CO exposure levels. <u>Ann NY Acad Sci</u> 174: 425-8, 1970.   | 6       |
| ENVIRONMENTAL HEALTH SERVICE : Air quality criteria for carbon monoxide. <u>Public Health Service</u> , Washington 1-140, 1970.  | 7       |
| FINCK P A : Exposure to carbon monoxide: review of the literature and 567 autopsies. <u>Milit Med</u> 131: 1513-39, 1966.  | 8       |
| GIEVER P M : Significance of carbon monoxide as an air pollutant. <u>J Occup Med</u> 9: 265-70, 1967.  | 9       |
| GOLDSMITH J R : Uses of medical and epidemiological research in the control and prevention of air pollution. <u>Proc Roy Soc Med</u> 57:Suppl: 1034-40, 1964.  | 10      |
| GOLDSMITH J R and COHEN S I : Epidemiological bases for possible air quality criteria for carbon monoxide. <u>J Air Pollut Contr Ass</u> 19: 704-13, 1969.   | 11      |
| GOLDSMITH J R and LANDAW S A : Carbon monoxide and human health. <u>Science</u> 162: 1352-9, 1968.   | 12      |
| GRUT A, ASTRUP P, CHALLEN P J R and GERHARDSSON G : Threshold limit values for carbon monoxide. <u>Arch Environ Health(Chicago)</u> 21: 542-4, 1970.   | 13      |
| JECH J. : Kyslicnik Uhelnaty a Jeho Vyznam v Ovzdusi Sidlist. (Carbon monoxide and its importance in the urban atmosphere. <u>Ces Hyg</u> 17(2/3): 93-99, 1972.  | 14      |
| KAYES : The menace of carbon monoxide. <u>Bol Asoc Med P Rico</u> 57: 626-30, 1965   | 14a     |
| KILLICK E M : Carbon monoxide anoxemia. <u>Physiol Rev</u> 20: 313-44, 1940.   | 15      |
| LARSON P S, HAAG H B and SILVETTE H : Tobacco. Experimental and clinical studies. A comprehensive account of the world literature. <u>Williams &amp; Wilkins Company</u> , Baltimore, 6; 107-9; 252; 479; 497; 651-2, 1961.    | 16      |
| LARSON P S and SILVETTE H : Tobacco. Experimental and clinical studies. A comprehensive account of the world literature. Supplement I. <u>The Williams &amp; Wilkins Company</u> , Baltimore 62-3; 136; 244; 254, 1968.        | 17      |
| LARSON P S and SILVETTE H : Tobacco. Experimental and clinical studies. A comprehensive account of the world literature. Supplement II. <u>The Williams &amp; Wilkins Company</u> , Baltimore 44-45; 116; 204; 212; 338, 1971. | 18      |
| LAWTHER P J and COMMINS B T : Cigarette smoking and exposure to carbon monoxide. <u>Ann NY Acad Sci</u> 174: 135-147, 1970.  | 18a     |

1005051072

- LECLERQ A and PROTEAU J: Intoxication par l'oxyde de carbone. (Poisoning by carbon monoxide.) Concours Med 92:8955-8966, 1970 19
- LILIENTHAL J L Jr: Carbon monoxide. Pharmacol Rev 2: 324-54, 1950. 20
- NATIONAL ACADEMY OF SCIENCES & NATIONAL ACADEMY OF ENGINEERING: Effects of chronic exposure to low levels of carbon monoxide on human health, behavior, and performance. National Academy of Sciences and National Academy of Engineering, Washington 1-66, 1969. 21
- NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS: Smoking and Health. Report of the advisory committee to the Surgeon General of the Public Health Service. Public Health Service, Washington 60; 343; 344, 1964. 22
- NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS: The health consequences of smoking. A Public Health Service review: 1967. Public Health Service, Washington 62-64, 1967. 23
- NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS: The health consequences of smoking. 1968 supplement to the 1967 Public Health Service Review. Public Health Service, Washington 38-40, 1968. 24
- NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS: The Health consequences of smoking. 1969 supplement to the 1967 Public Health Service Review. Public Health Service, Washington 26-9; 80, 1969 25
- NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS: The health consequences of smoking. A report of the Surgeon General: 1971. Public Health Service, Washington 8-9; 59-62, 1971. 26
- NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS: The health consequences of smoking: A report of the Surgeon General: 1972. Public Health Service, Washington 21-3; 125-8, 1972. 27
- NATIONAL INSTITUTE FOR OCCUPATIONAL SAFETY AND HEALTH: Criteria for a recommended standard: occupational exposure to carbon monoxide. Public Health Service, Washington 1-66, 1972. 28
- ROOT W S: Carbon monoxide. Handbook of Physiology, Section 3, Respiration. Edited by Fenn, W D and Rohn, H. Williams & Wilkins Company, Baltimore 1087-98, 1962. 29
- ROSSIN D A and ROBERTS J J: Episode control criteria and strategy for carbon monoxide. J Air Pollut Cont Ass 22: 254-9, 1972. 30
- SAYERS R R and DAVENPORT S J: Review of carbon monoxide poisoning. US Pub Health Bull 195: 1-97, 1930. 31
- SCHIEVELBEIN H, Editor: Nikotin. Pharmakologie und Toxikologie des Tabakrauches. Georg Thieme Verlag, Stuttgart 293-320, 1968. 32
- THEODORE J, O'DONNELL R D and BACK K: Toxicological evaluation of carbon monoxide in humans and other mammalian species. J Occup Med 13: 242-55, 1971. 33
- WYNDER E L and HOFFMANN D: Tobacco and tobacco smoke. Studies in experimental carcinogenesis. Academic Press, New York and London 442-444, 1967. 34

1005051073

## II. CARBOXYHEMOGLOBIN BLOOD LEVELS AND CIGARETTE SMOKING

The most important single measurement for assessing the amount of exposure to carbon monoxide is the blood level of carboxyhemoglobin. The portion of hemoglobin present in the form of carboxyhemoglobin cannot combine with oxygen and therefore cannot carry oxygen from the lungs to the tissues. The techniques for measuring carboxyhemoglobin are discussed in the articles under additional bibliography list no. 1. The nature of the binding between carbon monoxide and hemoglobin resulting in a shift of the oxyhemoglobin dissociation curve to the left is covered by additional bibliography list no. 2. The release of hemoglobin from carboxyhemoglobin during the elimination of carbon monoxide and its oxidation by living tissues are covered by the publications in additional bibliography list no. 3. The accumulation of carbon monoxide in enclosed spaces, such as an aircraft, submarine or tunnel, is covered by additional bibliography list no. 4. This section will discuss the details of the investigation that relate directly to cigarette smoking and its consequences on the level of carboxyhemoglobin in the blood.

**1005051074**

### A. Habitual Smokers

The first identification of carboxyhemoglobin in blood of smokers was accomplished by Hartridge (1919-1920). He estimated the level in one smoker to be 6% and in another to be absent. A more extensive investigation was completed by Gettler and Mattice (1933), who compared 4 groups of habitual cigarette smokers. The group of 12 rural dwellers had a mean value

of 1.2% carboxyhemoglobin in the blood, while 18 New York City residents had a mean of 1.4%, 12 New York City cleaners a mean of 3.5%, and 2 New York City taxi drivers a mean of 13.5%. These results proved for the first time that the carboxyhemoglobin contained in blood of habitual smokers was not only the result of cigarette smoking but was also the outcome of inhaling an atmosphere containing carbon monoxide released from automobile exhaust and other sources.

There have been 28 additional reports of carboxyhemoglobin in blood levels among cigarette smokers and the results are summarized in Table 1. The mean values do not represent the effect of cigarette smoking because the contribution of carbon monoxide in the atmosphere has to be subtracted. The last column in Table 1 is the net level of carboxyhemoglobin which can be attributed to cigarette smoking and was obtained by subtracting the mean blood levels for controls who were nonsmokers. The following generalizations can be made from the published results.

1. The 30 investigations summarized in Table 1 were performed in various cities in the United States and Europe. The highest mean level is 16.2% for a group of 6 U. S. Army enlisted men (Meigs, 1948). This represents an error in the analysis, since the blood levels for nonsmokers were also high. All the other mean levels were below 10% carboxyhemoglobin.

2. The overall mean level for 2,054 subjects reported in the 30 investigations is 3.76%. This represents the average blood level for smokers in the morning, 4 to 12 hours after they smoked a cigarette.

**1005051075**

3. The contribution of atmospheric pollution to the increased blood levels of carboxyhemoglobin can be derived by subtracting the blood levels for controls



who were nonsmokers. Twenty-one of the investigations included nonsmokers, so that it was possible to subtract their mean levels from those of habitual smokers. The net difference between 2 groups represents the contribution from smoking alone, which amounted to a mean of +2.19% of carboxyhemoglobin for 2,781 subjects. This mean value was calculated regardless of the number of cigarettes consumed.

4. Ten investigators related the amount of cigarettes consumed daily to blood levels of carboxyhemoglobin. The consumption of 20 or less cigarettes per day showed the following net change in blood carboxyhemoglobin levels in each case: +1.6% (Schmidt, 1939); +2.4% (Schrenk, 1942); +2.1% (Parmeggiani and Gilardi, 1952); +1.0% (Goldsmith et al, 1963); +1.9% (Curphey et al, 1965); +0.3% (Balbo et al, 1966); and +0.9% (Rouch et al, 1971). The net changes in blood carboxyhemoglobin levels for subjects consuming one or more cigarettes were respectively: +2.9% (Schmidt, 1939); +3.7% (Schrenk, 1942); +3.5% (Parmeggiani and Gilardi, 1952); +1.1% (Goldsmith et al, 1963); +3.6% (Curphey et al, 1965); +2.0% (Balbo et al, 1966); and +8.0 (Yacoub et al, 1970). The last-mentioned value represents the highest net level of carboxyhemoglobin, next to the +11.0% referred to above as reported for the U. S. Army enlisted men.

(Table 1 appears on the next page.)

1005051076

Table 1 Carboxyhemoglobin levels in the blood of habitual smokers.<sup>1</sup>

Reference (Year)	Nature of habitual smokers (Cigarette consumption)	No. of Subjects	Carboxyhemoglobin blood level %		
			Smokers Mean $\pm$ SD (Range)	Non-smokers <sup>2</sup> Mean	Net $\Delta$
Hartridge (1919-1920)	London volunteers	2	3.0 (0-6)		
Gettler and Mattice (1933)	New York residents	18	1.4 (1.0-4.1)		
	New York street cleaners	12	3.5 (1.2-0.9)		
	New York taxi drivers	2	13.5 (8.0-19)		
	Rural dwellers	12	1.2 (0.5-3.6)		
Ruhl and Lin (1936)	Berlin volunteers				
	(non-inhalers in morning)	21	0.6		
	(inhalers in morning)	25	0.5		
	(heavy inhalers in morning)	13	1.3		
Schmidt (1939; 1940)	Bonn volunteers				
	(20-30/day)	3	3.5	0.6	+2.9
	(> 30/day)		7.2		+6.6
	(10-20/day)		2.2		+1.6
Schrenk (1942); Sievers, Edwards Murray and Schrenk (1942)	Holland tunnel workers				
	(<20/day)	39	4.1 $\pm$ 1.9	1.7	+2.4
	(>20/day)	21	5.4 $\pm$ 1.8		+3.7
	(pipe)	5	2.5		
Wennesland 1945	(cigar)		3.2		
	Stockholm volunteers				
	(<15/day)	35			
	(pipe)	7			
Meigs 1948	US army enlisted men				
	(6-20/day)	6	16.2 (1.9-45)	5.2	+11.0
Parmeggiani and Gilardi (1952)	Italian volunteers				
	(10-12/day)	14	4.9 (1-9)	2.8	+2.1
	(15-25/day)	6	6.3 (2-9.5)		+3.5
	(30-40/day)	3	9.3 (6-14)		+6.5
Barthe, Paris, Duchemin and Thomas (1953)	Paris workers	100	(0.8-2.0)		
Ruel and Barthe (1954)	Paris workers				
	(<10/day)		1.0		
	(10-15/day)		1.7		
Valic and Duric (1954)	(>20/day)		2.7		
	Yugoslavian workers	75	3.8 $\pm$ 1.9	0.9	+2.9
Gaensler, Cadigan, Illicott, Jones and Mark (1957)	Boston workers	9	4.7 (3.1-7.9)	0.9	+3.8
	Stockholm volunteers	6	2.1 (1.2-3.0)	0.7	+1.4
	Öhrberg and Rothschild (1958)				

1005051077

Table 1 (continued)

page 15

Reference (Year)	Nature of habitual smokers (Cigarette consumption)	No. of Subjects	Carboxyhemoglobin blood level %			
			Smokers Mean $\pm$ SD (Range)	Non-smokers <sup>2</sup> Mean	Net	
Trinder and Harper (1962)	Sunderland volunteers	3	3.8 (3.2-4.8)	0.9	+2.9	
Hofreuter, Catcott and Xintaras (1962)	Cincinnati volunteers	19	2.9	1.9	+1.0	
Goldsmith, Schuette and Novick (1963)	San Francisco longshoremen ( $<10$ /day)	429	2.3	1.3	+1.0	
	(10-40/day)	1035	3.4		+1.1	
	( $>40$ /day)	233	5.5		+4.2	
Curphrey, Hood and Perkins (1965)	Los Angeles longshoremen (light)	55	2.3	0.4	+1.9	
	(medium)	153	3.0		+2.6	
	(heavy)	29	4.0		+3.6	
Ayres, Gianelli and Armstrong (1965)	New York volunteers	25	4.2	0.9	+3.3	
Balbo, Marucci and Ronchi (1966)	Paris workers (5/day)	34	2.1 (0-9.8)	1.9	+0.2	
	(10/day)	32	2.2 (0-9.5)		+0.3	
	(15/day)	16	2.2 (0-5.9)		+0.3	
	(20/day)	20	2.2 (0-6.9)		+0.3	
	(30/day)	7	2.8 (0-5.2)		+2.0	
McIlvaine, Nelson and Bartlett (1969)	Durham workders	5	3.8	1.6	+2.2	
Bhown, Maitrya and Haq (1969)	Indian beedi smokers ( $<10$ /day)	7	4.8 (3.4-5.6)			
	(10-19/day)	8	5.9 (5.6-6.1)			
	(20-29/day)	8	6.9 (6.6-7.3)			
	( $>29$ /day)	7	9.4 (8.5-10.5)			
Yacoub, Faure, Mallion and Cau (1970)	Paris workers (20/day inhaled)	90	9.5	1.5	+8.0	
	(20/day non-inhaled)	97	6.0		+4.5	
Rouch, Rioufol and Bourbon (1971)	Toulouse volunteers ( $<10$ /day)	5	2.5 (1.0-5.5)	1.6	+0.9	
	( $>10$ /day)	15	4.25 (1.0-11.0)		+2.6	
Brewer, Eaton, Weil and Grover (1970); Brewer, Eaton, Grover and Weil (1971)	Leadville volunteers	20	6.6 $\pm$ 2.7			
Weiss, Slawsky and Desforges (1971)	Boston patients with fibrosis	8	4.2 $\pm$ 2.7	1.5	+4.2	
Butley (1971)	Los Angeles residents	81	5.7 (3.2-14.2)	1.5	+4.2	
Hansen, Wilke, Malorny and Gøthert (1972)	Hamburg workers	40	4.9 $\pm$ 1.0	0.75	+4.2	

1005051078

Reference (Year)	Nature of habitual smokers (Cigarette consumption)	No. of Subjects	Carboxyhemoglobin blood level %			
			Smokers Mean $\pm$ SD (Range)		Non-smokers <sup>1</sup> Mean	Net $\Delta$
Curson, Garby, Robert and Zaar (1972)	Uppsala volunteers	6	2.7	(1.5-4.4)	1.27	+1.4
Mean (overall for number of subjects)			3.76 (2954 subjects)			+2.19 (2781 subjects)

<sup>1</sup> Some of the values were reported in volumes % and are expressed in this table as saturation % assuming normal hemoglobin values.

<sup>2</sup> See Table 4 for details.

1005051079

(7) ABELSON P H : A damaging source of air pollution. Science 158: 1527, 1967.

One of the toxic products of the automobile is carbon monoxide. Exposure for 1 hour to a concentration of this gas of 120 parts per million causes inactivation of about 5 percent of the body's hemoglobin and commonly leads to dizziness, headache, and lassitude. Concentrations of carbon monoxide as high as 100 ppm often occur in garages, in tunnels, and behind automobiles. Such concentrations are tiny in comparison with those (42,000 ppm) found in cigarette smoke. The smoker survives because most of the time he breathes air not so heavily polluted. However, in a poorly ventilated, smoke-filled room, concentrations of carbon monoxide can easily reach several hundred parts per million, thus exposing smokers and nonsmokers present to a toxic hazard.

In this article, the comparison of concentration of carbon monoxide is as follows: 100 ppm for garages and tunnels and 42,000 ppm in cigarette smoke. The latter represents 4.2 carbon monoxide in pure cigarette smoke. This concentration is not inhaled continuously, but is diluted by air in the lungs at the time of inhalation of cigarette smoke. After exhalation, atmospheric air enters to replace the cigarette smoke. Ringold et al. (1962) analyzed the expired air to determine the integrated concentration of carbon monoxide therein: that of heavy smokers had a concentration of 16.4 ppm, light smokers 7.7 ppm, and non-smokers 0.08 ppm (see page 19).

1005051205

- McILVAINE P M, NELSON W C and BARTLETT D: Temporal variation of carboxyhemoglobin concentrations. Arch Environ Health 19: 8-91, 1969. 50
- MEIGS J W: Carbon monoxide poisoning. Bull US Army 8: 542-6, 1948. 51
- MOTLEY H L: Environmental air pollution effect on pulmonary function. Aerosp Med 42: 1108-10, 1971. 52
- PARMEGGIANI L and GILARDI F: Rilievi sulla ossicarbonemia fisiologica. Med Lav 43: 179-83, 1952. 53
- ROUCH Y, RIOUFOL F and BOURBON P: Oxycarbonemie spontanee et apres exposition pendant quelques minutes a une atmosphere a un pour mille d'oxyde de carbone chez 43 sujets. (Carbon monoxide content in blood: normal and after a few minute exposure to an atmosphere containing 0, 1 p. 100 carbon monoxide/43 subjects). Arch Mal Profes Paris 32: 271-82, 1971. 54
- RUEL M H and BARTHE M R: Examen systematique de 132 ouvriers d'usines a gaz avec dosage du CO sanguin. Arch Mal Profes Paris 15: 234-5, 1954. 55
- RUHL A and LIN P: Zur frage der Kohlenoxydintoxikation bei starken Rauchern. Deutsch Med Wschr 62: 493-7, 1936. 56
- SCHMIDT O: Der gasanalytische nachweis von Kohlenoxyd im Blut, insbesondere bei Rauchern. Klin Wschr 18: 938, 1939. 57
- SCHMIDT O: Der Kohlenoxydgehalt des Blutes bei Rauchern. (The CO content of the blood of smokers). Reichs-Gesundhbl 15: 53-8, 1940. 58
- SCHRENK H H: Results of laboratory tests. Determination of concentration of carbon monoxide in blood. US Pub Health Bull 278: 36-49, 1942. 59
- SIEVERS R F, EDWARDS T I, MURRAY A L and SCHRENK H H: Effect of exposure to known concentrations of carbon monoxide. JAMA 118: 585-8, 1942. 60
- TRINDER P and HARPER F E: A colorimetric method for the determination of carboxyhaemoglobin over a wide range of concentrations. J Clin Path 15: 82-4, 1962. 61
- VALIC F and DURIC D: Concentration of carbon monoxide in the blood of smokers and non-smokers. Arh Hig Rada (Jugoslav) 5: 49-56, 1954. 62
- WEISS E B, SLAWSKY P and DESFORGES J F: Oxyhemoglobin affinity in chronic pulmonary granulomatosis (sarcoidosis) and fibrosis. Am Rev Res Dis 104: 694-702, 1971. 63
- WENNESLAND R: Erfaringer fra kulloksyd-gengass-undersokelser hos norske sjafforer og arbeidere. (Experiences from carbon monoxide -producer gas- investigations on Norwegian auto-drivers and workers). Svenska Lakorliden 42: 397-408, 1945. 64
- YACOB M, FAURE J, MALLION J M and CAU J: Peut-on determiner l'origine tabagique de l'oxycarbonisme chronique? (Is it possible to demonstrate chronic carbon monoxide intoxication specifically caused by smoking?). Med Leg Domm Corpor (Paris) 3: 262-5, 1970. 65

1005051081



## II B. Influence of Smoking on Alveolar Air and Blood Levels of Carboxyhemoglobin

Jongbloed (1939) analyzed the carbon-monoxide content in alveolar air. A trace of carbon monoxide (about 0.0004% or 4 ppm) was detectable in the alveolar air of nonsmokers. In one smoker the content was 10 ppm in the morning before smoking. Then the subject was given 4 cigarettes, each one taking 12 min to smoke with an intervening 12-minute rest period. The results were as follows:

before 1st cigarette	14.0 ppm
after " "	17.9 "
before 2nd " "	19.4 "
after " "	23.2 "
after 3rd " "	27.1 "
after 4th " "	31.5 "
32 min after 4th cigarette	29.0 "
52 min " " "	26.6 "
72 min " " "	23.7 "

There was a progressive increase in the level of carbon monoxide in the alveolar air up to the end of smoking the 4th cigarette. Then a fall occurred in the concentration of carbon monoxide in the alveolar air.

Other investigators have confirmed the observation of Jongbloed. Ringold et al. (1962) analyzed the expired air after a 20-second breath-holding period. The mean  $\pm$  SD levels of carbon monoxide were as follows:

Nonsmokers (93 subjects)	0.8 $\pm$ 4.7 ppm
Heavy smokers (41 subjects)	16.4 $\pm$ 12.6 "
Light smokers (25 subjects) -	7.7 $\pm$ 10.9 "
Pipe or cigar (9 subjects)	3.8 $\pm$ 4.9 "

Cohen et al. (1971) reported similar observations and correlated expired air levels of carbon monoxide with blood levels of carboxyhemoglobin.

The results of investigations relating to blood carboxyhemoglobin

1005051082

levels are summarized in Table 2. The essential features of the table are:

1. The 11 investigations consisting of measurements of carboxyhemoglobin in blood, resulted in a mean peak level of 5.26% in 29 pooled subjects. The net effect of smoking which was calculated in 15 subjects was an increase of 3.81% of carboxyhemoglobin.

2. There are isolated values as high as 12.3% (Earach et al, 1941), 10% (Ruhl and Lin, 1936) and 6.5% (Whitehead and Worthington, 1961). However, since the control value prior to smoking was not measured, it was not possible to calculate the net effect of smoking.

3. None of the investigations summarized in Table 2 determined the duration of the rise in carboxyhemoglobin. On the basis of work concerning experimental inhalation of carbon monoxide mixture, the duration depends on the concentration of inspired air. There is probably a slow fall in carboxyhemoglobin until the next cigarette is smoked.

After cessation of smoking, a reduction occurs in blood levels of carboxyhemoglobin (Belli and Guiliani, 1955; Ramsey, 1972; Rosenberg, 1968, 1971, 1972). The rate of fall in concentration was accelerated by exercise or muscular effort (Castellino, 1955; Crosetti et al, 1966; Shields, 1971). **1005051083**

Animals have been exposed to cigarette smoke for the investigation of its effect. The blood level of carboxyhemoglobin is elevated in mice (Leuchtenberger et al, 1965), hamsters (Dontenwill et al, 1966, 1967) and rats (Driscoll et al, 1972). In the last-mentioned species, the blood level of carboxyhemoglobin was influenced by the nature of cigarette used. With unfiltered smoke, the rat held its breath, whereas filtered smoke was inhaled with continuation of thoracic respiration.

(Table 2 appears on the next page.)

Table 2. Acute effects of cigarette smoking on carboxyhemoglobin levels.<sup>1</sup>

Reference (Year)	Number of Cigarettes	No of Subjects	Peak Carboxyhemoglobin blood levels %		
			Mean	(Range)	Δ
Manson and Hastings (1933)	10-15	6	4.3	(3.1-6.7)	
Ruhl and Lin (1936)	1-2	5		(3.2-10)	
Schmidt (1939, 1940)	1	1			+1.1
	5	2			+1.6; +4
Hsi-Pu and LiMing (1940)	1-5	43		(4.5 in 80% of subjects)	
Barach, Eckman and Molomut (1941)	20	18	5.7	(2.2-12.3)	
MacFarland, Roughton, Halperin and Niven (1944)	1	1			+2.0
	3	1			+4.0
Fabre, Truhaut and Berrod (1951)	24	5	4.85		+2.7
Parmeggiani and Gilardi (1952)	2	1			+3
Goldsmith, Terzaghi and Hackney (1963)	12	1			+11.9
Bowden and Woodhall (1964)	2	2			+1.5
Hamill and O'Neill (1969)	20 (cigars)	1			+13
Mean (overall for number of subjects)		5.26 (29 subjects)			+38 (15 subjects)

<sup>1</sup> Some of the values were reported in volumes % and are expressed in this table as saturation % assuming normal hemoglobin values.

1005051084

## BIBLIOGRAPHY

## II. CARBOXYHEMOGLOBIN BLOOD LEVELS AND CIGARETTE SMOKING

B. Influences of Smoking on Alveolar Air and Blood Levels of Carboxyhemoglobin.

- |   | Reprint |
|---|---------|
| BARACH A L, ECKMAN M and MOLOMUT N: Modification of resistance to anoxia with especial reference to high altitude flying. <u>Am J Med Sci</u> 202: 336-41, 1941.  | 66      |
| BELLI R and GUILLANO V: Ricerche sull'ossicarbonismo. Ossicarbonemia da fumo di tabacco e diagnosi di ossicarbonismo cronico. <u>Fol Med (Napoli)</u> 38: 351-8, 1955.  | 67      |
| BOWDEN C H and WOODHALL W R: The determination and significance of low blood carboxyhaemoglobin levels. <u>Med Sci Law</u> 4: 98-107, 1964.   | 68      |
| CASTELLINO N: Modifiche della carbossiemia in seguito alla inalazione del fumo di tabacco. <u>Fol Med (Napoli)</u> 38: 1014-24, 1955.   | 69      |
| COHEN S I, PERKINS N M, URY H K and GOLDSMITH J R: Carbon monoxide uptake in cigarette smoking. <u>Arch Environ Health</u> 22: 55-60, 1971.   | 70      |
| CROSETTI L, RUBINO G F and PETTINATI L: Osservazioni in tema di detossicazione dell'emoglobina nell'uomo esposto a rischio prolungato da ossido di carbonio. <u>Minerva Med</u> 57: 268-9, 1966.                  | 71.     |
| DONTENWILL W, RECKZEH G and STADLER L: Inhalationsexperimente mit Cigarettenrauch. <u>Beitr Tabakforsch</u> 3: 438-48, 1966.  | 72      |
| DONTENWILL W, RECKZEH G and STADLER L: Berauchungsapparatur fur Laboratoriumstiere. (Smoking apparatus for laboratory animals). <u>Beitr Tobakforsch</u> 4: 45-9, 1967.   | 73      |
| DRISCOLL P, DEUBER A, BAETTIG K and GRANDJEAN E: Effect of filtered cigarette smoke on rats. <u>Nature</u> 237: 37-8, 1972.   | 74      |
| FABRE R, TRUHAUT R and BERROD F: Une nouvelle methode de dosage de l'oxyde de carbone dans le sang. Applications a l'etude de l'oxycarbonemie normale et pathologique. <u>Ann Pharm (France)</u> 9: 625-38, 1951. | 75      |
| and HACKNEY J D   |         |
| GOLDSMITH J R, TERZAGHI J: Evaluation of fluctuating carbon monoxide exposures. <u>Arch Environ Health</u> 7: 647-63, 1963.   | 76      |
| HAMILL W and O'NEILL R P: Carbon monoxide intoxication in cigar smokers. <u>Irish J Med Sci</u> 8: 273-7, 1969.   | 77      |
| HANSON H B and HASTINGS A B: The effect of smoking on the carbon monoxide content of blood. <u>JAMA</u> 100: 1481, 1933.  | 78      |
| HSI-PU S and LI-MING: Carbon monoxide content of the blood after smoking. <u>J Oriental Med</u> 32: 1135-9, 1940.   | 79      |
| JONGBLOED J: Uber den Kohlenoxydgehalt der Alveolarluft beim Tabakrauchen. <u>Arch Int Pharmacodyn</u> 63: 346-58, 1939.  | 80      |
| LEUCHTENBERGER C, LEUCHTENBERGER R and WEISS S: Increase of carbon monoxide (CO) in blood of mice exposed to inhalation of cigarette smoke. <u>Proc Am Ass Canc Res</u> 6: 39, 1965.                              | 81      |
| MacFARLAND R A, ROUGHTON F J W, HALPERIN M H and NIVEN J L: The effects of carbon monoxide and altitude on visual thresholds. <u>J Aviation Med</u> 15: 381- , 1944.  | 82.     |
| PARMEGGIANI L and GILARDI F: Rilievi sulla ossicarbonemia fisiologia. <u>Med Lav</u> 43: 179-83, 1952.  | 83      |
| RAMSEY A: Fall in carbon monoxide blood levels after stopping smoking. <u>Lancet</u> I: 755, 1972.  | 84      |

1005051085

- RINGOLD A, GOLDSMITH J R, HELWIG H I, FINN R and SCHUETTE F: Estimating recent carbon monoxide exposures. A rapid method. Arch Environ Health 5: 308-18, 1962. 85
- ROSENBERG A: Kontrollert tobakafvaening. (Controlled tobacco detoxication). Ugeskrift for Laeger 130: 2014, 1968. 86
- ROSENBERG A: Synpunkter pa rokavvanjning - erfarenheter fran Kopenhamn. (Viewpoints on smoking withdrawal - Copenhagen experiences. Soc-Med Tidskrifts 2: 108-10, 1971. 87
- ROSENBERG A: Fall in carbon monoxide blood levels after stopping smoking. Lancet 1: 593, 1972. 88
- RUHL A and LIN P: Zur frage der Kohlenoxydintoxikation bei starken Rauchern. Deutsch Med Wschr 62: 493-7, 1936. 89
- SCHMIDT O: Der gasanalytische nachweis von Kohlenoxyd im Blut, insbesondere beim Rauchen. Klin Wschr 18: 938, 1939. 90
- SCHMIDT O: Der Kohlenoxydgehalt des Blutes bei Rauchern. (The CO content of the blood of smokers). Reichs-Gesundhbl 15: 53-8, 1940. 91
- SHIELDS C E: Elevated carbon monoxide levels from smoking in blood donors. Transfusion 11: 89-93, 1971. 92
- WHITEHEAD T P and WORTHINGTON S: The determination of carboxyhaemoglobin. Clin Chim Acta 6: 356-9, 1961. 93

1005051086

### II C. Passive Smoking

The phenomenon of "passive smoking" refers to the exposure of nonsmokers in an enclosed space containing smoke generated by cigarette smoking. The consequence to the individual in such a situation was first described by Jones et al. (1923), who studied the occupants of a non-ventilated room of approximately 1,000 cu ft capacity, filled with smoke liberated from cigars, cigarettes, stogies or pipe tobacco. The highest concentration of carbon monoxide was 200 ppm and the highest concentration of carboxyhemoglobin in the blood was 5%. Within 15 min, the occupant became uncomfortable and in 45 to 60 min it was necessary for him to wear goggles to prevent eye irritation.

Yaglou (1955) determined the rate of ventilation for a room of 1,410 cu ft, which contained 3 to 4 smokers consuming 24 cigarettes per hr. The passive smokers suffered from headache, watering of the eyes, irritation of the nose and throat, a feeling of depression, and loss of concentration power for reading, when the ventilation was less than 15 cfm per smoker. A ventilation of 25 cfm per smoker was considered as acceptable in preventing these complaints.

There were no analyses of blood reported for the subjects. **1005051087**

The consequences of passive smoking have been quantitatively measured by Oettel (1967), Markiewicz (1970) and Harke (1970). All three investigators measured the amount of nicotine absorbed by the passive smoker. Only Harke (1970) determined carboxyhemoglobin levels in the blood of active and passive smokers. In a non-ventilated room, 57 cubic meters in size, filled with smoke liberated from 42 cigarettes for 16 to 18 min, the level would reach as high as



48 ppm. In another room, with a size of 170 cu m, 11 smokers each consumed 9.5 cigarettes within 2 hours. In the presence of a carbon monoxide concentration of 30 ppm, the 11 active smokers had a mean carboxyhemoglobin level of  $7.5 \pm 0.8\%$  in the blood, while the 7 passive smokers had a level of  $3.3 \pm 1.4\%$ . With a room concentration of 5 ppm, the respective levels were  $5.8 \pm 1.6\%$  and  $3.3 \pm 1.4\%$ ; with concentrations of  $< 5$  ppm, the blood levels were respectively  $5.0 \pm 1.8\%$  and  $2.7 \pm 1.2\%$ . The effect of exposing the passive smokers to from 5 to 30 ppm caused a net increase of 0.5% in carboxyhemoglobin.

From the above results, Harke (1970) concluded that it is unlikely to find nonsmokers in a room absorbing a significant amount of cigarette smoke. The criticisms by Hess (1971), Schmidt (1971) and Portheine (1971) were largely directed against the measurement of nicotine absorption. The comments regarding carbon monoxide were answered in a rebuttal by Harke (1971), so that the carboxyhemoglobin levels remain valid.

The "passive smoking" situation has been determined by measurement of carbon monoxide in the air. Dublin (1972) reported a concentration of 20.5 to 32.5 ppm in a position next to the smoker in a room ventilated 12.5 times per hour. At the far end of the room the concentration was 13 to 17 ppm. Passive smoking has also been investigated in rats, mice and hamsters (Reckzeh et al., 1969). In these studies, nicotine was also absorbed by the passive smoker.

Harke 72

1005051088

## II. CARBOXYHEMOGLOBIN BLOOD LEVELS AND CIGARETTE SMOKING

C. Passive Smoking

- |  | Reprint |
|--|---------|
| DUBLIN W B : Unwilling smoking. <u>California Med</u> 117(1): 76-7, 1972.  | 94      |
| HARKE H P : Zum problem des "Passivrauchens". (On the problem of passive smoking).<br><u>Munch Med Wschr</u> 112: 2328-34, 1970.   | 95      |
| HARKE H P : Zum problem des "Passiv-Rauchens". (The problem of passive smoking). <u>Munch Med Wschr</u> 113: 710-3, 1971.  | 100     |
| HESS H : The problem of passive smoking. Remarks on H. P. Harke, <u>Munch Med Wschr</u> 112: 2328-34, 1970. <u>Munch Med Wschr</u> 113: 705-6, 1971.   | 101     |
| JONES G W, YANT W P and BERGER L B : Carbon monoxide hazards from tobacco smoke. <u>US Bur Mines Rep Invest</u> 2539: 6, 1973.   | 102     |
| MARKIEWICZ K : Bierni palacze. O szkodliwosci biernie wdychanego dymu z papierosow palonych przez innych. (Passive smokers. The harm from passively inhaled smoke of cigarettes smoked by others. <u>Polski Tygodnik Lek</u> 25(52) 2041-2, 1970.  | 103     |
| OETTEL H : Erkrankungsrisiko des passiven Rauchens? (Risk of sickness through passive smoking?) <u>Deutsch Med Wschr</u> 92: 2042-3, 1967.   | 104     |
| PORTHEINE F : The problems of passive smoking. Comments on H. P. Harke, <u>Munch Med Wschr</u> 112: 2328-34, 1970. <u>Munch Med Wschr</u> 113: 707-9, 1971.  | 105     |
| RECKZEH G, RUCKER K, HARKE H P and DONTENWILL W : Untersuchungen zur Bestimmung der akuten und chronischen Toxizitat von Zigarettenrauch bei passiver Berauchung von Versuchstieren. (Studies on determination of acute and chronic toxicity of cigarette smoke in passive smoke inhalation of experimental animals). <u>Arzneimittelforschung</u> 19: 237-41, 1969. | 106     |
| SCHMIDT F : The problem of passive smoking. An opinion about H. P. Harke, <u>Munch Med Wschr</u> 112: 2328-34, 1970. <u>Munch Med Wschr</u> 113: 702-5, 1971.  | 107     |
| YAGLOU C P : Ventilation requirements for cigarette smoke. <u>Trans Am Soc Heat Air Cond Eng</u> 61: 25-32, 1955.  | 108     |

1005051089

## II. D. Carbon Monoxide Content of Cigarette Smoke

The presence of carbon monoxide in cigarette smoke and its absorption by the blood were recognized as early as 1899 by Wahl. By shaking a few drops of blood in 2 or 3 mouthfuls of tobacco smoke, he studied the uptake of carbon monoxide from tobacco smoke by the blood of smokers. The presence of carbon monoxide was confirmed by spectroscopy. Wahl concluded that the quantity taken up was insufficient to produce any ill effects. A similar conclusion was arrived at by Lehmann (1908), Culverwell (1915) and Dixon (1927 a and b).

A more direct approach has been to measure the amount of carbon monoxide liberated from cigarette smoke. One gram of tobacco yields a variable amount of carbon monoxide, depending on the method of obtaining the tobacco smoke, the rate of smoking, the closeness of pack and the nature of use, i. e., by cigarette, pipe or cigar. Table 3 summarizes the results of 10 investigations of the amount of carbon monoxide emanating from cigarettes. The mean values range from less than 1 ml to 62 ml of carbon monoxide per gram of tobacco. A convenient median level is 25 ml, which would mean that a cigarette weighing about 1 g, burning in a space of 1 cu m, would result in a concentration of 25 ppm carbon monoxide. In a larger space of 100 cu m, it would require 100 cigarettes to approach a concentration of 25 ppm, or 200 cigarettes one of 50 ppm, assuming that there is no ventilation and that no carbon monoxide is absorbed by the smokers or by the objects inside the room. This calculation, based on the amount of carbon monoxide liberated from a cigarette, makes it unlikely that enough carbon monoxide can be contained in a room to be a hazard to nonsmoking occupants. Bridge and Corn

1005051090

(1972) have arrived at the same conclusion by monitoring carbon monoxide levels in a chamber containing cigarette smoke generated by a machine.

The concentration of carbon monoxide in the cigarette smoke ranges from 2 to 7% (Table 3). As the cigarette was burned and became shorter, the concentration of carbon monoxide was increased (Jarrell and Burde, 1965; Waltz and Hausermann, 1965; Baxter and Hobbs, 1967; Owen and Reynolds, 1967; Krusznynski and Henriksen, 1969).

The amount of carbon monoxide in cigarette smoke that is absorbed in the lung has been calculated in a number of ways. Bokhoven and Niessen (1961) measured the concentration of carbon monoxide in the expired air of the same individual with and without inhaling the cigarette smoke. The difference in carbon monoxide concentration in 3 individuals, expressed as % absorbed, averaged 82%.

Haebisch (1970) performed a similar analysis and concluded that approximately 7.5 ml of carbon monoxide is absorbed by smoking 45 mm of a cigarette. Dalhamn et al (1968) performed an analysis of smoke administered with a motorized syringe, which drew a 2-second 35 ml puff of smoke, and analyzed the sample before and after the smoke was exhaled by the subject. In 5 subjects,  $54 \pm 12.7\%$  was retained after deep inhalation, while only  $3 \pm 0.7\%$  was absorbed when the smoke was kept in the mouth and prevented from reaching the lungs.

(Table 3 appears on the next page.)

1005051091

Table 3. Measurement of carbon monoxide liberated from tobacco.

Reference (ear)	ml carbon monoxide/ g tobacco	% carbon monoxide in smoke
Toth (1907)	0.1-0.3	
Marcelet (1907)	20-80	
Lee 1908	4.1	
Lehmann (1909)	15-23.5	
Armstrong (1922)	0.89-1.24	
Baumberger (1923)	8.3	
Ehrismann and Abel (1934)	17.25-35	
Saruta (1937)	62.7	
Kohn-Abrest (1949)	40.0	
Osborne, Adamek and Hobbs (1956)		3.3-5.7%
Philippe and Hobbs (1956)		2.8-5.7%
Stenhagen (1959)	(12-53 mg/cigarette)	
Stumpower, Lewis and Touey (1962)		5.1%
Newsome and Keith (1965)		5-11%
Keith and Tesh (1965)		3.2%
Scassellatti, Sforzolini and Savino (1968)		(110 mg/m <sup>3</sup> )
Grob (1968)		3.2%
Harke and Drews (1968)		3.4-7.7%
Otsuka, Fujiwara, Ikawa and Hirayama (1970)		3.6%

1005051092

## BIBLIOGRAPHY

## II. CARBOXYHEMOGLOBIN BLOOD LEVELS AND CIGARETTE SMOKING

D. Carbon Monoxide Content of Cigarette Smoke.

- |  |                |
|--|----------------|
| ARMSTRONG H E : Carbonic oxide in tobacco smoke. <u>Brit Med J</u> 1: 992-3, 1922.   | Reprint<br>109 |
| BAUMBERGER J P : The carbon monoxide content of tobacco smoke and its absorption on inhalation. <u>J Pharm Exp Ther</u> 21: 23-34, 1923a.  | 110            |
| BAXTER J E and HOBBS M E : Investigation of some physico-chemical aspects of cigarette smoke using oxygen isotopes. I. CO and CO <sub>2</sub> from atmospheric oxidation. <u>Tobacco</u> 164: 26-32, 1967.   | 111            |
| BOKHOVEN C and NIESSEN H J : Amounts of oxides of nitrogen and carbon monoxide in cigarette smoke, with and without inhalation. <u>Nature</u> 192: 458-9, 1961.  | 112            |
| BRIDGE D P and CORN M : Contribution to the assessment of exposure of nonsmokers to air pollution from cigarette and cigar smoke in occupied spaces. <u>Environ Res</u> 5: 192-209, 1972.  | 113            |
| CULVERWELL G H : The toxic possibilities of tobacco smoke. <u>Dublin J Med Sci</u> 139, 241-9, 1915.   | 114            |
| DALHAMN T, EDFORS M L and RYLANDER R : Retention of cigarette smoke components in human lungs. <u>Arch Environ Health</u> 17: 746-8, 1968.   | 115            |
| DIXON W E : The tobacco habit. <u>Lancet</u> 2: 881-5, 1927a.  | 116            |
| DIXON W E : The tobacco habit. <u>Canad Med Ass J</u> 17: 1537-9, 1927b.   | 117            |
| EHRISMANN O and ABEL G : Uber den Kohlenoxydgehalt des Tabakrauches. <u>Zschr Hyg</u> 116: 4-10, 1934.   | 118            |
| GROB K : The gas phase of cigarette smoke. <u>Nat Canc Inst Mono</u> 28: 215-20, 1968.   | 119            |
| HAEBISCH H : Die Zigarette als Kohlenmonoxydquelle. (The cigarette as source of carbon monoxide). <u>Arch Toxikol Germany</u> 26: 251-61, 1970.  | 120            |
| HARKE H P and DREWS C J : Eine einfache Methode zur Gewinnung gasformiger Tabakrauchbestandteile. Ihre Anwendung zur Bestimmung des Kohlenmonoxids im Rauch. (A simple method for obtaining gaseous constituents of tobacco smoke. Its application for the determination of carbon monoxide in tobacco smoke. <u>Beitr Tabakforsch</u> 4: 275-7, 1968. | 121            |
| JARRELL J E : A study of the major gaseous constituents in the mainstream smoke of a cigarette. <u>Tobacco</u> 160: 26-32, 1965.   | 122            |
| KEITH C H and TESH P G : Measurement of the total smoke issuing from a burning cigarette. <u>Tobacco</u> 160(5): 26-9, 1965.   | 123            |
| KOHN-ABREST E : Sur l'oxycarbonemie chez les fumeurs. <u>Arch Mal Prof</u> 10: 37- , 1949.   | 124            |
| KRUSZYNSKI A J and HENRIKSEN A : The quantitative determination of carbon monoxide in tobacco smoke. <u>Beitr Tabakforsch</u> 5(1): 9-11, 1969.  | 125            |
| LEE W E : The action of tobacco smoke, with special reference to arterial pressure and degeneration. <u>Quar J Exp Physiol London</u> 1: 335-59, 1908.   | 126            |
| LEHMANN K B : Untersuchungen über das Tabakrauchen. <u>Munch Med Wschr</u> 55: 723-5, 1908.  | 127            |
| LEHMANN K B : Chemische und toxikologische Studien über Tabak, Tabakrauch und das Tabakrauchen. <u>Arch Hyg Munch</u> 68: 319-420, 1909.   | 128            |
| MARCELETH H : Sur le dosage de l'oxide de carbone en particulier dans les fumees de tabac. <u>These Montbellier</u> 32pp. 1902.  | 129            |

1005051093

- MUMPOWER R C, LEWIS J S and TOUEY G P : Determination of carbon monoxide in cigarette smoke by gas chromatography. Tobacco Sci 6: 142-5, 1962. 130
- NEWSOME J R and KEITH C H : Variation of the gas phase composition within a burning cigarette. Tobacco Sci 9: 65-9, 1965. 131
- OSBORNE J S, ADAMEK S and HOBBS M E : Some components of gas phase of cigarette smoke. Anal Chem 28: 211-5, 1956. 132
- OTSUKA S, FUJIWARA K, IKAWA H and HIRAYAMA K : Studies on carbon monoxide content in cigarette smoke. J Hyg Chem Japan 16: 150-3, 1970. 133
- OWEN W C and REYNOLDS M L : The diffusion of gases through cigarette paper during smoking. Tobacco Sci 16: 28-34, 1967. 134
- PHILIPPE R J and HOBBS M E : Some components of the gas phase of cigarette smoke. Anal Chem 28/12: 2002-6, 1956. 135
- SARUTA, N : Uber die Quantitative Bestimmung der Kohlenoxydmenge im Tabakrauche. Fuk Acta Med 20: 117-8, 1937. 136
- SCASSELLATTI /SFORZOLINI G and SAVINO A : Valutazione di un indice rapido di contaminazione ambientale da fumo di sigaretta, in relazione alla composizione della fase gassosa del fumo. (Evaluation of a rapid index of environmental pollution by cigarette smoke in relation to the composition of the gas phase of the smoke). Riv Ita Ig 28: 43-55, 1968. 137
- STENHAGEN E : On the chemistry of tobacco smoke. Acta Soc Med Upsal 64/3-4: 322-40, 1959. 138
- TOTH J : Beitrag zur Kohlenoxydbestimmung im Tabakrauche. Chem Zfg 31: 98-9, 1907. 139
- WAHL F : Ueber den Gehalt des Tabakruaches and Kohlenoxyd. Pflugers Arch 78: 262-85, 1899. 140
- WALTZ P and HAUSERMANN M : Betrachtungen uber die Veranderung des Tabakrauches in der Cigarette. (Reflections on the change in tobacco smoke in the cigarette. The yield of crude condensate, total moisture, pyridine nicotine, phenols, pyrocatechol, scopoletin, and carbon monoxide in cigarette smoke as a function of the puff number and of the smoke filter. Beitr Tabakforsch 3(3):169-202, 1965. 141

1005051094

### III. SOURCES OF CARBON MONOXIDE

The interpretation of the significance of carboxyhemoglobin levels in the blood included the contribution from exogenous carbon monoxide in the atmosphere, as well as endogenous carbon monoxide that is found in the tissues. The details of the contribution of these forms of carbon monoxide are discussed in the articles in additional bibliography list no. 5, dealing with atmospheric carbon monoxide, and additional bibliography list no. 6 on endogenous carbon monoxide. Pollution of the air by factories and by vehicular exhausts are covered respectively in additional bibliography lists nos. 7 and 8. This section will discuss those articles that include analysis of carboxyhemoglobin in the blood of individuals that have been exposed to the sources of carbon monoxide other than cigarette smoking.

#### A. Carboxyhemoglobin Levels in the Blood of Nonsmokers

It was stated in section II-A that the carboxyhemoglobin contained in the blood of habitual smokers included the contribution from exogenous and endogenous sources of carbon monoxide. Table 4 summarizes the results of 26 investigations of blood levels in nonsmokers. The overall mean for 1,662 subjects was 1.45 %. The individual averages varied from 0.4 to 5.2. In cities such as London, Los Angeles and Milan, the residents show the following mean levels of carboxyhemoglobin-respectively: 3.5, 2.3 and 2.8. These values represent a significant contribution from carbon monoxide pollution in the atmosphere.

(Table 4 appears on the next page.)

1005051095



Table 4. Carboxyhemoglobin levels in the blood of nonsmokers.<sup>1</sup>

page 33

Reference (Year)	Nature of nonsmoker	No of Subjects	Carboxyhemoglobin blood levels % Mean $\pm$ SD (Range)	
Conson and Hastings (1953)	Chicago residents	7	1.5	(1.2-1.9)
Schmidt (1939)	Bonn residents	14	0.6	(0.2-1.1)
Meigs (1948)	US army enlisted men	4	5.2	(0-8.8)
Parmeggiani and Gilardi (1952)	Milan residents	9	2.8	(0-4)
Barthe, Paris, Duchemin and Thomas (1953)	Paris residents	30		(0.6-1.2)
Ruel and Barthe (1954)	Paris residents		0.5	
Valic and Duric (1954)	Yugoslavian residents	75	0.9 $\pm$ 0.42	
Gaensler, Cadigan, Ellicott, Jones and Mark (1957)	Boston residents	9	0.9	(0.6-1.2)
Dahlström, Nordström- Öhrberg and Rothschild (1958)	Stockholm residents	14	0.7	(0.4-0.8)
Whitehead and Worthington (1961)	Warwick residents	6	0.8	(0. -1.5)
Trinder and Harper (1962)	Sunderland residents (adults)	23	0.9 $\pm$ 0.3	(0.4-2.1)
	(children)	24	0.6 $\pm$ 0.2	(0.3-1.0)
Hofreuter, Catcott and Xintaras (1962)	Cincinnati residents	6	1.9	
Goldsmith, Schuette and Novick (1963)	San Francisco longshoremen	764	1.3	
Goldsmith, Terzaghi and Hackney (1963)	Los Angeles resident	1	2.3	
Bowden and Woodall (1964)	Halton residents	42	0.7 $\pm$ 0.3	(0.2-1.5)
Curphrey, Hood and Perkins (1965)	Los Angeles longshoremen	115	0.4	
Ayres, Giannelli and Armstrong (1965)	New York residents	28	0.9	(0.1-1.9)
Ibo, Marucci and Ronchi (1966)	Paris residents	44	1.9	(0-7)
McIlvaine, Nelson and Bartlett (1969)	Durham residents	10	1.6	
Lawther and Commins (1970)	London residents	165	3.5	

1005051096

Table 4. (continued)

page 33 a

Reference (Year)	Nature of nonsmoker	No. of Subjects	Carboxyhemoglobin blood levels % Mean $\pm$ SD (Range)	
Lacoub, Faure, Mallion and Cau (1970)	Paris residents	113	1.5	
Rouch, Rioufol and Bourbon (1971)	Toulouse residents	23	1.6	(1.0-3.5)
Weiss, Slawsky and Desforbes (1971)	Boston residents	13	1.5 $\pm$ 0.8	
Motley (1971)	Los Angeles residents	115	1.5	(0.2-2.9)
Hansen, Wilke, Malorny and Göthert (1972)	Hamburg residents	37	0.75	
Arturson, Garby, Robert and Zaar (1972)	Uppsala residents	11	1.3	(0.9-1.7)
Mean (overall for number of subjects)			1.45	(1662 subjects)

<sup>1</sup> Some of the values were reported in volumes % and are expressed in this table as saturation % assuming normal hemoglobin values.

1005051097

## BIBLIOGRAPHY

## III. SOURCES OF CARBON MONOXIDE

A. Carboxyhemoglobin Levels in the Blood of Non-smokers.

- |  | Reprint |
|--|---------|
| ARTURSON G, GARBY L, ROBERT M and ZAAR B : Changes in carbon monoxide content of whole blood during gas equilibration in the radiometer dissociation curve analyzer. (DCA-1). <u>Upsala J Med Sci</u> 77: 22-4, 1972.  | 142     |
| AYERS S M, GIANNELLI S and ARMSTRONG R G : Carboxyhemoglobin: Hemodynamic and respiratory responses to small concentrations. <u>Science</u> 149: 193-4, 1965.  | 143     |
| BALBO W, MARUCCI V and RONCHI G U : Les valeurs de l'oxycarbonemie en differentes conditions individuelles et de milieu Recherches preliminaires. (Values of blood carbon monoxide under various individual and environmental conditions. Preliminary studies). <u>Acta Med Leg Soc Liege</u> 19: 187-9, 1966. | 144     |
| BARTHE R, PARIS J, DUCHEMIN M and THOMAS F : Valeur et interpretation des taux d'oxycarbonemie obtenus par la methode Nicloux-Endiometrie. Deuxieme note. - Oxy-carbonemie et intoxications exogenes: L'oxycarbonemie des fumeurs et des ethyliques. <u>Arch Mal Profes(Paris)</u> 14: 288-90, 1953.           | 145     |
| BOWDEN C H and WOODHALL W R : The determination and significance of low blood carboxy-haemoglobin levels. <u>Med Sci Law</u> 4: 98-107, 1964.  | 146     |
| CURPHEY T J, HOOD L P and PERKINS N M : Carboxyhemoglobin in relation to air pollution and smoking. <u>Arch Environ Health</u> 10: 179-85, 1965.   | 147     |
| DAHLSTRÖM H, NORDSTRÖM-ÖHRBERG G and ROTHSCILD A : The influence of tobacco smoking and increased initial carbon monoxide concentration on results of Sjostrand's method of total hemoglobin determination. <u>Acta Physiol Scand</u> 42: 174-84, 1958.  | 148     |
| GAENSLER E A, CADIGAN J B, ELLICOTT M F, JONES R H and MARKS A : A new method for rapid precise determination of carbon monoxide in blood. <u>J Lab Clin Med</u> 49: 945-57, 1957.   | 149     |
| GOLDSMITH J R, SCHUETTE F and NOVICK L : Appraisal of carbon monoxide exposure from analysis of expired air. <u>Ex Med Int Cong Ser</u> 62: 948-52, 1963.  | 150     |
| and HACKNEY J D:<br>GOLDSMITH J R, TERZAGHI J : Evaluation of fluctuating carbon monoxide exposures. <u>Arch Environ Health</u> 7: 647-63, 1963.   | 151     |
| HANSON H B and HASTINGS A B : The effect of smoking on the carbon monoxide content of blood. <u>JAMA</u> 100: 1481, 1933.  | 152     |
| HANSEN O, WILKE H, MALORNY G and GOTHERT M : Absorption and release of carbon monoxide during breathing of low CO concentrations by smokers and non-smokers. <u>Chem Abstr</u> 77/6: 43-6, 1972.   | 153     |
| HOFREUTER D H, CATCOTT E J and XINTARAS C : Carboxyhemoglobin in men exposed to carbon monoxide. <u>Arch Environ Health</u> 4: 81-5, 1962.   | 154     |
| LAWTHER P J and COMMINS B T : Cigarette smoking and exposure to carbon monoxide. <u>Ann NY Acad Sci</u> 174: 135-47, 1970.   | 155     |
| McILVAINE P M, NELSON W C and BARTLETT D : Temporal variation of carboxyhemoglobin concentrations. <u>Arch Environ Health</u> 19: 83-91, 1969.   | 156     |
| MEIGS J W : Carbon monoxide poisoning. <u>Bull US Army</u> 8: 542-6, 1948.   | 157     |

1005051098

Bibliography III. A

- MOTLEY H L: Environmental air pollution. Effect on pulmonary function. Aerospace Med 42: 1108-10, 1971. 158
- PARMEGGIANI L and GILARDI F: Rilievi sulla ossicarbonemia fisiologica. Med Lav 43: 179-83, 1952. 159
- ROUCH Y, RIOUFOL F and BOURBON P: Oxycarbonemie spontanee et apres exposition pendant quelques minutes a une atmosphere a un pour mille d'oxyde de carbone chez 43 sujets. (Carbon monoxide content in blood: normal and after a few minutes exposure to an atmosphere containing 0. 1p. 100 carbon monoxide/43 subjects). Arch Mal Prof (Paris) 32: 271-82, 1971. 160
- RUEL M H and BARTHE M R: Examen systematique de 132 ouvriers d'usines a gaz avec dosage du CO sanguin. Arch Mal Prof Paris 15: 234-5, 1954. 161
- SCHMIDT O: Der gasanalytische nachweis von Kohlenoxyd im Blut, insbesondere bei Rauchern. Klin Wschr 18: 938, 1939. 162
- TRINDER P and HARPER F E: A colorimetric method for the determination of carboxyhaemoglobin over a wide range of concentrations. J Clin Path 15: 82-4, 1962. 163
- VALIC F and DURIC D: Concentration of carbon monoxide in the blood of smokers and non-smokers. Arh Hig Rada (Jugoslavia) 5: 49-56, 1954. 164
- WEISS E B, SLAWSKY P and DESFORGES J F: Oxyhemoglobin affinity in chronic pulmonary granulomatosis (sarcoidosis) and fibrosis. Am Rev Res Dis 104: 694-702, 1971. 165
- WHITEHEAD T P and WORTHINGTON S: The determination of carboxyhaemoglobin. Clin Chim Acta 6: 356-9, 1961. 166
- YACOB M, FAURE J, MALLION J M and CAU J: Peut-on determiner l'origine tabagique de l'oxycarbonisme chronique? (Is it possible to demonstrate chronic carbon monoxide intoxication specifically caused by smoking? Med Leg Domm Corpor Paris 3: 262-5, 1970. 167

1005051099

### III B. Vehicular Traffic and Smoking

One major source of air pollution in general and of carbon monoxide in particular is automobile exhaust. Blood samples collected from vehicular drivers, traffic policemen, garage operators and miscellaneous workers show an increase in content of carboxyhemoglobin after exposure. The results of 19 investigations summarized in Table 5 indicate an overall average in 84 subjects of an increase of 1.1% of carboxyhemoglobin following 4 to 8 hours of exposure to vehicular traffic. The average peak level for 1,481 subjects was 3.9%.

The combination of exposure to vehicular traffic and cigarette smoking has been investigated by 8 groups and is summarized in Table 6. The peak blood levels showed an overall average of 6.8% of carboxyhemoglobin for 446 subjects. There is an increase of 2.9% in carboxyhemoglobin as a result of smoking and this value is close to the increase of 3.8% reported for subjects not exposed to vehicular traffic (see section IIB).

The most extreme example of combining cigarette smoking and exposure to automobile exhaust was reported by Srch (1967), who placed 4 subjects in an automobile with doors and windows closed inside a closed garage. Two of the subjects were smokers of 20 to 30 cigarettes daily, who were asked to smoke 5 cigarettes. Before smoking, the blood showed 5% of carboxyhemoglobin. After smoking 5 cigarettes, the blood levels had a mean value of 10%. The two nonsmokers in the automobile had 2% of carboxyhemoglobin before and 5% after the test. This situation rarely

1005051100

occurs in the ordinary course of events.

There is no reason to suspect that the blood levels of carboxyhemoglobin attained by the combination of cigarette smoking and exposure to vehicular traffic are responsible for the occurrence of vehicular accidents and poor driving (Boeck, 1958; Williams, 1964). The mechanical properties of the lung do not show any abnormality attributable to the exposure to cigarette smoking and vehicular traffic (Reichel et al, 1970). The incidence of sick rates among smokers was significantly higher than among nonsmokers who were exposed to carbon monoxide in the atmosphere (Fiandaca and Vercellotti, 1964; Mountain et al, 1968). There is a reduction in reaction time in drivers exposed to 90 minutes of commuting traffic (Ramsey, 1970). In all of these reports, although an elevation of carboxyhemoglobin is mentioned, a cause and effect relationship has not been established. The exposure to vehicular traffic is associated with inhalation of lead, ozone, hydrocarbons and nitrogen oxides, each of which can also contribute to the effect attributed to carbon monoxide alone. Literature on the toxicity of automobile exhaust is listed as additional bibliography No. 8.

1005051101

- PARKHURST and GIBSON 1967 (A 131) 184  
 PARMEGGIANI and GILARDI 1952 (53) 18, 13, 14;  
 (83) 22, 21; (159) 35, 33  
 PARROT, STUPFEL, ROMARY and MORDELET-  
 DAMBRINE 1971 (224) 50, 47  
 PASECHNIK, SHTUMM, VLADISLAVLEV and  
 ZAMAYATNIN 1971 (A 133) 184  
 PATTONO, MARCHIARO, CAPELLARO and  
 ORIONE 1964 (A 160) 186  
 PATZ 1949 (261) 55, 51  
 PAULEIKOFF, MULLER-FAHBUSCH, MESTER  
 and MEIBNER 1971 (A 521) 209  
 PAULET and CHEVRIER 1966 (A 134) 184  
 PAULET and CHEVRIER 1969 (A 765) 227  
 PAULI, TRUNIGER, LARSEN and MULHAUSEN  
 1968a (557) 101, 99  
 PAULI, TRUNIGER, LARSEN and MULHAUSEN  
 1968b (558) 101, 100  
 PEARCE 1968 (A 522) 209  
 PECORA 1964 (A 766) 227  
 PEDRERO and RODRIGO 1964 (A 347) 198  
 PERRELLI, PREVOT and SULOTTO 1970 (418)  
 77, 74  
 PERRELLI and ROSETTANI 1964 (A 297) 195  
 PERRELLI, ROSETTANI and BRAGUZZI 1965  
 (A 278) 195  
 PETERSON, SIGGAARD-ANDERSEN,  
 KRISTENSEN and KJELDSEN 1968 (348) 67,  
 66  
 PETERSON and STEWART 1970 (A 161) 186  
 PETIT, PETIT and GEILLE 1970 (A 692) 220  
 PETRILLI and KANITZ 1970 (186) 41, 38, 39  
 PETROVA, DALAKMANSKI and BAKALOV  
 1966 (A 299) 195  
 PETROVIC 1970 (419) 77, 75  
 PETTER, BOURBON, MALTIER and JOST  
 1971 (A 135) 184  
 PETTY 1969 (A 832) 231  
 PHELPS and ANTONINI 1969 (A 136) 184  
 PHILIPPE and HOBBS 1956 (135) 31, 29  
 PLACE 1979 (A 767) 227  
 PICKWELL 1970 (A 195) 189  
 PIEDELIEVRE, BRETON and DEROBERT  
 1969 (591) 106, 102  
 PIERCE and COLLINS 1971 (A 343) 198  
 PINCHERLE and SHANKS 1967 (420) 77, 75  
 PIPER, PFEIFER and SCHEID 1969 (A 433) 204  
 PIPER and SIKAND 1966 (A 432) 204  
 PIRNAY, DEROANNE, DUJARDIN and PETIT  
 1971b (299) 60, 56  
 PIRNAY, DUJARDIN, DEROANNE and PETIT  
 1971a (298) 60, 56  
 PIRNAY, FASSOTTE, DEROANNE and PETIT  
 1968 (A 137) 184  
 PIRNAY, FASSOTTE, GAZON, DEROANNE  
 and PETIT 1969 (A 434) 204  
 PIRNAY, PETIT and ROBERTS 1970 (A 435)  
 204  
 PODLESCH and STEVANOVIC 1966 (A 436) 204  
 POGRUND 1969 (609) 110, 108  
 POLITZER 1968 (A 138) 184  
 POLLARD 1970 (A 693) 220  
 PORTHEINE 1971 (105) 26, 25  
 POWER 1968 (A 162) 186  
 POWER, AOKI, LAWSON and GREGG 1971  
 (A 438) 204  
 POWER, HYDE, SEVER, HOPPIN and NAIRN  
 1965 (A 437) 204  
 PRELLWITZ, SCHUSTER, SCHYLLA, BAUM,  
 SCHONBORN, UNGERN-STERMBERG,  
 BRODERSEN and POEPLAU 1970 (421) 77, 74;  
 (A 638) 217  
 PREROVSKA and DRDKOVA 1967a (384) 73, 70  
 PREROVSKA and DRDKOVA 1967b (385) 73, 70  
 PREROVSKA and DRDKOVA 1971 (386) 73, 71  
 PREZIOSI, LINDENBERG, LEVY and  
 CHRISTENSON 1970 (521) 95, 93  
 PROKOP and WABNITZ 1970 (453) 83, 81  
 PUKHOV 1964 (610) 110, 108  
 PUKHOV 1965 (A 768) 227  
 PUREC and KRASNA 1967 (A 605) 215  
 QUINTANA, MIRETE and GARCIA 1969 (559)  
 101, 99  
 RADUSHVICH 1968 (A 769) 227  
 RAMSEY 1972 (84) 22, 20  
 RAMSEY 1966 (A 300) 195  
 RAMSEY 1967 (A 344) 198  
 RAMSEY 1969 (422) 77, 75  
 RAMSEY 1970 (188) 41, 37  
 RAMSEY 1972 (493) 91, 89  
 RAMSEY 1967 (187) 41, 38, 39  
 RANDOWA 1967 (A 440) 204  
 RANDOWA and SIERAWSKI 1964 (A 439) 204  
 RAPOPORT 1967 (A 770) 227  
 RAUSA, DIANA and PERIN 1968 (A 139) 184  
 RAUSA, PERIN and DIANA 1967 (A 300) 195  
 RAY and ROCKWELL 1970 (494) 91, 89  
 RAY 1967 (A 606) 215  
 RAYFIELD 1967 (A 301) 195  
 RECKZEH and DONTENWILL 1970 (423) 77, 75  
 RECKZEH, RUCKER, HARKE and DONTENWILL  
 1969 (106) 26, 25  
 REDDEMANN, AMENDT and JAHRIG 1970  
 (A 639) 217  
 REED and TROTT 1971 (A 302) 195  
 REED 1970 (A 140) 184  
 RIECHEL, WOBITH and ULMER 1970 (189) 41, 37  
 REJSEK 1971 (A 771) 227  
 REMMERS and MITHOEFER 1969 (A 441) 204  
 REPLOH, KLOSTERKOTTER and  
 EINK-ROSSKAMP 1966 (A 710) 223  
 REVOL, MONIER, COURJON, FOURMET and  
 GERIN 1966 (543) 97, 96  
 REVSIN and BRODIE 1969 (A 607) 215  
 RHODES 1971 (225) 50, 48  
 RICCI, CAPELLARO and GAIDO 1964 (424) 77, 74  
 RI 1966 (630) 112, 111  
 RIKANS and VAN DYKE 1971 (A 577) 213  
 RINGEL and KLAUWANS 1972 (A 523) 209  
 RINGOLD, GOLDSMITH, HELWIG, FINN and  
 SCHUETTE 1962 (85) 23, 19

1005051309

Table 6. Carboxyhemoglobin blood levels of smokers following exposure to vehicular traffic.

Reference (Year)	Nature of subjects	No of subjects	Carboxyhemoglobin blood levels %				Net $\Delta$ compared Nonsmokers
			Before Mean	After smoking and exposure Mean $\pm$ SD	(Range)	$\Delta$	
DeBruin, Bult and Van Haeringen (1965)	Amsterdam police- men	14	4.6	4.9		+0.3	+2.9
Chovin (1967)	Paris policemen	35	3.7				+2.5
Ramsey (1967)	Dayton parking attendants	24	2.9	9.3 $\pm$ 3.16		+6.4	+2.0
Buchwald (1969)	Alberta garage operators (1-9/day)	21		6.4	(1-15)		+1.4
	(10-20/day)	138		8.5	(0.5-19)		+3.5
	(>20/day)	76		9.2	(1-18)		+9.2
Breyse and Bovce (1969)	Seattle fork lift drivers ( $\leq$ 20/day)	44		3.5	(0-17)		+2.3
	(20-40/day)	38		5.5	(1-20)		+4.3
	(>40/day)	6		6.5	(3-14)		+5.3
Gothe, Fristedt, Sundell, Kolmodin, Ehrner-Samuel and Gothe(1969)	Stockholm policemen	28		3.5 $\pm$ 1.17			+2.3
	Malmö policemen	6		5.0 $\pm$ 2.4			+4.2
	Orebro policemen	5		2.4 $\pm$ 1.1			+1.8
Petrilli and Kanitz (1970)	Genoa vehicle drivers	20			(6.0-8.5)		
Cohen, Dorion, Golds- mith, Permutt (1971)	US-Mexican border inspector	11	4.8	6.4		+1.6	+2.8
Mean (overall for number of subjects)				6.8 (446 subjects)			+4.07 (446 subjects)

1005051103



## III. SOURCES OF CARBON MONOXIDE

## B. Vehicular Traffic and Smoking

- ALIVISATOS G P, BAZAS B N, ALEXOPOULOS J and VERYKOKAKIS E : Air pollution in the city of Athens and surrounding territory. Ig Mod 60/1-2: 3-25, 1967. 168 Reprint
- AYRES S M and BUEHLER M E : The effects of urban air pollution on health. Clin Pharm Ther 11: 337-71, 1970. 169
- BOECK J K : Traffic safety. Research Review 2: 2- , 1958. 170
- BREYSSE P A and BOVEE H H : Use of expired air-carbon monoxide for carboxyhemoglobin determinations in evaluating carbon monoxide exposures resulting from the operation of gasoline fork lift trucks in holds of ships. Am Industr Hyg Ass J 30: 477-83, 1969. 171
- BUCHWALD H : Exposure of garage and service station operatives to carbon monoxide: a survey based on carboxyhemoglobin levels. Am Industr Hyg Ass J 30: 570-5, 1969. 172
- CHOVIN P : Carbon monoxide: analysis of exhaust gas investigations in Paris. Environ Res 1: 198-216, 1967. 173
- COHEN S I, DORION G, GOLDSMITH J R and PERMUTT S : Carbon monoxide uptake by inspectors at a United States-Mexico border station. Arch Environ Health 22: 47-54, 1971. 174
- DeBRUIN A : Carboxyhemoglobin levels due to traffic exhaust. Arch Environ Health 15: 384-9, 1967. 175
- BULT A R and VAN  
DeBRUIN A / HAERINGEN A : Het koolmonoxydegehalte in het bloed bij verkeersagenten. (The carbon monoxide content of the blood in traffic police). T Soc Geneesk 43: 775-7, 1965. 176
- VAN  
DeBRUIN A, VROEGE D and HAERINGEN A : Onderzoek naar de opname van koolmonoxyde bij verkeersagenten. (Study of carbon monoxide uptake in traffic policemen). T Soc Geneesk 43: 146-51, 1965. 177
- DESOILLE H : Une pseudo-intoxication-collective chronique par l'oxyde de carbone dans un garage. (A chronic collective pseudo-intoxication by carbon monoxide in a garage), Arch Mal Profes Paris 28(4/5): 472, 1967. 178
- FIANDACA S and VERCELLOTTI E : Aspetti della morbidita e della infortunabilita nei soggetti esposti a rischio di ossicarbonismo cronico. (Aspects of the morbidity and accident rates in subjects exposed to the risk of chronic carbon monoxide poisoning). Rass Med Industr 33: 360-79, 1964. 179
- GOLDSMITH J R, TERZAGHI J and HACKNEY J D : Evaluation of fluctuating carbon monoxide exposures. Arch Environ Health 7: 647-63, 1963. 180
- GOTHE C J, FRISTEDT B, SUNDELL L, KOLMODIN B, EHRNER-SAMUEL H and GOTHE K : Carbon monoxide hazard in city traffic. Arch Environ Health 19: 310-4, 1969. 181
- LUDERITZ P : Experimentelle Untersuchungen uber die Wirkungen von Kohlenmonoxid aus Kraftfahrzeugabgasen auf den Gesundheitszustand von Verkehrspolizisten. (Experimental studies on the effect of carbon monoxide from motor vehicle exhaust gases on the state of health of traffic police). Z Ges Hyg 17(9): 645-6, 1971. 182
- MORANDO A E and ROVIDA S : Sull' esposizione di un campione di popolazione (Vigili urbani) all'ossidio di carbonio atmosferico derivante da traffico motorizzato. (On the exposure of a sample of the population (policemen) to atmospheric carbon monoxide deriving from motorized traffic). G Ig Med Prev 6: 78-90, 1965. 183

1005051104

- MOUNTAIN I M, CASSELL E J, WOLTER D W, MOUNTAIN J D, DIAMOND J R and McCARROLL J R: Health and the urban environment. VII. Air pollution and disease symptoms in a "normal" population. Arch Environ Health 17: 343-52, 1968. 184
- MOUREU H: Carbon monoxide as a test for air pollution in Paris due to motor-vehicle traffic. Proc Roy Soc Med 57 Suppl: 1015-20, 1964. 185
- PETRILLI F L and KANITZ S: Problemes de sante publique qui se posent a la suite du trafic motorise et de la pollution de l'air qui en resulte. (Problems of public health as a result of motor traffic and the resulting air pollution). G Ig Med Prev 11: 215-35, 1970. 186
- RAMSEY J R: Carboxyhemoglobinemia in parking garage employees. Arch Environ Health 15: 580-3, 1967. 187
- RAMSEY J M: Oxygen reduction and reaction time in hypoxic and normal drivers. Arch Environ Health 20: 597-601, 1970. 188
- REICHEL G, WOBITH F and ULMER W T: Akute und chronische Wirkung von Straßenluft an verkehrsreicher Kreuzung auf die Lungenfunktion des Menschen, den CO-, Hb- und Bleigehalt des Blutes. (Acute and chronic effects of air pollution produced by traffic at a busy crossing on lung function in humans. Determination of CO, Hb and Pb in blood. Int Arch Arbeitsmed 26: 84-97, 1970. 189
- SRCH M: Über die Bedeutung des Kohlenoxyds beim Zigarettenrauchen im Personenkraftwageninneren. (On the significance of carbon monoxide in cigarette smoking in an automobile). Deutsch Z Ges Gerichtl Med 60: 80-9, 1967. 190
- SZADKOWSKI D, MASTALL V, SCHALLER K H and LEHNERT G: Pilot-study zur beruflichen Kohlenmonoxid-Gefährdung in Großstadtstraßen. (Pilot study on occupational exposure to carbon monoxide in city streets). Int Arch Arbeitsmed 26: 224-30, 1970. 191
- WILLIAMS N: Traffic accidents - epidemiology and medical aspects of prevention. Canad Med Ass J 90: 1099-1104, 1964. 192

1005051105

- (26) DOYLE J T : Smoking and myocardial infarction. Circulation 39 & 40: Suppl 4: 136-43, 1969.

This review contains a paragraph on the role of carbon monoxide in pathogenesis of atherosclerosis. The author views the problem in the proper perspective.

The manner in which cigarette smoking accelerates atherosclerosis and its complications is, in short, unexplained. It is possible that in some way cigarette smoking damages the arterial intima. Carbon monoxide is the likeliest immediate candidate for such a role. Some presently mysterious interference with the normal mechanism of transport of lipids from the plasma through the vascular tunics to the lymphatics secondary to the inhalation of cigarette smoke is an alternative possibility: in all populations yet scrutinized, the prevalence and incidence of CHD rise with the serum cholesterol concentration.<sup>22</sup> It is, accordingly, a plausible hypothesis that inordinate cigarette smoking may be associated with an increased serum cholesterol concentration. Such a relationship does, indeed, exist, but is unimpressive. Although the serum cholesterol concentration in both men and women is consistently higher in cigarette smokers, the influence of increasing age is substantially greater (figs. 1 and 2).<sup>23</sup> The observation that heavy cigarette smokers have far more atheroma than nonsmokers is, possibly, complemented by Astrup's observation that fat-fed rabbits exposed to high tensions of carbon monoxide exhibit extreme hyperlipidemia and cholesterosis as compared to controls not exposed to carbon monoxide.<sup>24-26</sup> This interesting experimental model has, however, no recognized counterpart in human epidemiological studies. Obesity as a coronary risk factor is not related to cigarette smoking.<sup>2</sup> Lastly, the arterial blood pressure is not associated with cigarette habit.<sup>2</sup>

1005051226

Habitual smokers who also suffer from chronic lung disease are known to have an elevated carboxyhemoglobin level. Such patients show a reduction in pulmonary diffusion capacity (Chosy et al., 1963; Clauzel et al., 1966; Trinquet et al., 1971). Chevalier et al. (1966) compared such patients with nonsmokers inhaling 0.5 % carbon monoxide. The resulting elevation in levels of carboxyhemoglobin caused a reduction in pulmonary diffusing capacity in nonsmokers. On this basis, it was proposed that the reduction in pulmonary diffusion capacity in smokers is due to elevation of the carboxyhemoglobin level. This explanation has not been supported by any other form of experimental data.

1005051107

## IV. RESPIRATORY SYSTEM

A. Acute Respiratory Effects in Humans

- AYRES S M, GIANNELLI S Jr and ARMSTRONG R G : Carboxyhemoglobin: Hemodynamic and respiratory responses to small concentrations. Science 149: 193-4, 1965. Reprint 193
- CHEVALIER R B, KRUMHOLZ A and ROSS J G : Reaction of nonsmokers to carbon monoxide inhalation. JAMA 198: 1061-4, 1966. 194
- CHIODI H, DILL D B, CONSOLAZIO F and HORVATH S M : Respiratory and circulatory responses to acute carbon monoxide poisoning. Am J Physiol 134: 683-93, 1941. 195
- CHOSY L, GEE J B L and RANKIN I : The effects of cigarette smoking on the pulmonary diffusing capacity. Clin Res 11: pt. 3: 301, 1963. 196
- CLAUZEL A M, TRINQUET G, CARRE R and MEYER A : Influence du tabac sur la ventilation et la diffusion pulmonaire des sujets normaux. Etude statistique de 194 sujets. (Influence of tobacco on ventilation and pulmonary diffusion of normal subjects. Statistical study of 194 subjects). Rev Med Aeron 5(20): 13-6, 1966. 197
- FISHER A B, HYDE R W, BAUE A E, REIF J S and KELLY D F : Effect of carbon monoxide on function and structure of the lung. J Appl Physiol 26: 4-12, 1969. 198
- TRINQUET G and MEYER A : Role de l'oxyde de carbone dans le transfert de l'oxygene au cours des bronchopathies obstructives. (The role of carbon monoxide in the transfer of oxygen during obstructive bronchopathies). Poumon Coeur 27: 695-704, 1971. 199

1005051108

#### IV B. Chronic Respiratory Effects in Humans

The medical examination of traffic officers stationed at the Holland Tunnell in New York has provided an opportunity to determine the effects of chronic exposure to 70 ppm carbon monoxide. Sievers et al. (1942) examined 156 such officers and failed to find any evidence of injury to health that was attributable to carbon monoxide exposure. There were no signs or symptoms of respiratory abnormalities. Other reports of elevated carbon monoxide levels in highway tunnels have appeared, but the clinical examination of the traffic officers has not been included (Braja and Trompeo, 1964; D'Arca et al. 1964; Miranda et al., 1967; Yamate and Matsumura, 1968).

Astrup et al. (1968) and Klausen et al. (1968) exposed 8 male subjects to inhalation of 0.5 % carboxyhemoglobin, resulting in a blood level of 10 %. There were no changes in ventilation, circulation or metabolism.

1005051109

## BIBLIOGRAPHY

## IV. RESPIRATORY SYSTEM

B. Chronic Respiratory Effects in Humans

- |  | Reprint |
|--|---------|
| ASTRUP P, PAULI H G, KJELDSSEN K and PETERSEN C E : Introduction and general description of the study and of the procedures for prolonged exposure to carbon monoxide and hypoxia. <u>Scan J Clin Lab Invest</u> 22: Suppl 103: 5-8, 1968.   | 200     |
| BRAJA M and TROMPEO C : Contributo alla conoscenza delle concentrazioni di ossido di carbonio nelle gallerie stradali. (Contribution to the knowledge of concentrations of carbon monoxide in road tunnels). <u>Rass Med Industr</u> 33: 411-3, 1964.  | 201     |
| D'ARCA S U, GUALDI G and ARCIERI G : Indagini e considerazioni sulla presenza di ossido di carbonio in alcune gallerie autostradali della città di Roma. (Findings and considerations on the presence of carbon monoxide in some highway tunnels in the city of Rome). <u>Nuovi Ann Ig Microbiol</u> 15: 200-20, 1964. | 202     |
| KLAUSEN K, RASMUSSEN B, GJELLEROD H, MADSEN H and PETERSEN E : Circulation, metabolism and ventilation during prolonged exposure to carbon monoxide and to high altitude. <u>Scand J Clin Lab Invest</u> 22: Suppl 103: 26-37, 1968.   | 203     |
| MIRANDA J M, KONOPINSKI V J and LARSEN R I : Carbon monoxide control in a high highway tunnel. <u>Arch Environ Health</u> 15: 16-25, 1967.   | 204     |
| SIEVERS R F, EDWARDS T I, MURRAY A L and SCHRENK H H : Effect of exposure to known concentrations of carbon monoxide. A study of traffic officers stationed at the Holland Tunnel for thirteen years. <u>JAMA</u> 118: 585-8, 1942.  | 205     |
| YAMATE N and MATSUMURA T : (Determination of air pollutants inside the tunnel under construction). <u>Bull Nat Inst Hyg Sci Tokyo</u> 86: 148-50, 1968.  | 206     |

1005051110

#### IV C. Acute and Chronic Respiratory Effects in Animals

The lack of respiratory stimulation during acute exposure to carbon monoxide has also been demonstrated in animals (Korner, 1965). In perfusion of the carotid body chemoreceptors there was no evidence of direct stimulation by carbon monoxide (Duke et al., 1952; Joels and Neil, 1961; Meyer et al., 1972). With extremely high levels of carboxyhemoglobin a fall in blood oxygen tension occurs, which may cause activation of chemoreceptors. However, the respiratory centers are depressed directly by the reduction in blood oxygen content, so that chemoreceptor-induced hyperpnea is not apparent (Mills and Edwards, 1968).

Carbon monoxide in a concentration of 2% provokes bronchoconstriction in the guinea pig (Parrot et al., 1971). In the cat lung, ventilation with 1 to 20% carbon monoxide in air caused bronchodilatation with a fall in pulmonary arterial pressure (Duke and Killick, 1952). In the dog there is a rise in pulmonary arterial blood pressure following inhalation of carbon monoxide (Rubino, 1964). The differences in bronchomotor and pulmonary vascular responses between animal species have not been explained. Carbon monoxide is not toxic to bronchial cilia of various animals (Okeson and Divertie, 1970).

There are other differences in responses of various animal species. In the rat lung, exposure to 0.5 or 1.0% carbon monoxide in air caused swelling of alveolar epithelial mitochondria and nucleoplasm, swelling of capillary endothelial cells and capillary platelet thrombosis (Niden and Schulz, 1965). Mice exposed to carbon monoxide showed similar alteration

1005051111



in the alveolar tissue (Bils and Romanovsky, 1967; Rhodes, 1971). Experiments on the dog indicate that administration of a mixture of 8% to 14% carbon monoxide did not cause ultrastructural changes in the lung that could be attributed to the gas per se (Norman et al, 1966; Fisher et al, 1969). The observations on primates do not include electron microscopy. So far, the only published report based on gross and histopathological examination of lungs and other organs of the cynomolgus monkey does not reveal any abnormalities resulting from continuous exposure to 19.86 and 65.46 ppm carbon monoxide for 2 years (Eckardt et al, 1972).

The chronic exposure of animals to cigarette smoke reported by Dontenwill et al, (1966); Dontenwill (1967, 1970), and by Campbell (1936) does not relate directly to the effects of carbon monoxide alone. Experiments regarding exposure to automobile exhaust reported by Vaughan et al (1969) are difficult to interpret in terms of identifying the effects of carbon monoxide. Chronic exposure to carbon monoxide in air caused an aggravation of pulmonary tuberculosis in rabbits (Kiriachko, 1966) and did not influence the rate of tumor development in rats (Shintani, 1968).

1005051112

## BIBLIOGRAPHY

page 49

## IV. RESPIRATORY SYSTEM

C. Acute and Chronic Respiratory Effects in Animals

- BILS R F and ROMANOVSKY J C : Ultrastructural alterations of alveolar tissue of mice. II. Synthetic photochemical smog. Arch Environ Health 14: 844-58, 1967. 207 Reprint
- CAMPBELL J A : The effects of exhaust gases from internal combustion engines and of tobacco smoke upon mice, with special reference to incidence of tumours of the lung. Brit J Exp Path 17: 146-58, 1936. 208
- DONTENWILL W, RECKZEH G and STADLER L : Inhalationsexperimente mit cigarettenrauch. (Inhalation experiments with cigarette smoke). Beitr Tabakforsch 3:438-48, 1966. 209
- DONTENWILL W : Inhalationsexperimente mit Tabakrauch und Nitrosaminen. (Inhalation experiments with tobacco smoke and nitrosamines). Alkylierend wirkende Verbindungen. Zweite Konferenz uber aktuelle Probleme der Tabakforschung: 51, 1967. 210
- DONTENWILL W : Experimental investigations on the effect of cigarette smoke inhalation on small laboratory animals. Inhalation Carcinogenesis. A. C. S. Symposium Series No. 18: 389-412, 1970. 211
- DUKE H N, GREEN J H and NEIL E : Carotid chemoceptor impulse activity during inhalation of carbon monoxide mixtures. J Physiol 118: 520-7, 1952. 212
- DUKE H N and KILLICK E M : Pulmonary vasomotor responses of isolated perfused cat lungs to anoxia. J Physiol 117: 303-16, 1952. 213
- ECKARDT R E, MacFARLAND H N, ALARIE Y C E and BUSEY W M : The biologic effect from long-term exposure of primates to carbon monoxide. Arch Environ Health 25: 381-7, 1972. 214
- FISHER A B, HYDE R W, BAUE A E, REIF J S and KELLY D F : Effect of carbon monoxide on function and structure of the lung. J Appl Physiol 26: 4-12, 1969. 215
- JOELS N and NEIL E : The action of high tensions of carbon monoxide on the carotid chemoreceptors. Arch Int Pharmacodyn 139: 528-34, 1962. 216
- KIRIACHKO B A : (Effect of chronic carbon monoxide intoxication on the course of pulmonary tuberculosis). Vrach Delo 4: 95-9, 1966. 217
- KORNER P I : The role of the arterial chemoreceptors and baroreceptors in the circulatory response to hypoxia of the rabbit. J Physiol 180: 279-303, 1965. 218
- MEYER J R, GROVER R F and WEIL J V : Carbon monoxide and control of ventilation. Clin Res 20: 196, 1972. 219
- MILLS E and EDWARDS M W Jr : Stimulation of aortic and carotid chemoreceptors during carbon monoxide inhalation. J Appl Physiol 25: 494-502, 1968. 220
- NIDEN A H and SCHULZ H : The ultrastructural effects of carbon monoxide inhalation on the rat lung. Virchow Arch Path Anat 339: 283-92. 1965. 221
- NORMAN J N, DOUGLAS T A and SMITH G : Respiratory and metabolic changes during carbon monoxide poisoning. J Appl Physiol 21: 848-52, 1966. 222
- OKESON G C and DIVERTIE M B : Cilia and bronchial clearance: The effects of pharmacologic agents and disease. Mayo Clin Proc 45: 361-73, 1970. 223

1005051113

- PARROT J L, STUPFEL M, ROMARY F and MORDELET-DAMBRINE M: Etude sur les polluants atmospheriques. Modification de la bronchomotricite chez le cobaye anesthesie, sous l'influence de l'oxyde de carbone et de l'hypoxie. (Study on atmospheric pollutants. Modification of bronchomotoricity in the anesthetized guinea pig under the effect of carbon monoxide and hypoxia). Bull Acad Natl Med Paris 155: 627-33, 1971. 224
- RHODES M L: The effect of carbon monoxide on mitochondrial respiratory enzymes in pulmonary tissue. Am Rev Resp Dis 103: 906-7, 1971. 225
- RUBINO G F: Modificazioni emodinamiche durante l'intossicazione acuta da monossido di carbonio. (Hemodynamic changes during acute carbon monoxide poisoning). Rass Med Industr 33: 268-74, 1964. 226
- SHINTANI K: Effects of O<sub>2</sub> or CO gas exposure on carcinogenesis. Nagoya-Shiritsu Daigaku Igakkai Zasshi 19: 465-524, 1968. 227
- VAUGHAN T R, JENNELLE L F and LEWIS T R: Long-term exposure to low levels of air pollutants. Arch Environ Health 19: 45-50, 1969. 228

1005051114

## V. CIRCULATORY SYSTEM

The investigation of circulatory effects of carbon monoxide has been more extensive than that of its effects on the respiratory system discussed in the preceding section. There has been increasing concern that chronic exposure to carbon monoxide present in cigarette smoke would lead to diseases of the heart and blood vessels and abnormalities in the composition of the blood. However, the problem has not been solved by direct experimentation relating to the carbon monoxide in cigarette smoke. There are numerous observations regarding the effects of sublethal concentrations of carbon monoxide in man and animals. These are reviewed in the following paragraphs, although they are only remotely related to the small amount of carbon monoxide contained in cigarette smoke.

### V A. Heart Rate

The acceleration of heart rate known to occur during inhalation of cigarette smoke is explained by the nicotine content. The amount of carbon monoxide in the smoke does not influence heart rate, since experiments consisting of administering carbon monoxide alone in amounts even exceeding that produced by cigarettes failed to alter the electrocardiogram.

The electrocardiograms of patients suffering from acute carbon monoxide poisoning or chronic exposure to carbon monoxide show the following alterations: depression of S-T segment (Steinmann, 1937; Störmer, 1938; Wendt, 1941; Graybiel, 1942; Breu, 1943; Patz, 1949; Saracoglu, 1951); sinus arrhythmia (Breu, 1942); premature systole (Parade and Franke, 1939); atrial flutter

1005051115

(Donatelli, 1940; atrial fibrillation (Dvorak et al., 1951; Cosby and Bergeron, 1963); distortion of QRS-T complex (Hegglin, 1944; Kostyukova, 1951; Lorente and Varela de Seijas, 1953) Wolff-Parkinson-White syndrome (Doumer and Merlen, 1946; Selig, 1966); atrioventricular block (Ehrich et al., 1944; Altmann, 1953; Kledecki and Winiarski, 1963); bundle-branch block (Graziani et al., 1957); and cardiac arrest (Brunner, 1939). Multiple abnormalities in the electrogram have been reported (Faivre et al., 1954; Faivre et al., 1959; Capellaro and Gandolfo, 1964; Hayes and Hall, 1964; Medvedowsky et al., 1965; Shafer et al., 1965; Zanardi et al., 1966; Kuroiwa et al., 1968; Orinius, 1968; Lang et al., 1969; Mosinger et al., 1969; Lustman and Geerts, 1971; Thiels et al., 1972). Most of these abnormalities have been reproduced in animals inhaling concentrations larger than 2% carbon monoxide in air (von Bergmann, 1934; Motta, 1940; Loeper et al., 1942; Lewey and Drabkin, 1944; Hundt and Grünberg, 1960; Mainardi, 1964; Datsenko, 1966; Mosinger et al., 1969)

1005051116

## BIBLIOGRAPHY

page 53

## V. CIRCULATORY SYSTEM

## A. Heart Rate

- ALTMANN R : Elektrokardiographische Beobachtungen beim Kohlenoxydvergifteten. (Electrocardiographic observations in patients with CO-poisoning). Zeit Ges Inn Med 8:39-43, 1953. Reprint 229
- BREU W : Elektrokardiographische Beiträge und vergleichende Betrachtungen akuter Vergiftungen unter besonderer Berücksichtigung der Kohlenmonoxyd-Vergiftung. (Electrocardiographic studies and comparative considerations of acute poisoning, with special reference to carbon monoxide poisoning). Arch Kreislaufforsch 11: 107-35, 1942. 230
- BREU W : Der Kohlenoxydhamoglobingehalt des Blutes bei Kohlenmonoxydvergiftungen und dessen Einfluß auf das Herz, gesehen im Elektrokardiogramm. (The carbon monoxide-hemoglobin content of the blood in carbon monoxide poisoning and its influence on the heart as seen in the electrocardiogram). Wein Klin Wochsch 56: 103-6, 1943. 231
- BRUNNER J : Ein Fall schwerster CO-Vergiftung durch andauernde Wiederbelebungsversuche gerettet. (A case of severe carbon monoxide poisoning successfully counteracted. Wein Klin Wochsch 52: 1116-7, 1939. 232
- CAPELLARO FRANCO and GANDOLFO SEBASTIANO : Indagini elettrocardiografiche in un gruppo di operai esposti al CO. (Electrocardiographic investigations of a group of workers exposed to CO). Rass Med Industr 33: 443-5, 1964. 233
- COSBY R S and BERGERON M : Electrocardiographic changes in carbon monoxide poisoning. Am J Cardiol 11: 93-6, 1963. 234
- DATSENKO I I : (Changes in cardiac activity in experimental chronic carbon monoxide poisoning). Gig Sanit 31: 102-5, 1966. 235
- DONATELLI L : L'azione cardiaca di alcuni derivati dell'AC. Barbiturico usati come narcotici. (Cardiac effect of some barbiturates and used as narcotics. Boll Soc Ital Biol Sper 15: 839-40, 1940. 236
- DOUMER E and MERLEN J F : Syndrome de Wolff, Parkinson et White au cours d'une intoxication aiguë par l'oxyde de carbone. (Wolff-Parkinson-White syndrome in the course of an acute intoxication by carbon monoxide. Acta Cardiol 1: 302-4, 1946. 237
- DVORAK L, PROKSAN F and ZITKA M : Míhání síní ve světle nové teorie a jeho klinický výskyt. (New therapy of auricular fibrillation and its clinical aspect). Cas Lek Ces 90: 1235-8, 1951. 238
- EHRLICH W E, BELLET S and LEWEY F H : Cardiac changes from CO poisoning. Am J Med Sci 208: 511-23, 1944. 239
- FAIVRE, M M, DUREUX, VINCENT and MULLER : Alterations électrocardiographiques du type coronarien dans l'intoxication aiguë par l'oxyde de carbone. (Electrocardiographic alterations of the coronary type in acute intoxication by carbon monoxide. Rev Prat 4: 3068, 1954. 240
- FAIVRE G, GILGENKRANTZ J M and HUEBER J M : Les alterations électrocardiographiques dans l'intoxication oxycarbonée aiguë. (Electrocardiographic changes in acute carbon monoxide poisoning. Presse Med 67: 502-5, 1959. 241
- GRAYBIEL A : Diseases of the Heart. A review of significant contributions made during 1941. The heart in wartime, with special reference to neurocirculatory asthenia. Arch Intern Med 70: 303-42, 1942. 242
- GRAZIANI G, ROSSI L, CASTELLINO N and SILVERSTRONI A : Gli elettroliti nella intossicazione sperimentale da CO nei loro rapporti col quadro elettrocardiografico. (Electrolytes in experimental carbon monoxide intoxication in relation to electrocardiographic alterations. Boll Soc Ital Biol Sper 33: 1002-4, 1957. 243

100505117

## Bibliography V. A

- HAYES J M and HALL G V : The myocardial toxicity of carbon monoxide. Med J Aust 56(1): 865-8, 1964. 244
- HEGLIN R : Die verlängerte QT-Dauer im Elektrokardiogramm. (The prolonged QT period in the electrocardiogram). Arch Kreislaufforsch 13: 173-286, 1944. 245
- HUNDT H J and GRUNBERG H : Vergleichende Untersuchungen über das Ratten-Ekg bei Sauerstoffmangelzuständen verschiedener Genese. (Comparative study of electrocardiograms in rats with hypoxia of varied etiology). Zeit Kreislaufforsch 49: 769-80, 1960. 246
- KLEDECKI Z and WINIARSKI W : Elektrokardiogram w ostrym zatruciu tlenkiem wegla u ludzi. (Electrocardiogram in acute poisoning with carbon monoxide). Pol Tyg Lek 18: 1101-3, 1963. 247
- KOSTYUKOVA S Z : (Electrocardiographic observations in acute intoxication with carbon monoxide). Klin Med 29: 56-60, 1951. 248
- M, CHEN C M and UEDA H  
KUROIWA A, MURAO S, HARUMI K, KATAYAMA S, YAMAMOTO/: (The electrocardiogram in acute carbon monoxide poisoning). Naika 22: 1124-9, 1968. 249
- LANG K F, SCHUSTER H P, UNGERN-STERMBERG A, BAUM P and KNOLLE J : Häufigkeit und prognostische Bedeutung von Rhythmusstörungen bei Patienten einer internistischen Intensivstation. (Frequency and prognosis of cardiac arrhythmias in patients of an intensive care unit for internal diseases). Z Kreislaufforsch 58: 1025-45, 1969. 250
- LEWEY F H and DRABKIN D L : Experimental chronic carbon monoxide poisoning of dogs. Am J Med Sci 208: 502-11, 1944. 251
- LOEPER M, VARAY A and COTTET J : Le coeur dans l'intoxication oxycarbonée aiguë. Etude clinique et expérimentale. Arch Mal Coeur 35: 123-7, 1942. 252
- VARELA D E and  
LORENTE L, SEIJAS J V : Alteraciones electrocardiograficas en la intoxicacion por el oxido de carbono. (Electrocardiographic alterations in the intoxication with carbon monoxide). Rev Clin Espan 50: 103-5, 1953. 253
- LUSTMAN F and GEERTS L : Manifestations électrocardiographiques dans 114 cas d'intoxication par l'oxyde de carbone. (Electrocardiographic manifestations in 114 cases of carbon monoxide poisoning). Acta Clin Belg 26: 131-8, 1971. 254
- MAINARDI M : Alterazioni elettrocardiografiche in casi di intossicazione acuta da ossido di carbonio. (Electrocardiographical changes in cases of acute carbon monoxide poisoning). Rass Med Industr 33: 420-9, 1964. 255
- MEDVEDOWSKY J L, SACCO J J and BELZUNCE M : Intoxication oxycarbonée et modifications électrocardiographiques. A propos de dix-sept observations. (Carbon monoxide poisoning and electrocardiographic changes. A propos of 17 cases). Marseille Med 102: 115-22, 1965. 256
- MOSINGER M, BISSCHOP G and LUCCIONI R : Les manifestations cardiaques dans l'intoxication oxycarbonée. (Cardiac manifestation of carbon monoxide poisoning). Arch Mal Prof Paris 30: 5-12, 1969. 257
- MOTTA G : L'elettrokardiogramma nell'avvelenamento da CO. (Electrocardiogram in CO poisoning). Boll Soc Ital Biol Sper 15: 854, 1940. 258
- ORINIUS E : The late cardiac prognosis after acute carbon monoxide intoxication. Acta Med Scand 183: 239-41, 1968. 259
- PARADE G W and FRANKE H : Beiträge zur Frage der Kohlenmonoxydvergiftung des Herzens. (Carbon monoxide poisoning of the heart). Deutsch Arch Klin Med 185: 294-3-2, 1939. 260

1005051118

- PATZ A : Über elektrokardiogramm-veränderungen bei Kohlenoxyd-und Schlafmittelvergiftungen. (Electrocardiographic changes in carbon monoxide poisoning and soporific poisonings). Arztl Wochschr 4: 653-6, 1949. 261
- SARACOGLU : Modifications électrocardiographiques enregistrees au cours des intoxications, des narcoses et des chocs. (Electrocardiographic modifications during intoxication, narcosis and shock). Arch Mal Coeur 44: 832-8, 1951. 262
- SELING A : WPW-Syndrom nach akuter Intoxikation mit Kohlenmonoxyd. (Wolff-Parkinson-White syndrome following acute intoxication with carbon monoxide). Med Klin 61: 499-501, 1966. 263
- SHAFFER N, SMILAY M G and MacMILLAN F P : Primary myocardial disease in man resulting from acute carbon monoxide poisoning. Am J Med 38: 316-20, 1965. 264
- STEINMANN B : Über das Elektrokardiogramm bei Kohlenoxydvergiftung. (The electrocardiogram in carbon monoxide poisoning). Zeit Kreislaufforsch 29: 281-99, 1937. 265
- STÖRMER A : Über traumatische Herzerkrankungen. (Traumatic heart disease). Deutsch Med Wochsch 64: 235-7, 260-3, 1938. 266
- THIELS H, DURME J P, VERMEIRE P and PANNIER R : Modifications électrocardiographiques immediates et tardives au cours de l'intoxication oxycarbonée aiguë. (Immediate and late electrocardiographic changes in the course of acute carbon monoxide poisoning). Lille Med 17: 191-5, 1972. 267
- VON BERGMANN : Erstickung im Herzmuskel als Ursache der Angina pectoris. (Anoxia of the heart muscle as a cause of angina pectoris). Med Klin 30: 979, 1934. 268
- WENDT L : Die ST-Depression des elektrokardiogramms, ein Symptom des anaeroben Herzstoffwechsels. (ST depression of electrocardiogram, a symptom of anaerobic cardiac metabolism). Arch Kreislaufforsch 8: 74-136, 1941. 269
- ZANARDI S, VILLA A and MONTI G : Modificazioni elettrocardiografiche in operai esposti abitualmente alla inalazione di ossido di carbonio. (Electrocardiographic changes in workers habitually exposed to carbon monoxide). Med Lavore 57: 761-70, 1964. 270

1005051119



- DROGICHINA E A and RYZHKOVA M N : (The clinical picture and diagnosis of diencephalic pathology in occupational poisonings.) Gig Tr Prof Zabol 11: 20-4, 1967. A 479
- FAURE J, VINCENT D, ESCHAPASSE P, LOISEAU P and CASTAING R : Confrontation des signes électrocliniques observés au cours d'intoxications par oxyde de carbone. (Comparison of electro-clinical signs observed in carbon monoxide poisoning.) Rev Neurol (Paris) 112: 287-92, 1965. A 480
- GARLAND H and PEARCE J : Neurological complications of carbon monoxide poisoning. Q J Med 36: 445-55, 1967. A 481
- GARREL S, PERRET J, PELLAT J and ARNOULD P : Syndrome neuro-psychiatrique d'allure frontale: complication post-intervallaire d'une intoxication oxycarbonée. (Neuro-psychiatric syndrome of frontal aspect: delayed complications of carbon monoxide poisoning.) Rev Neurol (Paris) 122: 445-7, 1970. A 482
- GARREL S, PERRET J, PELLAT J and ARNOULD P : Neuro-psychiatric syndrome following carbon monoxide poisoning. Electroenceph Clin Neurophysiol 29: 534, 1970. A 483
- GAULTIER M, FOURNIER E, GERVAIS P and BODIN F : Encéphalopathie pancréatique survenue au décours d'une intoxication oxycarbonée. Comparaison avec l'encéphalopathie post-intervallaire de l'intoxication oxycarbonée. (Pancreatic encephalopathy occurring after carbon monoxide poisoning. Comparison with the post-interval encephalopathy of carbon monoxide poisoning.) Presse Med 72: 3263-5, 1964. A 484
- GIRARDI G, CIS C and PIATTI A : La sindrome nucleo-reticolare cronica nelle intossicazioni professionali. (Chronic nucleo-reticular syndrome in occupational poisonings.) Arch Ital Otol 78: 756-70, 1967. A 485
- GORALSKI H and JANUSZKO L : Zespoły neurologiczne i psychiatryczne po zatruciu tlenkiem węgla. (Neurological and psychiatric syndromes after carbon monoxide poisoning.) Neurol Neurochir Pol 2:633-7, 1968. A 486
- GORDON E B : Carbon monoxide encephalopathy. Brit Med J 5444: 1343, 1965. A 487
- GROHME S, SCHNEIDER H and MASSHOFF W : Encephalopathien bei Vita reducta. (Encephalopathies in vita reducta.) Internist (Berlin) 10: 430-42, 1969. A 488
- GUNTHER K D : Lange verkannte CO-Vergiftungen mit schweren neurologischen Symptomen. (Chronically misdiagnosed CO poisoning with severe neurological symptoms.) Psychiat Neurol Med Psychol (Leipz) 23: 368-77, 1971. A 489
- HAMEL-PUSKARIC H, BERITIC T, JUSIC A and FRANJIC F : Neurološke komplikacije otrovanja ugljičnim monoksidom. (Neurologic complications of carbon monoxide poisoning.) Neuropsihijatrija 18: 147-55, 1970. A 490
- HANSEN D : Berufshedingte Riechstörung infolge chronischer Kohlenoxydeinwirkung. (Occupationally induced anosmia due to chronic carbon monoxide effect.) HNO 5: 140-2, 1970. A 491
- HARADA M and KOZUMA Z : (A case of carbon monoxide poisoning with so-called Sudeck's syndrome, and various neurologic symptoms.) Brain Nerve (Tokyo) 20: 1095-9, 1968. A 492
- HIRAI T : (The relationship between the disturbance of consciousness and behavior disorders.) Adv Neurol Sci (Tokyo) 14: 712-22, 1971. A 493
- IKUTA T : (Somatosensory evoked responses in patients with carbon monoxide poisoning as compared with those in Schizophrenics.) Fol Psychiat Neurol Jap 23: 285-9, 1969. A 494

1005051271

(Gorski, 1962), reduced response of left ventricle to the nitroglycerin test (Gokina, 1971), and elevation of serum enzymes which reflect myocardial injury (Jaffe, 1965). The postmortem examination of the heart in patients dying from carbon monoxide poisoning reveals necrosis and hemorrhagic infarction, fibrosis and fatty degeneration (Koelsch, 1936; Nagel, 1937; Monau, 1940; Breu, 1942; Binet and Bétourné, 1951; Holm, 1950; Caccuri, 1955; Ritter, 1956; Klavis and Schulz, 1966; Borst, 1967; Sobotka and Sobotka, 1969; Caroff et al., 1970).

The rat has been the laboratory animal most extensively used to investigate myocardial effects of carbon monoxide. Asmussen and Paulsen (1953) exposed immature rats for 60 days to an atmosphere containing carbon monoxide. The blood levels were kept at 50 to 60 % carboxyhemoglobin. Compared with control rats, the carbon-monoxide-treated rats were inferior in their ability to swim till exhausted and to withstand low oxygen tension. The carbon-monoxide-treated rats showed cardiac hypertrophy and a slight but significant increase in the relative number of coronary capillaries. Suzuki (1969) administered 1% carbon monoxide for 10 min to mature rats and the animals were sacrificed from 10 min to 24 hours after cessation of inhalation. The electron microscopic examination of the heart revealed intracellular edema, swelling of mitochondria and sarcoplasmic reticula, disruption and reduction of cristae, disappearance of mitochondria, appearance of lipofuscin pigment granules and lysosomes and increase of glycogen granules and fat droplets. The author concluded that the effects of carbon monoxide on the heart result not only from hypoxemia but also from the direct toxic effects on the specific respiratory enzymes.

1005051121

Holczabek (1971) arrived at a similar conclusion following exposure of rats to 3% carbon monoxide. Slater (1950) demonstrated inhibition of dihydrocozymase oxidase of heart muscle exposed to carbon monoxide in vitro.

The direct effects of carbon monoxide on the monkey heart have not been investigated. Since there is a species difference relating to pulmonary effects, it is reasonable to suspect that this may also apply to the heart. The rabbit heart shows cardiac necrosis, which could be interpreted to be the result of hypoxemia rather than the direct effect of carbon monoxide (Veith, 1940)..

1005051122

## BIBLIOGRAPHY

## V. CIRCULATORY SYSTEM

## B. Myocardium

- ANDERSON W H : Acute exposure to cigarette smoke as a cause of hypoxia. Chest 59: 33S-34S, 1971. Reprin 271
- ASMUSSEN E and PAULSEN N V : Cardiac hypertrophy in CO-treated young rats and their ability to withstand stress. Acta Physiol Scand 29: 307-13, 1953. 272
- BINET L and BÉTOURNÉ C : Intoxication oxycarbonée à localisation musculaire. Sem Hop Paris 27: 2859-60, 1951. 273
- BREU W : Die Kohlenmonoxydvergiftung des Herzens. (Carbon monoxide poisoning of the heart). Wien Klin Wochschr 55: 867-70, 1942. 274
- BORST J R : De cardiale aspecten van de chronische koolmonoxydevergiftiging. (The cardiac aspects in chronic carbon monoxide poisoning). Nerl T Geneesk 11: 573-9, 1967. 275
- BRODY J S and COBURN R F : Carbon monoxide-induced arterial hypoxemia. Science 164: 1297-8, 1969. 276
- BRODY J S and COBURN R F : Effects of elevated carboxyhemoglobin on gas exchange in the lung. Ann NY Acad Sci 174: 255-60, 1970. 277
- CACCURI S : L'apparato circolatorio nell' intossicazione da ossido di carbonio. (The cardiovascular system intoxicated by carbon monoxide). Rif Med 69: 649-56, 1955. 278
- CAROFF J, DEHOUE P and DEROBERT L : Cardiopathie congénitale (Taussig-Bing) et intoxication par l'oxyde de carbone. (Congenital cardiopathy (Taussig-Bing) and carbon monoxide poisoning). Med Leg Domm Corpor Paris 3: 84-7, 1970. 279
- CHEVALIER R B, KRUMHOLZ R A and ROSS J C : Effect of carbon monoxide inhalation on the cardiopulmonary responses of nonsmokers to exercise. J Lab Clin Med 62: 867, 1963. 280
- DIAMANT-BERGER F, GAJDOS P, RAPIN M and GOULON M : Aspects hemodynamiques de l'intoxication oxycarbonée aiguë massive humaine. (Hemodynamic aspects of acute massive carbon monoxide poisoning in humans). Eur J Toxicol 3: 211-26, 1970. 281
- FLAXMAN N : Cardiac Review of 1938. Illinois Med J 76: 182-97, 1939. 282
- GOKINA M S : Myocardial contractile function in workers exposed to prolonged action of small carbon monoxide concentrations). Vrach Delo 8: 130-2, 1971. 283
- GORSKI J : Balistokardiogram w ostrym zatruciu tlenkiem wegla. (Ballistocardiogram in acute poisoning with carbon monoxide). Pol Tyg Lek 17: 872-76, 1962. 284
- GUILLERM R, BADRE R and GAUTIER H : Effets du séjour dans une atmosphère à faible concentration d'oxyde de carbone sur les réactions circulatoires et respiratoires à l'effort musculaire et sur l'acuité visuelle nocturne. Biometeorology, Tromp, Pergamon Press, London 2: 306-13, 1967. 285
- HOLCZABEK W : Über die zyanochrome myelinige Entmischung des Herzfleisches nach Tod an Kohlenoxydvergiftung und nach Tod an Hypoxämie. (Cyanochromic myelin disintegration of the myocardium after death from carbon monoxide poisoning and death from hypoxemia). Zentralbl Allg Pathol 114: 83-9, 1971. 286
- HOLM K F : Dauerschaden des Herzens nach CO-Vergiftung. (Permanent heart damage after CO poisoning). Med Klin 45: 1427-9, 1950. 287
- JAFFE N : Cardiac injury and carbon monoxide poisoning. S Afr Med J 39: 611-5, 1965. 288
- JAGIĆ N and ZIMMERMANN O : Zur Klinik und Therapie der Myokarderkrankungen. (The clinic and therapy of myocardial diseases). Wien Klin Wochschr 47: 1217-21, 1955-8, 1954. 289

1005051123

## Bibliography V. B

- KALIAEVA S I: (Changes in the cardiovascular system after accidental acute intoxications with carbon monoxide). Klin Med 29: 60-4, 1951. 290
- KATZCHMANN R: Das Herz-Kreislauf-System bei Schwerarbeitenden (Physiologie-Pathologie-Prophylaxe. (Cardiovascular system in workers. Physiology-Pathology-Prevention). Z Gesamte Inn Med 25: 738-46, 1970. 291
- KLAUSEN K, RASMUSSEN B, GJELLEROD H, MADSEN H and PETERSEN E: Circulation, Metabolism and ventilation during prolonged exposure to carbon monoxide and to high altitude. Scand J Clin Lab Invest 22 Suppl: 103: 26-38, 1968. 292
- KLAVIS G and SCHULZ L C: Herzscheiden bei der akuten Kohlenmonoxyd-Vergiftung. (Heart damage in acute carbon monoxide poisoning). Arch Toxikol 21: 250-60, 1966. 293
- KOELSCH : (Circulatory changes in industrial poisonings). Med Klin 32: 916-7, 1936. 294
- KROETZ C: Herzscheiden nach Kohlenoxydvergiftungen. (Cardiac damage after carbon monoxide intoxication). Dtsch Med Wochenschr 62: 1365-9, 1414-17, 1936. 295
- MONAUI J: Myokardschädigung als Spätfolge einer Kohlenoxydvergiftung. (Myocardial damage as a sequel to carbon monoxide poisoning). Munch Med Wochenschr 87: 659, 1940. 296
- NAGEL H G: Zur Frage der Koronarscheiden nach Leuchtgasvergiftungen. (Coronary damage in illuminating gas poisoning). Dtsch Med Wochenschr 63: 301-2, 1937. 297
- PIRNAY F, DUJARDIN J, DEROANNE R and PETIT J M: Muscular exercise during intoxication by carbon monoxide. J Appl Physiol 31: 573-5, 1971. 298
- PIRNAY F, DEROANNE R, DUJARDIN J and PETIT J M: Exerice musculaire maximum sous intoxication oxycarbonée. (Maximal muscular exertion under carbon monoxide poisoning). J Physiol Paris 63: 87A-8, 1971. 299
- RITTER U: Ekg-Veränderungen bei Vergiftungen. (Changes in electrocardiogram due to intoxications). Arztl Wochenschr 11: 721-6, 1956. 300
- ROSKRAMM K: Vermindert das Rauchen die sportliche Leistungsfähigkeit? (Does smoking lessen the capacity for athletic activities?) Med Klin 59(14): 591, 1964. 301
- SLATER E C: The succinic oxidase and dihydrocozymase oxidase systems in heart muscle and kidney preparations. Nature 165: 674-5, 1950. 302
- SOBOTKA W and SOBOTKA S: Uszkodzenie miesnia sercowego w przebiegu zatrucia tlenkiem wegla u dzieci. (Damage of the cardiac muscle in the course of carbon monoxide poisoning in children.) Przegl Lek 25: 251-2, 1969. 303
- SUZUKI T: Effects of carbon monoxide inhalation on the fine structure of the rat heart muscle. Tohoku J Exp Med 97: 197-211, 1969. 304
- VEITH G: Experimentelle Untersuchungen zur Wirkung von Adrenalin auf den Herzmuskel. (Experimental investigations on the effect of epinephrine on cardiac muscle. Arch Kreislaufforsch 6: 335-60, 1940. 305
- VOGEL J A and GLESER M A: Effect of carbon monoxide on oxygen transport during exercise. J Appl Physiol 32: 234-9, 1972. 306
- GLESER M A,  
VOGEL J A, WHEELER R C and WHITTEN B K: Carbon monoxide and physical work capacity. Arch Environ Health 24: 198-203, 1972. 307
- ZIEGLER K: Kohlenoxydgasvergiftung und Myokard. (Carbon monoxide intoxication and the myocardium.) Dtsch Med Wochenschr 62: 389-91, 1936. 308

1005051124

### V C. Coronary Circulation

In recent years there has been an increasing number of publications associating coronary heart disease with the carbon monoxide contained in cigarette smoke (Jaffe, 1968; Dinman, 1969; Robin *et al.*, 1969; Goldsmith, 1970; Sz8118si *et al.*, 1970; Tibblin, 1971; Schievelbein and Eberhardt, 1972; and Bersay Marland/ 1972). The evidence for stating that the carbon monoxide content of cigarette smoke caused coronary heart disease is indirect. A review of the investigations concerned reveals that the levels of carboxyhemoglobin in the blood of habitual smokers do not cause coronary heart disease.

The effect of exposure to lower concentrations of carbon monoxide in high-pollution areas of Los Angeles has been examined by Cohen *et al.* (1969). The case fatality rates for patients admitted with myocardial infarction to 35 hospitals during 1958 were examined. The results indicate that there was an increase in fatality rate in high-pollution areas and that this difference was evident during periods of relatively increased carbon monoxide pollution. However, it was not possible to prove cause and effect relationship between carbon monoxide and high fatality rate, since there are other pollutants involved. In the same city, Haywood *et al.* (1972) examined 34 patients with acute myocardial infarction and 35 control patients with diverse diseases. Carboxyhemoglobin levels averaged 5.14 % for the infarct patients and 4.8 % for the controls; there was no clear-cut relationship between carbon monoxide levels and acute infarction. For patients with angina pectoris, exposure to the heavy morning freeway traffic in Los Angeles

1005051125

caused a decrease in exercise performance that initiated the onset of angina (Aronow et al., 1972). The mean blood levels of carboxyhemoglobin in % were  $1.12 \pm 1.20$  before,  $5.08 \pm 1.19$  immediately after leaving the freeway, and  $2.91 \pm 0.93$  two hours later. Any one of the pollutants other than carbon monoxide may be responsible for quicker development of angina after less cardiac work. Ten patients with angina pectoris were examined by Aronow and Rokaw (1971) and Aronow et al. (1971) following the smoking of low-nicotine cigarettes. After each subject had smoked 8 cigarettes, at the rate of one every 30 min, the carboxyhemoglobin level in the blood rose from 1.58 to 7.79%. This was accompanied by a decrease in exercise tolerance. These results cannot be interpreted to mean that carbon monoxide alone is the cause of the decrease in exercise tolerance. The only direct proof would be to repeat similar observations on patients inhaling carbon monoxide mixture. De Bias et al. (1972) exposed dogs with myocardial infarction to 100 ppm carbon monoxide for 14 weeks. The elevation of the blood carboxyhemoglobin level to 14% did not influence the electrocardiogram nor the serum enzymes that would be expected to accompany increasing severity of hypoxia. Carbon monoxide alone, producing up to 14% saturation of carboxyhemoglobin, does not appear to exaggerate myocardial infarction in dogs.

months,  
Exposure of rabbits for up to 14/ resulting in a blood level of 15 to 40% carboxyhemoglobin, causes myocardial damage (Andersson, 1972). The lesions are similar to those reported for patients who have recovered from acute carbon monoxide poisoning. Some of these patients manifested anginal

1005051126

attacks (Kroetz, 1936a; Beck and Suter, 1938; Hubert, 1943; Zeh, 1960) and myocardial infarction (Kroetz, 1936b; Wiktor, 1954; Anderson et al., 1967).

The effects of inhalation of 0.1 or 5 % carbon monoxide, sufficient to raise the carboxyhemoglobin level to between 5 and 25 % in dogs and humans, were reported by Ayres et al. (1969, 1970). There was an increase in coronary blood flow and alteration of lactate and pyruvate metabolism. Most of these changes could be accounted for by hypoxemia, although a direct effect of carbon monoxide on the coronary vessels has not been excluded.

1005051127



## V. CIRCULATORY SYSTEM

## C. Coronary Circulation

- |  | Reprint |
|--|---------|
| ANDERSON R F, ALLENSWORTH D C and DeGROOT W J: Myocardial toxicity from carbon monoxide poisoning. <u>Ann Intern Med</u> 67: 1172-82, 1967.  | 309     |
| ANDERSSON A: A study of cardio-vascular alterations in animals exposed to carbon monoxide during long time. <u>Opuscula Med</u> 17/5: 203-9, 1972.   | 310     |
| ARONOW W S, DENDINGER J and ROKAW S N: Heart rate and carbon monoxide level after smoking high-, low-, and non-nicotine cigarettes. A study in male patients with angina pectoris. <u>Ann Int Med</u> 74: 697-702, 1971.   | 311     |
| ARONOW W S, HARRIS C N, ISBELL M W, ROKAW S N and IMPARATO B: Effect of freeway travel on angina pectoris. <u>Ann Int Med</u> 77: 669-76, 1972.  | 312     |
| ARONOW W S and ROKAW S N: Carboxyhemoglobin caused by smoking nonnicotine cigarettes. Effects in angina pectoris. <u>Circulation</u> 44: 782-8, 1971.  | 313     |
| AYRES S M, GIANNELLI S Jr, and MUELLER H: Myocardial and systemic responses to carboxy-hemoglobin. <u>Ann NY Acad Sci</u> 174: 268-93, 1970.   | 314     |
| AYRES S M, MUELLER H S, GREGORY J J, GIANNELLI S and PENNY J L: Systemic and myocardial hemodynamic responses to relatively small concentrations of carboxyhemoglobin (COHb). <u>Arch Environ Health</u> 18: 699-709, 1969.  | 315     |
| BECK H G and SUTER G M: Role of carbon monoxide in the causation of myocardial disease. <u>JAMA</u> 110: 1982-6, 1938.   | 316     |
| COHEN S I, DEANE M and GOLDSMITH J R: Carbon monoxide and survival from myocardial infarction. <u>Arch Environ Health</u> 19: 510-7, 1969.   | 317     |
| DeBIAS D A, BIRKHEAD N C, BANERJEE C M, KAZAL L A, HOLBURN R R, GREENE C H, HARPER W V, ROSENFELD L M, MENDUKE H, WILLIAMS N and FRIEDMAN M H F: The effects of chronic exposure to carbon monoxide on the cardiovascular and hematologic systems in dogs with experimental myocardial infarction. <u>Int Arch Arbeitsmed</u> 29:253-67, 1972. | 318     |
| DINMAN B D: Discussion. Toxicological appraisal of carbon monoxide. <u>J Air Pollut Contr Ass</u> 19: 727-9, 1969.   | 319     |
| GOLDSMITH J R: Carbon monoxide research- recent and remot. <u>Arch Environ Health</u> 21: 118-20, 1970.  | 320     |
| HAYWOOD L F, WALBERG C B, KERR F, MOHSENIN M and MOHLER J: Carbon monoxide levels in acute myocardial infarction. <u>J Na Med Ass</u> 64(2): 114-6, 1972.  | 321     |
| HUBERT G: Kritische Betrachtungen zum Begriff Koronarinsuffizienz. (Critical notes on coronary insufficiency.) <u>Z Kreislaufforsch</u> 35: 145-56, 1943.  | 322     |
| JAFFE N: Role of carbon monoxide in coronary disorders. <u>New Eng J Med</u> 279: 111, 1968.   | 323     |
| KROETZ C: Kohlenoxyd und Herzinfarkt. (Carbon monoxide and heart infarct.) <u>Munch Med Wochschr</u> 83: 951, 1936a  | 324     |
| KROETZ C: Angina pectoris nach Rauchgasvergiftung. (Angina pectoris after inhaled gas poisoning.) <u>Med Klin</u> 32: 1521-4, 1936b  | 325     |
| MARLAND P and BERSAY C: De l'intérêt d'être coronariens. (The advantage of being a coronary patient.) <u>Nouv Presse Med</u> 1: 1097-8, 1972.  | 326     |

1005051428

- ROBIN E, RAVENS K G and BING R J : Die Wirkung von Alkohol, Nikotin und Zigarettenrauchen auf das Herz. (The effect of alcohol, nicotine and cigarette smoking on the heart.) Deutsch Med J 20: 19-29, 1969. 327
- SCHIEVELBEIN H and EBERHARDT R : Cardiovascular actions of nicotine and smoking. J Natl Canc Ins 48: 1785-94, 1972. 328
- SZÖLLÖSI E, MEDVE F and JENEY E : Angaben zur Wirkung des niedrigen Kohlenmonoxyd-Gehaltes in der Luft auf den Menschen. (Data on the effect of a low carbon monoxide content in the air on man.) Z Arbeitsmed 20: 263-8, 1970. 329
- TIBBLIN G : Hjärtinfarkt och rökning. (Harmful clinical effects of smoking. Myocardial infarct and smoking.) Soc Med Tid 2: 65-7, 1971. 330
- WIKTOR Z : Sprawozdanie z posiedzen naukowych wroclawskiego oddzialu towarzystwa internistow polskich w r. 1952. (Report on the scientific session of the Wroclaw branch of the Polish Society of Internal Medicine in 1952. Pol Arch Med Wewn 24: 596-7, 1954. 331
- ZEH E : (Heart function disorders after carbon monoxide or E605 poisoning. Med Welt 1: 339-40, 1960. 332

1005051129

#### V D. Systemic Circulation

Cigarette smoking causes vasoconstriction of most vascular beds. These effects are brought about by nicotine contained in the smoke. The carbon monoxide absorbed during smoking does not contribute to the vascular effects. In animals, the pattern of action of carbon monoxide is vasodilatation with elevation of body temperature (Binet and Burstein, 1948; Coret and Hughes, 1964; Nielsen, 1971).

Acute carbon monoxide poisoning is accompanied by a fall in aortic blood pressure (Litzner, 1936; Deviatka, 1956; Navratil, 1956; Vyskocil and Novotny, 1956; Chudzikiewicz, 1957; Mihai and Weber, 1964; Heidrich et al., 1970). Hypotension has also been noted following exposure to carbon monoxide in dogs (Brewer, 1937; von Oettingen et al., 1941), cats (Kayser, 1939; Maurer, 1941), rabbits (Nishigori, 1932; Süpfle, 1934) and rats (Truhaut et al., 1968). The fall in blood pressure is entirely due to vasodilatation, which has been demonstrated in dogs (Sulotto et al., 1969 a and b). In man vasoconstriction of the hand reflexly induced by cold is reduced by levels of 19 and 25 % carboxyhemoglobin (Heistad and Wheeler, 1972).

The influence of carbon monoxide on capillary permeability has been investigated in humans and animals. In man, exposure to carbon monoxide for 8 days caused an increase in the permeability of the capillaries to albumin (Siggaard-Andersen et al., 1968, 1969). The increase in permeability could not be demonstrated in the calf muscle (Petersen et al., 1968). In rabbits, guinea pigs and rats there is an increase in permeability in the peritoneal cavity (Güthert et al., 1970) and subcutaneous tissue (Van Liew, <sup>1968 a</sup> and b, 1970).

1005051130

## V. CIRCULATORY SYSTEM

## D. Systemic Circulation

- BINET L and BURSTEIN M : Intoxication par l'oxyde de carbone et tonus des vaisseaux periphériques. (Intoxication by carbon monoxide and peripheral vascular tone.) C R Soc Biol 142: 1487-8, 1948. Reprint 333
- BREWER N R : Blood-pressure responses to acute carbon monoxide poisoning. Am J Physiol 120: 91-9, 1937. 334
- CHUDZIKIEWICZ T : Uszkodzenie miesnia sercowego w przebiegu zatrucia tlenkiem wegla. (Myocardial injury in carbon monoxide poisoning.) Przeg Lek 13: 88-9, 1957. 335
- CORET I A and HUGHES M J : A further study of hypoxic smooth muscle. Arch Int Pharmacodyn 149: 330-53, 1964. 336
- DEVIATKA D G : (Etiologic role of carbon monoxide on the development of hypotensive condition.) Terap Arkh 28: 29-32, 1956. 337
- GÖTHERT M, LUTZ F and MALORNY G : /monoxide partial pressure in tissue of different animals. Environ Res 3: 303-9, 1970. Carbon 338
- HEIDRICH H, BARCKOW D and FRISIUS H : Untersuchungen über den Einfluß von Actihaemyl auf den peripheren Widerstand und das Herzzeitvolumen. (Studies on the effect of actihaemyl on peripheral vascular resistance and cardiac output.) Z Kreislaufforsch 59: 251-61, 1970. 339
- HEISTAD D D and WHEELER R C : Effect of carbon monoxide on reflex vasoconstriction in man. J Appl Physiol 32: 7-11, 1972. 340
- KAYSER H W : Der Einfluß des Kohlenoxyds auf vasomotorische Reaktionen. (Effect of carbon monoxide on vasomotor reactions.) Arch Exp Path Pharmacol 192: 625-33, 1939. 341
- LITZNER S : Über Kreislauf- und Herzscheidigungen bei der Kohlenoxydvergiftung. (Circulatory and cardiac damage in carbon monoxide poisoning.) Med Klin 32: 630-1, 1936. 342
- MAURER F W : The effects of carbon monoxide anoxemia on the flow and composition of cervical lymph. Am J Physiol 133: 170-9, 1941. 343
- MIHAI N and WEBER A : Cercetari asupra oxicarbonismului acut si cronic Modificarile tensiunii arteriale. (Research on acute and chronic carbon monoxide poisoning. Changes in arterial pressure.) Med Intern Bucur 16: 1113-9, 1964. 344
- NAVRATIL M : Vliv kyslicniku uhelnateho na krevni obch. (The effect of carbon monoxide on the vascular apparatus.) Prakt Lek 36: 89-90, 1956. 345
- NIELSEN B : Exercise temperature plateau shifted by a moderate carbon monoxide poisoning. J Physiol Paris 63: 362-5, 1971. 346
- NISHIGORI H : (The effect of carbon monoxide on blood pressure and the electrocardiogram.) Nippon Nai Gak Zasshi 20: 603-9, 1932. 347
- PETERSEN F B, SIGGAARD-ANDERSEN J, KRISTENSEN J H and KJELDSEN K : Capillary filtration rate on the human calf during exposure to carbon monoxide and hypoxia (3454m). Scan J Clin Lab Invest 22 Suppl 103: 49-54, 1968. 348
- SIGGAARD-ANDERSEN J, PETERSEN F B, HANSEN T I and MELLEMGAAARD K : Plasma volume and vascular permeability during hypoxia and carbon monoxide exposure. Scan J Clin Lab Invest 22 Suppl 103: 39-48, 1968. 349

1005051131

Bibliography V. D

page 00

- SIGGAARD-ANDERSEN J, PETERSEN F B, HANSEN T I and MELLEMGAAARD K: Vascular permeability and plasma volume changes during hypoxia and carbon monoxide exposure. Angiology 20: 356-8, 1969. 350
- SULOTTO F, MEO G, POLI G and RUBINO G F: Studio delle modificazioni emodinamiche nell'intossicazione sperimentale acuta da ossido di carbonio. Circolo sistemico. (Hemodynamic changes in acute experimental carbon monoxide poisoning. Systemic circulation.) Med Lav 60: 97-108, 1969. 351
- SULOTTO F, BONZANINO A, MEO G and RUBINO G F: Studio delle modificazioni emodinamiche nell'intossicazione sperimentale acuta da ossido di carbonio. 2. Circoli distrettuali. Coronarico, carotideo, mesenterico, renale, iliaco. (Hemodynamic changes in acute experimental carbon monoxide poisoning. 2. Regional circulation. Coronary, carotid, mesenteric, renal, iliac.) Med Lav 60: 109-17, 1969. 352
- SÜPFLE K: Zur Frage der chronischen Kohlenoxydvergiftung. (Chronic carbon monoxide poisoning.) Dtsch Med Wochenschr 60: 1263-7, 1934. 353
- TRUHAUT R, BOUDÈNE C, CLAUDE J R, JACOTOT B: Recherches sur les effets de l'exposition prolongée du lapin et du rat à de très faibles concentrations d'oxyde de carbone. III. Etude de l'action sur le système cardiovasculaire (1). (Research of the effects of prolonged exposure of rats and rabbits to very low concentrations of carbon monoxide. III. The effect on the cardiovascular system.) Arch Mal Prof Paris 29: 189-96, 1968. 354
- VAN LIEW H D: Interaction of CO and O<sub>2</sub> with hemoglobin in perfused tissue adjacent to gas pockets. Res Physiol 5: 202-10, 1968. 355
- VAN LIEW H D: Coupling of diffusion and perfusion in gas exit from subcutaneous pocket in rats. Amer J Physiol 214: 1176-85, 1968. 356
- VAN LIEW H D: Interaction of CO and O<sub>2</sub> with hemoglobin in perfused tissue adjacent to gas pockets. USAF Aerospace Med 21: 212-20, 1970. 357
- VON OETTINGEN W F, DONAHUE DD and VALAER P J: On the mechanism of carbon monoxide poisoning. J Pharmacol Exp Ther 72: 42, 1941. 358
- VYSKOCIL J and NOVOTNY S: Nase zkušenosti akutními otravami kyslíkem uhelnatým. (Our experience with acute carbon monoxide poisonings.) Prakt Lek 36: 88-9, 1956. 359

1005051132

V E. Arteries

case

Although no specific of arterial disease caused by carbon monoxide associated with cigarette smoking has been reported, there have been repeated suggestions of cause and effect relationship. The facts are as follows:

1. Patients who have been exposed to acute carbon monoxide poisoning develop skeletal muscle necrosis. (Mautner, 1955). Volkmann's contracture (Ortizaga, 1967) or venous thrombosis (Heidrich and Klems, 1969). Similar lesions have not been reported following exposure to low levels of carbon monoxide.
2. In patients with thromboangiitis obliterans or Buerger's disease, Astrup (1964) pointed out a connection between smoking and increased affinity of hemoglobin. Astrup (1966 a and b) and Astrup et al. (1966) showed the increase in affinity for oxygen to be associated with carbon monoxide present in tobacco smoke, since higher carboxyhemoglobin levels were observed in smokers with thromboangiitis obliterans than in healthy smokers. Mulhausen et al. (1967) confirmed this observation in another group of patients. Kjeldsen and Mozes (1969) and Kjeldsen (1969) noted in a third group of patients that the carboxyhemoglobin saturations and cholesterol levels are higher in controls. Birnstingl et al. (1966) demonstrated that patients with thromboangiitis obliterans did not show a greater alteration in oxygen affinity produced by smoking/compared with normal smokers. The possibility that carboxyhemoglobin increases blood viscosity and therefore reduces the velocity of blood circulation and hastens the tendency to thrombus formation, has been excluded by measurements performed by Solvsteen and Kristjansen (1968).

1005051133

3. That exposure to carbon monoxide could lead to arteriosclerosis was proposed by Hueper (1944) as part of his anoxemia theory. Astrup and his collaborators have attempted to find experimental support for this theory in cholesterol-fed rabbits — see reviews by Astrup (1967, 1969, 1970, 1972) and by Astrup and Kjeldsen (1970). The exposure to carbon monoxide enhanced the development of atheromatosis (Astrup et al., 1970 a and b). The appearance of lesions was accompanied by elevation of serum lipid levels (Truhaut et al., 1968; Kjeldsen, 1970a), change in lactate dehydrogenase isoenzymes of the aortic arch (Hellung-Larsen et al., 1968), increased endothelial permeability (Wanstrup et al., 1969), and ultrastructural intimal changes (Kjeldsen et al., 1972). In human subjects exposed to carbon monoxide, an increase in capillary filtration rate (Siggaard-Andersen et al., 1967) and elevation of serum lipid levels (Kjeldsen and Damgaard, 1965, 1968; Kjeldsen, 1970b) have been demonstrated. It has been suggested that carbon monoxide inhibits synthesis of cholesterol, leading to accumulation of lanosterol (Gibbons and Mitropoulos, 1972). Another effect of carbon monoxide is an increase in mitochondrial enzymic activity, which stimulates lipid synthesis within the artery (Whereat, 1970). It has not been possible to develop atherosclerosis in animals exposed to carbon monoxide without supplemental cholesterol feeding.

**1005051134**

4. Examination of individuals who have been exposed to an environment of up to 1,000 ppm carbon monoxide with carboxyhemoglobin levels of blood between 2 and 26 % for an average duration of 10.5 years did not reveal any early development of arteriosclerosis (Prerovská and Drdková, 1967 a and b;

1971). The average values of serum lipid levels did not exceed the normal range. The results of experiments on rabbits do not apply to epidemiologic surveys in humans.

1005051135



## BIBLIOGRAPHY

page 14

## V. CIRCULATORY SYSTEM

## E. Arteries

- ASTRUP P: An abnormality in the oxygen-dissociation curve of blood from patients with non-specific myocarditis. Lancet 2: 1152-4, 1964. Reprint 360
- ASTRUP P: Den kliniske betydning af forskydninger i oksihæmoglobinet's dissociationskurve. Nordisk Med Stockholm 76: 1039-41, 1966a. 361
- ASTRUP P: Hæmmet iltafgift fra blodet og udviklingen af oblittererende arteriesygdomme. (Impeded oxygen release from the blood and the development of obliterating arterial diseases.) Ug Læger 128: 701-6, 1966b. 362
- ASTRUP P: Carbon monoxide and peripheral arterial disease. Scan J Clin Lab Invest 93: 193-7, 1967. 363
- ASTRUP P: Effects of hypoxia and of carbon monoxide exposures on experimental atherosclerosis. Ann Int Med 71: 426-7, 1969. 364
- ASTRUP P: Karbeskadigende virkning af CO og hypoxi. (Smoking and coronary disease. Vessel injuring effect of CO and hypoxia.) Lakartidningen 67: 256-61, 1970. 365
- ASTRUP P: Some physiological and pathological effects of moderate carbon monoxide exposure. Brit Med J 4: 447-52, 1972. 366
- ASTRUP P, HELLUNG-LARSEN P, KJELDSSEN K and MELLEMGAAARD K: The effect of tobacco smoking on the dissociation curve of oxyhemoglobin. Investigations in patients with occlusive arterial diseases and in normal subjects. Scand J Clin Lab Invest 18: 450-7, 1966. 367
- ASTRUP P, KJELDSSEN K and WANSTRUP J: Enhancing influence of carbon monoxide on the development of atheromatosis in cholesterol-fed rabbits. J Atheroscler Res 7: 343-54, 1967. 368
- ASTRUP P, KJELDSSEN K and WANSTRUP J: The effects of exposure to carbon monoxide, hypoxia and hyperoxia on the development of experimental atheromatosis in rabbits. Atherosclerosis Proceedings, 2nd International Symposium, R.J. Jones (Editor), Chicago, Springer-Verlag 108-11, 1970. 369
- ASTRUP P, KJELDSSEN K and WANSTRUP J: Effects of carbon monoxide exposure on the arterial walls. Ann NY Acad Sci 174: 294-300, 1970. 370
- BIRNSTINGL M A, COLE P J and HAWKINS L: Variation in blood oxygen dissociation with age, smoking, and Buerger's disease. Brit J Surg 53: 986, 1966. 371
- GIBBONS, G F/ MITROPOULOS K A: Inhibition of cholesterol biosynthesis by carbon monoxide: accumulation of Lanosterol and 24, 25-Dihydrolanosterol. Biochem J 127: 315-7, 1972. 372
- HEIDRICH H and KLEMS H: Doppelseitige Thrombose der Vena poplitea mit diffuser Muskelnekrose nach CO-Intoxikation. (Bilateral thrombosis of the popliteal vein with diffuse muscular necrosis following CO intoxication.) Deutsch Med Wschr 94: 1367-70, 1969. 373
- HELLUNG-LARSEN P, LAURSEN T, KJELDSSEN K and ASTRUP P: Lactate dehydrogenase isoenzymes of aortic tissue in rabbits exposed to carbon monoxide. J Altheroscler Res 8: 343-9, 1968. 374
- HUEPER W C: Arteriosclerosis. Arch Path 38: 161-81, 245-85, 350-64, 1944. 375

1005031136

- KJELDSSEN K : Smoking and atherosclerosis. Investigations on the significance of the carbon monoxide content in tobacco smoke in atherogenesis. Munksgaard, Copenhagen 1-145, 1969. 376
- KJELDSSEN K : Carboxyhemoglobin and serum cholesterol levels in smokers correlated to the incidence of occlusive arterial disease. Atherosclerosis Proceedings of the 2nd International Symposium, R. J. Jones, Editor, Springer-Verlag, New York 378-81, 1970. 377
- KJELDSSEN K : CO-eksposition og aterosklosefrekvens. (Smoking and coronary disease. CO exposure and frequency of arteriosclerosis.) Lakartidningen 67: 262-5, 1970. 378
- KJELDSSEN K, ASTRUP P and WANSTRUP J : Ultrastructural intimal changes in the rabbit aorta after a moderate carbon monoxide exposure. Atherosclerosis 16: 67-82, 1972. 379
- KJELDSSEN K and DAMGAARD F : Influence of prolonged carbon monoxide exposure and altitude hypoxia on serum lipids in man. Scand J Clin Lab Invest 22 Suppl 103: 16-9, 1968. 380
- KJELDSSEN K and MOZES M : Buerger's disease in Israel. Investigations on carboxyhemoglobin and serum cholesterol levels after smoking. Acta Chir Scand 135: 495-8, 1969. 381
- MAUTNER L S : Muscle necrosis associated with carbon monoxide poisoning. Arch Path 60 136-8, 1955. 382
- MULHAUSEN R, ASTRUP P and KJELDSSEN K : Oxygen affinity of hemoglobin in patients with cardiovascular diseases, anemia and cirrhosis of the liver. Scand J Clin Lab Invest 19:291, 1967. 382a
- ORIZAGA M, DUCHARME F A, CAMPBELL J S and EMBREE G H : Muscle infarction and Volkmann's contracture following carbon monoxide poisoning. J Bone Joint Surg 49: 965-70, 1967. 383
- PREROVSKA I and DRDKOVA S : Vliv chronickeho pusobeni kyslicniku uhelnateho na biochemicke zmeny v seru vzhledem k ateroskleroze. (The effect of chronic exposure to carbon monoxide on biochemical changes in the blood with respect to atherosclerosis.) Prac Lek 19: 1-4, 1967a. 384
- PREROVSKA I and DRDKOVA S : Vliv chronickeho pusobeni prumyslovych skodlivin na exponovane pracovníky vzhledem k rozvoji aterosklerozy. (Influence of the chronic action of industrial anoxic agents on exposed workers in relation to the development of atherosclerosis.) Cas Lek Cesk 106: 754-9, 1967b. 385
- PREROVSKA I and DRDKOVA S : Der Einfluß der chronischen Einwirkung von Kohlenoxyd auf den klinischen Zustand und biochemische Veränderungen im Serum exponierter Personen in Hinsicht auf die vorzeitige Entwicklung der Atherosklerose. (Influence of chronic action of carbon monoxide on the clinical status of biochemical changes in the serum of exposed persons on development of atherosclerosis with influence to the premature.) Int Arch Arbeitsmed 28: 175-88, 1971. 386
- SIGGAARD-ANDERSEN J, KJELDSSEN K, PETERSEN F B and ASTRUP P : A possible connection between carbon monoxide exposure, capillary filtration rate and atherosclerosis. Acta Med Scand 182: 397-9, 1967. 387
- SÖLVSTEEN P and KRISTJANSEN P F : Carbon monoxide, blood viscosity and development of Buerger's disease. Z Kreislaufforsch 57: 790-2, 1968. 388
- TRUHAUT R, BOUDENE C and CLAUDE J R : Sur quelques reflets humoraux de l'intoxication chronique par l'oxyde de carbone chez le Lapin. (On some humoral effects of chronic carbon monoxide poisoning in rabbits.) Ann Biol Clin Paris 26: 1249-60, 1968. 389
- WANSTRUP J, KJELDSSEN K and ASTRUP P : Acceleration of spontaneous intimal-subintimal changes in rabbit aorta by a prolonged moderate carbon monoxide exposure. Acta Path Microbiol Scand 75: 353-62, 1969. 390
- WHEREAT A F : Is atherosclerosis a disorder of intramitochondrial respiration? Ann Intern Med 73: 126-7, 1970. 391

1005051137

## V F. Blood Cells and Plasma

Exposure to carbon monoxide causes an increase in platelet adhesiveness in atherosclerotic patients (El-Ebrashy et al, 1967) and in rabbits (Birnstingl et al, 1971). There is also an increase in fibrinolytic activity in patients suffering from carbon monoxide poisoning (El-Attar, 1968 a and b) and in experimental animals (Candura and Craveri, 1964). However, none of these effects have been encountered as a result of cigarette smoking.

The concentration of enzymes in the blood reflecting cardiac and liver damage have been reported in patients suffering from acute carbon monoxide poisoning (Ricci et al, 1964; Van Vugt, 1968; Duplay et al, 1967; Afans'ev, 1967; Gramer and Ruof, 1968; Müller and Voigt, 1968; Prellwitz et al, 1970; Antos and Sevcik, 1971).

Four groups of situations are characterized by a similar hematologic response. They are as follows:

1. Exposure to carbon monoxide in man causes an increase in the red cell and reticulocyte counts and the serum globulin fraction (Coscia et al, 1964; Tkachenko et al, 1966; Glass et al, 1968; Kjeldsen and Damgaard, 1968; Bethlenfalvay, 1971a). The erythrocytes manifest some differences from normal erythrocytes in staining (Blackmore, 1970; Bethlenfalvay, 1971b), denaturation of hemoglobin (Perrelli et al, 1970) and a change in the shape of the oxyhemoglobin dissociation curve to the left (Brody and Coburn, 1969).

2. In dogs, exposure to carbon monoxide increases hemoglobin

1005051138

concentration (Asmussen and Vinther-Paulsen, 1949). This change is regarded as an important mechanism for tolerance or acclimatization to carbon monoxide (Otis, 1970). A similar increase in hemoglobin and the red cell count has been noted in other species, such as rabbits (Truhaut et al., 1968), rats (Ramsey, 1969), rhesus monkeys (Theodore et al., 1971), and squirrel monkeys (Jones et al., 1971).

3. Cigarette smoking causes an increase in the red cell count and in the content of these hemoglobin, whereas abstinence caused a fall/in humans (Eisen and Hammond, 1956; Pincherle and Shanks, 1967) and hamsters (Reckzeh and Dontenwill, 1970). In another investigation involving smokers, the elevated hemoglobin and red cell count were correlated with carboxyhemoglobin levels (Petrovic, 1970). A shift in the oxygen dissociation curve to the left has been noted in cigarette smokers (Gutenkauf et al. 1967; Birnstingl et al., 1967).

4. Patients suffering from myocardial ischemia with normal coronary arteriograms have been shown to have abnormal hemoglobin-oxygen dissociation (Likoff et al., 1967; Eliot and Bratt, 1969; Guy et al., 1971). There is no evidence that these patients sustain carboxyhemoglobinemia, nor is the ischemia associated with cigarette smoking.

The above examples are clearly different entities. Cigarette smoking is included only in situation 3 and carbon monoxide in 1 to 3.

1005051139

## BIBLIOGRAPHY

## V. CIRCULATORY SYSTEM

## F. Blood Cells and Plasma

- |  | Reprint |
|--|---------|
| AFANAS'EV B G, ANTONICHIS J L, ARTEM'EV V P and SLUZEVSKY A E:<br>(On the effect of the air medium on the indices of peripheral blood in naval specialists.)<br><u>Voen Zh</u> 3: 60-1, 1967.  | 392     |
| ANTOS V and SEVČIK M: Aktivita enzymu v séru u nemocných s prudkou otravou kysličníkem uhelnatým. (Serum enzyme activity in patients with acute carbon monoxide poisoning.) <u>Vnitr Lek</u> 17: 1151-4, 1971.   | 393     |
| ASMUSSEN E and VINTHER-PAULSEN N V: On the circulatory adaptations to arterial hypoxemia. (Co-poisoning). <u>Acta Physiol Scand</u> 19: 115-24, 1969.  | 394     |
| BETHLENFALVAY N C: Resistance of reticulocytes to oxidation by nitrite: a cytologic demonstration. <u>J Lab Clin Med</u> 77: 361-5, 1971a.   | 395     |
| BETHLENFALVAY N C: Cytologic demonstration of carboxyhemoglobin. Clinical and in vitro studies in man. <u>J Lab Clin Med</u> 77: 543-50, 1971b.  | 396     |
| BIRNSTINGL M, BRINSON K and CHAKRABARTI B K: The effect of short-term exposure to carbon monoxide on platelet stickiness. <u>Brit J Surg</u> 58: 837-9, 1971.  | 397     |
| BIRNSTINGL M, COLE P and HAWKINS L: Variations in oxyhaemoglobin dissociation with age, smoking, and Buerger's disease. <u>Brit J Surg</u> 54: 615-9, 1967.  | 398     |
| BLACKMORE D J: Distribution of HbCO in human erythrocytes following inhalation of CO. <u>Nature</u> 227: 386, 1970.  | 399     |
| BRODY J S and COBURN R F: Carbon monoxide-induced arterial hypoxemia. <u>Science</u> 164: 1297-8, 1969.  | 400     |
| CANDURA F and CRAVERIA A: Valore e significato della fibrinolisi nell'ossicarbonismo sperimentale. (Degree and significance of fibrinolysis in experimental carbon monoxide poisoning.) <u>Rass Med Industr</u> 33: 404-6, 1964.   | 401     |
| COSCIA G C, PERRELLI G, GAIDO P C and CAPELLARO F: Il comportamento del glutathione del glutathione stabile e della glucosio-6-fosfato-deidrogenasi in soggetti esposti ad inalazione cronica d'ossido di carbonio. (The effect of glutathione, stable glutathione and glucose -6-phosphate dehydrogenase in subjects exposed to chronic inhalation of carbon monoxide.) <u>Rass Med Industr</u> 33: 446-51, 1964. | 402     |
| DUPLAY H, ZIEGLER G, GARD A M and PINTO J: Valeur pronostique du dosage des transaminases dans les intoxications par l'oxyde de carbone. 160 observations. (Prognostic value of the measurement of transaminases in carbon monoxide poisoning. 160 cases.) <u>Presse Med</u> 75: 469-72, 1967.   | 403     |
| EISEN M E and HAMMOND E C: The effect of smoking on packed cell volume, red blood cell counts, haemoglobin and platelet counts. <u>Canad Med Ass J</u> 75: 520, 1956.  | 404     |
| EL-ATTAR O A: Effect of carbon monoxide on the whole fibrinolytic activity. <u>Industr Med Surg</u> 37: 774-7, 1968a.  | 405     |
| EL-ATTAR O A: Effect of carbon monoxide on the whole fibrinolytic activity. <u>J Egypt Med Ass</u> 51: 591-7, 1968b.   | 406     |
| EL-EBRASHY N, EL-ASHMAWY S and ALY A: Effect of smoking on the index of platelet adhesiveness and blood glucose level in atherosclerotic patients. <u>J Egypt Med Ass</u> 50/3: 157-68, 1967.  | 407     |

1005051140

- ELIOT R S and BRATT G : The paradox of myocardial ischemia and necrosis in young women with normal coronary arteriograms. Relation to abnormal hemoglobin-oxygen dissociation. Am J Cardiol 23: 633-8, 1969. 408
- GLASS H I, GARRETA A C, LEWIS S M GRAMMATICOS P and SZUR L : Measurement of splenic red-blood-cell mass with radioactive carbon monoxide. Lancet 1: 669-70, 1968. 409
- GRAMER L and ROUF H : Enzym und eiweißveränderungen im Serum bei schwerer akuter Kohlenmonoxidvergiftung. (Enzyme and protein changes in the serum in severe acute carbon monoxide poisoning.) Deutsch Med Wschr 93: 2275-8, 1968. 410
- GUTENKAUF J J, BRATT G T and ELIOT R S : Effect of cigarette smoking on the hemoglobin-oxygen dissociation curve. Circulation 36: Suppl II: 129, 1967. 411
- GUY C R, SALHANY J M and ELIOT R S : Disorders of hemoglobin-oxygen release in ischemic heart disease. Am Heart J 82: 824-32, 1971. 412
- JONES R A, STRICKLAND J A, STUNKARD J A and SIEGEL J : Effects on experimental animals of long-term inhalation exposure to carbon monoxide. Toxicol Appl Pharmacol 19: 46-53, 1971. 413
- KJELDSSEN K and DAMGAARD F : Influence of prolonged carbon monoxide exposure and high altitude on the composition of blood and urine. Scand J Clin Lab Invest 22 Suppl 103: 20-5, 1968. 414
- LIKOFF W, SEGAL B L and KASPARIAN H : Paradox of normal selective coronary arteriograms in patients considered to have unmistakable coronary heart disease. New England J Med 276: 1063-6, 1967. 415
- MULLER H E and VOIGT I M : Quantitativ immunologische Serumproteinbestimmungen bei verschiedenen Krankheitszuständen. Deutsch Med Wschr 93: 216-8, 1968. 416
- OTIS A B : The physiology of carbon monoxide poisoning and evidence for acclimatization. Ann NY Acad Sci 174: 242-5, 1970. 417
- PERRELLI G, PREVOT P A and SULOTTO F : Recenti contributi in tema di patologia enzimatica del globulo rosso nelle intossicazioni professionali. (Recent contributions on enzymatic pathology of the erythrocyte in occupational poisonings.) Minerva Med 61: 2620-5, 1970. 418
- PETROVIC D : Uvecanje broja eritrocita i kolicine hemoglobina u krvi pusaca. (Increase in the number of erythrocytes and in the amount of hemoglobin in the blood of smokers.) Med Glasnik 24: 470-3, 1970. 419
- PINCHERLE G and SHANKS J : Haemoglobin values in business executives. Brit J Prev Soc Med 21: 40-2, 1967. 420
- PRELLWITZ W, SCHUSTER H P, SCHYLLA G, BAUM P, SCHONBORN H, UNGERN-STERNBERG A, BRODERSEN H C and POEPLAU W : Zur differentialdiagnose von Organbeteiligungen bei exogenen Intoxikationen mit Hilfe klinischer und klinisch-chemischer Untersuchungen. (Differential diagnosis of organ involvement in exogenous poisoning by means of clinical and clinico-chemical studies.) Klin Wochschr 48: 51-3, 1970. 421
- RAMSEY J M : The immediate haematological response in the rat to experimental exposures of carbon monoxide. J Physiol 202: 297-304, 1969. 422
- RECKZEH G and DONTENWILL W : Effects of cigarette smoke on hamsters. Arch Environ Health 20: 7-15, 1970. 423
- RICCI C, CAPELLARO F and GAIDO P C : Indagini elettroforetiche ed immunoelettroforetiche sul siero di individui esposti a rischio protratto da CO. (Electrophoretic and immunophoretic studies of the blood of persons exposed to prolonged risk of CO.) Rass Med Industr 33: 414-6, 1964. 424

1005051141

- THEODORE J, O'DONNELL R D and BACK K C : Toxicological evaluation of carbon monoxide in humans and other mammalian species. J Occup Med 13: 242-55, 1971. 425
- TKACHENKO Z A, TISHCHENKO A N, ZATSEPIN A T and DIMITROVA E Z : (External respiration and blood morphology in persons working in conditions with an insignificantly increased concentration of carbon monoxide.) Vrach Delo 12: 78-81, 1966. 426
- TRUHAUT R, BOUDÈNE C and CLAUDE J R : Recherches sur les effets de l'exposition prolongée du lapin et du rat a de très faibles concentrations d'oxyde de carbone. Arch Mal Prof 29: 97-108, 1968. 427
- VAN VUGT H : Enzymveranderingen bij koolmonoxyde-intoxicatie. Ned crl T Geneesk 112: 1993-8, 1968. 428

1005051142

WALD and FENTON 1972 (A 49) 178  
 WALTZ and HAUSERMANN 1965 (141) 31, 28  
 WANSTRUP, KJELDSEN and ASTRUP  
 1969 (390) 73, 70  
 WARBURG, GEISSLER and LORENZ 1967  
 (A 579) 213  
 WATABE 1969 (A 780) 227  
 WATANABE, KITAGUCHI, KIYOFUJI,  
 MORISAKI, MASUDA, NOGUCHI and  
 MATSUMOTO 1970 (A 681) 221  
 WATSON 1968 (A 682) 221  
 WAYLAND and MOHAFER 1971 (A 616) 215  
 WEASE 1967 (A 359) 199  
 WEATHERBURN and LOGAN 1969 (A 50)  
 178  
 WEAVER 1971 (A 313) 196  
 WEBER, MORET and CHAUVET 1967  
 (A 456) 205  
 WEDERKINCH 1964 (A 51) 178  
 WEIGT 1967 (A 683) 221  
 WEINSTOCK 1969 (A 201) 189  
 WEINSTOCK and NIKI 1972 (A 202) 189  
 WEISKOPF and SEVERINGHAUS 1972 (A 457)  
 205  
 WEISS, SLAWSKY and DESFORGES 1971  
 (63) 18, 15; (165) 35, 34  
 WENDER 1963 (A 540) 210  
 WENDT 1941 (269) 55, 51  
 WENNELAND 1945 (64) 18, 14  
 WERNITSCH 1969 (A 844) 232  
 WESTBERG and COHEN 1971 (A 203) 189  
 WHARTON 1964 (A 146) 185  
 WHEREAT 1970 (391) 73, 70  
 WHITE 1970 (A 238) 191  
 WHITE, COBURN, WILLIAMS, GOLDWEIN,  
 ROTIIFER and SHAFER 1967 (A 237) 191  
 WHITEHEAD and WORTHINGTON 1961  
 (93) 23; (166) 35, 33  
 WHO CHRONICLE 1971 (A 268) 193  
 WHO EXPERT COMMITTEE 1963 (A 264) 193  
 WHO EXPERT COMMITTEE 1964 (A 265) 193  
 WHO EXPERT COMMITTEE 1969 (A 267) 193  
 WHO SCIENTIFIC GROUP 1968 (A 266) 193  
 WIECZOREK 1968 (A 52) 178  
 WIETHAUP 1968 (A 315) 196  
 WIJDEVELD 1968 (561) 101, 99  
 WIKTOR 1954 (331) 65, 63  
 WILLIAMS 1964 (192) 41, 37  
 WILSON, RICH and MESSMA 1972 (A 684)  
 221  
 WILSON and HARDING 1970 (613) 110, 108  
 WILSON, NELSON and HARDING 1965  
 (A 580) 213  
 WINTER and SHATIN 1970 (A 845) 232  
 WINTERHALTER, AMICONI and ANTONINI  
 1968 (A 147) 185  
 WITTENBERG, ANTONINI, BRUNORI, NOBLE,  
 WITTENBERG and WYMAN 1967 (A 148)  
 185  
 WITTENBERG, BRUNORI, ANTONINI,  
 WITTENBERG and WYMAN 1965 (633) 112,  
 111  
 WITTCENS 1966 (614) 110, 108  
 WITUSIK 1971 (A 541) 210  
 WOHLERS, NEWSTEIN and DAUNIS 1967  
 (A 727) 224  
 WOHLRAB and OGUNMOLA 1971 (A 581) 213  
 WOJAHN 1967 (A 53) 179  
 WOJCZUK and CHYLAK 1971 (562) 101, 99  
 WOLFGANG 1970 (A 204) 189  
 WOLKONSKY 1969 (A 269) 193; (A 458) 205  
 WOODRUFF 1970 (A 316) 196  
 WOOLF 1964 (A 459) 205  
 WRANNE 1967 (A 164) 186  
 WRANNE 1967 (A 239) 191  
 WRANNE 1969 (A 149) 185  
 WYNDER and HOFFMANN 1967 (34) 10, 7  
 XINTARAS, JOHNSON, ULRICH, TERRILL  
 and SOBECKI 1966 (A 360) 199  
 YACOB, FAURE, MALLON and CAU  
 1970 (65) 18, 13, 15; (167) 35, 34  
 YAGLOU 1955 (108) 26, 24  
 YAMATE and MATSUMURA 1968 (206)  
 46, 45  
 YAMATE and MATSUMURA 1971 (A 317) 196  
 YAMAZAKI, OSHISHI and YAMAZAKI 1970  
 (A 150) 185  
 YASUKOCHI and YASUOKA 1967 (A 542) 210  
 YASUOKA 1970 (550) 98, 96  
 YOUNG and PUGH 1963 1963 (598) 107, 104  
 YOUNOSZAI, KACIC and HAWORTH 1968  
 (596) 106, 104  
 YOUNOSZAI, PELOSO and HAWORTH 1969  
 (597) 107, 103  
 YUKITAKE 1970 (A 543) 210  
 ZAFFIRI 1964 (A 781) 228  
 ZAFFIRI, CALA, CENTI and SALICONE 1971  
 (A 782) 228  
 ZANARDI, VILLA and MONTI 1964 (270)  
 55, 52  
 ZARIVAISKALA 1966 (A 728) 224  
 ZEH 1960 (332) 65, 63  
 ZENK 1964 (477) 87, 85  
 ZENK 1965 (478) 87, 85  
 ZIBEROV 1966 (A 783) 228  
 ZIEGLER 1936 (308) 60, 56  
 ZOLOTUKHIN 1968 (457) 84, 81  
 ZORN 1964 (A 685) 221  
 ZORN 1968 (A 846) 232  
 ZORN 1969 (A 54) 179

1005051313



## BIBLIOGRAPHY

## VI. NERVOUS SYSTEM

A. Cerebral Vessels

- BREITENECKER L: Hirnblutung und Kohlenoxydvergiftung. (Cerebral hemorrhage and carbon monoxide poisoning. Wien Klin Wochschr 51: 217, 1938. Reprint 429
- BRETON J, CAROFF J, MARTIN R, DEHOUE A and DEHOUE P: Obstruction mortelle du tronc basilaire consécutive à une intoxication oxycarbonée bénigne. (Fatal obstruction of the basilar trunk following benign carbon monoxide poisoning.) Med Leg Domm Cor-por Paris 2: 409-11, 1969. 430
- FODOR G G: Tierexperimentelle Untersuchungen über die Auswirkung von Kohlenoxidvergiftungen bei Beeinträchtigung des Hirnkreislaufs durch Carotisligatur. (Experimental investigations on the effect of carbon monoxide poisoning on the cerebral circulation impaired by carotid ligation.) Naunyn Schmiedeberg Arch Pharm 264: 229-30, 1969. 431
- GLOWACKI B, GRUDZINSKA B and WACLAWIK P: Ostre zaburzenia naczyniowe mozgu po zatruciu CO. (Acute disturbances of the cerebral blood vessels in carbon monoxide poisoning.) Prezeg Lek 14: 306-8, 1958. 432
- HALL G H: Effects of nicotine, carbon monoxide and tobacco smoke on regional blood flow in the cerebral cortex. Europ J Pharmacol 19: 385-8, 1972. 433
- MAURER F W: The effects of anoxemia due to carbon monoxide and low oxygen on cerebrospinal fluid pressure. Am J Physiol 133: 180-8, 1941. 434
- SAITA G and LUSSANA S: Sindrome coronaro-cerebrale tardiva da intossicazione ossicarbonica acuta. (Late coronary-cerebral syndrome caused by acute carbon monoxide poisoning.) Med Lav 62: 185-95, 1971. 435
- SJOSTRAND T: Brain volume, diameter of the blood-vessels in the pia mater, and intracranial pressure in acute carbon monoxide poisoning. Acta Physiol Scand 15: 351-61, 1948. 436
- WAJGT A: Zmiany naczyniopochodne w przypadku leczenia elektrowstrzasem zespołu psychicznego po zatruciu tlenkiem węgla. (Cerebral vascular lesions after use of electroshock in a case of psychotic syndrome following carbon monoxide intoxication.) Neurol Neurochir Pol 5: 587-90, 1971. 437

1005051144

## VI B. Eye and Visual Pathways

Visual threshold is influenced by administration of carbon monoxide (see review by McFarland, 1970). McFarland et al. (1944) used a visual discriminometer and noted a change in visual sensitivity following exposure to 0.01% carbon monoxide. There was a fall in threshold with a rise of 5% in carboxyhemoglobin level in 2 subjects. Lilienthal and Fugitt (1946) noted in 5 subjects that increments of 5 to 10% carboxyhemoglobin impaired the critical flicker fusion frequency. Halperin et al. (1959) observed the recovery from the effects of carbon monoxide on visual function. Administration of pure oxygen resulted in an immediate improvement, and the addition of 7% carbon dioxide to the oxygen hastened the elimination of carbon monoxide from the body, compared with the inhalation of air.

In patients suffering from carbon monoxide poisoning the following effects have been reported: change in optically evoked response in the electroencephalogram (Helmchen and Kunkel, 1964; Hosko, 1970); impairment of visual perception relating to lesions in the occipital lobe (Kuroiwa et al., 1967; Szliwowski and Klees-Delange, 1970), visual agnosia (Zolotukhin, 1968; Benson and Greenberg, 1969), retinopathy (Heydenreich, 1970; Bilchik et al., 1971) and conjunctival hemorrhages (Prokop and Wabnitz, 1970).

In laboratory animals inhalation of carbon monoxide leads to lesions which relate to those observed in human poisoning. In rabbits there is sludging of blood in the ophthalmic vessels (Taccola et al., 1965). Diminution of evoked potentials in the visual pathways occurs in cats (Ikeda, 1969) and in

1005051145

## Additional Bibliography List No. 17

## THERAPY OF CARBON MONOXIDE POISONING

- AQUINAS M : Coal gas poisoning: a nursing care study. Nurs Times 60: 1208-10, 1964. Reprint A 729
- BEGHE' RAOUL : Esperienze in tema di trattamento dell'ossicarbonismo acuto. (Experience in the treatment of acute carbon monoxide poisoning). Rass Med Industr 33: 337-48, 1964. A 730
- BELAÏSCH J : La plasmathérapie. (Plasmatherapy.) Vie Med Paris 35: 999-1002, 1954. A 731
- BORBÉLY F : Die behandlung der kohlenoxydvergiftungen. (Treatment of carbon monoxide poisoning). Deutsch Med Wschr 90: 1963-4, 1965. A 732
- BOULETREAU P and MOTIN J : Etude d'une série homogène de 50 comas oxycarbonés traités par l'oxygénothérapie hyperbare. (Study of a homogeneous series of 50 cases of coma due to carbon monoxide poisoning and treated with hyperbaric oxygenation). Moroc Med 50: 386, 1970. A 733
- BOUR H, PASQUIER P and BERTRAND-HARDY J M : Le coma oxycarboné. Étude générale, clinique, biologique et thérapeutique de 290 cas. (Carbon monoxide coma. General study, clinical, biological and therapeutics of 290 cases). Sem Hop Paris 42: 1839-61, 1966. A 734
- BRANDON S : Treatment of carbon monoxide poisoning. Lancet 1: 626, 1970. A 735
- BURMEISTER H, BARCKOW D, HUMPERT U, IBE K and LERCHE D : Die künstliche beatmung. Ein erfahrungsbericht. (Artificial respiration. Report based on personal experience). Deutsch Med Wschr 93: 517-22, 1968. A 736
- BURMEISTER H and HEUHAU G A : Die behandlung der schweren subakuten leuchtgasvergiftung beim menschen. (Treatment of severe, subacute natural gas poisoning in humans). Arch Toxikol 26: 277-92, 1970. A 737
- CARUSO G and BARNABA A : Trattamento con diazepam e remissione della sintomatologia extrapiramidale in un caso di parkinsonismo da intossicazione con ossido di carbonio. (Treatment with diazepam and remission of the extrapyramidal symptoms in a case of parkinsonism caused by carbon monoxide poisoning). Acta Neurol Napoli 23: 103-10, 1968. A 738
- CENTI R and ZAFFIRI O : Considerazioni su di un caso gravissimo di ossicarbonismo acuto favorevolmente risolto con acido ascorbico ad alte dosi e con L-dopa. (On a very grave case of acute carbon monoxide poisoning favorably resolved with high doses of ascorbic acid with L-dopa). Minerva Anestesiol 37: 406-14, 1971. A 739
- CHUKHRIENKO D P and LULKO A V : (Extrarenal hemodialysis in carbon monoxide poisoning). Vrach Delo 12: 18-21, 1968. A 740
- CIOCATTO E and PATTONO R : Organizzazione di un centro di rianimazione respiratoria terapia dell'intossicazione acuta da ossido di carbonio. (Organization of a respiratory resuscitation center. Therapy of acute carbon monoxide poisoning). Rass Med Industr 33: 330-3, 1964. A 741
- COLLINS J V and GOULDING R : Treatment of acute poisoning at Guy's Hospital: October 1969 to September 1970. Guys Hosp Rep 120: 31-46, 1971. A 742
- DAMIA G, SIBILLA E and DAMIA G : Considerazioni sulla prognosi e trattamento di alcuni casi di avvelenamento da barbiturici e da CO. (Considerations on the prognosis and treatment of some cases of poisoning due to barbiturates and carbon monoxide). Minerva Anest 31: 564-8, 1965. A 743

1005051289

## VI. NERVOUS SYSTEM

## B. Eye and Visual Pathways

- |   | Reprint |
|---|---------|
| BENSON D F and GREENBERG J P : Visual form agnosia. <u>Arch Neurol</u> 20: 82-9, 1969.  | 438     |
| BILCHIK R C, MULLER-BERGH H A and FRESHMAN M E : Ischemic retinopathy due to carbon monoxide poisoning. <u>Arch Ophthal</u> 86: 142-4, 1971.  | 439     |
| HALPERIN M H, McFARLAND R A, NIVEN J I and ROUGHTON F J W : The time course of the effects of carbon monoxide on visual thresholds. <u>J Physiol</u> 146: 583-93, 1959.   | 440     |
| HELMCHEN H and KÜNKEL H : Befunde zur rhythmischen Nachschwankung bei optisch ausgelösten Reizantworten (evoked responses) im EEG des Menschen. (Findings on rhythmic potential fluctuation in optically evoked responses in the human EEG.) <u>Arch Psychiat Nervenkr</u> 205: 397-408, 1964.          | 441     |
| HEYDENREICH A : Erkrankungen des Auges in Industrie und Beruf. (Eye diseases in industry and professions.) <u>Z Aerztl Fortbild Jena</u> 64: 918-21, 1970.  | 442     |
| HOSKO M J : The effect of carbon monoxide on the visual evoked response in man. <u>Arch Environ Health</u> 21: 174-80, 1970.  | 443     |
| IKEDA T : Experimental carbon monoxide poisoning: its electrophysiological effects on the visual pathway in cats. <u>Fol Psychiat Neurol Jap</u> 23: 135-42, 1969.  | 444     |
| KATSURA T : (Effect of carbon monoxide on the evoked potential of the visual neurons.) <u>Nippon Ika Dai Zasshi</u> 38: 69-76, 1971.  | 445     |
| KUROIWA Y, SHIDA K, NAGAMATSU K, KATO M and SANTA T : Involvement of cerebral functions in acute carbon monoxide poisoning with special reference to occipital lobe functions. <u>Fol Psychiat Neurol Jap</u> 21: 189-97, 1967.   | 446     |
| LILIENTHAL J L Jr. and FUGITT C H : The effect of low concentrations of carboxyhemoglobin on the 'altitude tolerance' of man. <u>Am J Physiol</u> 145: 359-64, 1946.  | 447     |
| McFARLAND R A : The effects of exposure to small quantities of carbon monoxide on vision. <u>Ann NY Acad Sci</u> 174: 301-12, 1970.   | 448     |
| McFARLAND R A, ROUGHTON F J W, HALPERIN M H and NIVEN J I : The effects of carbon monoxide and altitude on visual thresholds. <u>J Aviation Med</u> 15: 381-94, 1944.   | 449     |
| MELANOWSKI W H : Wpływ palenia tytoniu na narząd wzroku człowieka. (The influence of tobacco smoking on human sight.) <u>Pol Tyg Lek</u> 18: 1937-9, 1963.  | 450     |
| NIEBROJ T : Wpływ zatrucia tlenkiem węgla na zmiany cytochemiczne w siatkówce świnki morskiej. (Effect of intoxication with carbon monoxide on the cytochemical changes in the retina of guinea pigs.) <u>T Klin Oczna</u> 39: 7-11, 1969.  | 451     |
| OHARA M : (The electro microscopic studies on the optic nerves of mice poisoned with carbon monoxide gas or potassium cyanide solution.) <u>Folia Ophthal Jap</u> 19: 99-123, 1968.   | 452     |
| PROKOP O and WABNITZ R : Vorkommen von Bindehautblutungen bei Lebenden und Toten, dargestellt in 10 Tabellen. (Incidence of conjunctival hemorrhage in the living and dead presented in 10 tablets.) <u>Z Reichtmed</u> 67: 249-57, 1970.   | 453     |
| SCHMIDT B : Einfluß des Zigarettenrauchens auf das EOG. (Effect of cigarette smoking in the EOG.) <u>Klin Monatsbl Augenheilkd</u> 156: 523-31, 1970.   | 454     |
| SZLIWOWSKI H B and KLEES-DELANGE M : Aspects visuomoteurs de la perception: mise au point théorique et application aux intoxications oxycarbonées. (Viso-motor aspects of perception: a theoretical discussion and application to carbon monoxide poisoning.) <u>Acta Pediatr Belg</u> 24: 295-9, 1970. | 455     |

1005051147

TACCOLA A, JEDRYCHOWKI W and CAVALLERI A : La microcircolazione nella congiuntiva bulbare dopo intossicazione acuta, sperimentale, da CO. (The microcirculation in the bulbar conjunctiva after experimental acute CO poisoning.) Folla Med Napoli 48: 1023-33, 1965.

456

ZOLOTUKHIN A N : (On the combined effect of small concentrations of carbon monoxide and hypoxia on the visual analyzer.) Voen Med Zh 4:60-4, 1968.

457

1005051148

## VI C. Hearing and Auditory Pathways

The auditory and vestibular apparatus is less sensitive than the eye to carboxyhemoglobin. Tibbling (1969) tested 25 human volunteers by the smoking of cigarettes. A rise in the blood level of carboxyhemoglobin to 2.5 % did not influence vestibular nystagmus. Guest et al. (1970) administered carbon monoxide to produce levels of 10 % carboxyhemoglobin in 8 subjects. No depression of the auditory flutter fusion threshold or of the critical flicker fusion threshold occurred. In the same subjects the oral administration of phenobarbital depressed both thresholds.

Patients suffering from acute carbon monoxide poisoning develop the following disorders relating to the ear: abnormalities in hearing (Taniewski and Kugler, 1964a; Hansz and Styperek, 1968; Kawamura, 1971); deafness (Morris, 1969; Fortunato and Catalano, 1970); distortion of the audiometric curve (Taniewski and Kugler, 1964b; Fritzsche, 1969); and disturbance in cochleovestibular function (Cis and Perani, 1964; Mounier-Kuhn et al., 1968). Industrial workers exposed chronically to carbon monoxide develop similar abnormalities in hearing and vestibular function (Stanković et al., 1964; Strzelczyk and Zenk, 1964; Zenk, 1964, 1965; Sato, 1966; Mesolella et al., 1970).

In experimental animals the administration of carbon monoxide causes interference with the auditory and olfactory systems. There is depression of action potentials in the olfactory bulb of the cat (Hall, 1970) and in the cochlea of the guinea pig (Freigang et al., 1968). The morphologic examination of the inner ear shows degeneration in the rabbit (Küttner, 1968) and in other animal species (Kittel and Theissing-Erlangen).

1005051149

## VI. NERVOUS SYSTEM

## C. Hearing and Auditory Pathways

- CIS C and PERANI G: La funzionalita cocleo-vestibolare nell'ossicarbonismo. (Cochleo-vestibular function in carbon monoxide poisoning.) Arch Ital Otol 75: 635-43, 1964. 458 Reprint
- FORTUNATO V and CATALANO G B: Ipoacusie e sordita da intossicazioni esogene. (Bradyacusia and deafness caused by exogenous intoxications.) Minerva Otorinolaringol 20: 193-217, 1970. 459
- FREIGANG B, SEIDEL P and FLACH M: Zur Frage der Kohlenmonoxydauswirkung auf die Mikrophon- und Summenaktionspotentiale der Meerschweinchencochlea. (On the problem of carbon monoxide effect on the microphonics and sum of action potentials of the guinea pig cochlea.) Arch Klin Exp Ohr Nas Kehlk 190: 24-35, 1968. 460
- FRITSCH F: Richtungsaudiometrische Untersuchungen im freien Schallfeld bei U-förmig verlaufenden Horschwellenkurven. (Directional audiometric studies in a free field on patients with U-shaped audiograms.) Z Lar Rhin Otol 48: 291-302, 1969. 461
- GUEST A D L, DUNCAN C and LAWTHORPE P J: Carbon monoxide and phenobarbitone: A comparison of effects on auditory flutter fusion threshold and critical flicker fusion threshold. Ergonomics 13: 587-94, 1970. 462
- HALL G H: Effects of nicotine and tobacco smoke on the electrical activity of the cerebral cortex and olfactory bulb. Brit J Pharmacol 38: 271-86, 1970. 463
- HANSZ J and STYPEREK J: Jednostronne uszkodzenie nerwu sluchowego w ostrym zatruciu tlenkiem wegla. (Unilateral damage of the acoustic nerve during acute carbon monoxide poisoning.) Pol Tyg Lek 23: 1441-2, 1968. 464
- KAWAMURA S: (Clinical observation of tinnitus caused by extrinsic causes - head trauma, acoustic trauma, etc.) J Otol Jap 74: 1007-27, 1971. 465
- KITTEL G and THEISSING-ERLANGEN G: Histologische Untersuchungen der Cochlea an Häutchen-Präparaten und Treppen-Serienschnitten nach hochgradiger protrahierter Hypoxydase. (Histological studies of the cochlea on cuticle preparations and scaliform sections following severe protracted hypoxia.) Arch Klin Exp Ohr Nas Kehlk 191: 534-8, 1968. 466
- KÜTTNER K: Zur Pathomorphologie der Veränderungen am peripheren Hörorgan bei wiederholter experimenteller Kohlenmonoxydintoxikation. (On the pathomorphology of changes in the peripheral organ of hearing during repeated experimental carbon monoxide intoxication.) Z Laryng Rhinol Otol 47: 779-85, 1968. 467
- MESOLELLA C, PERRELLA F, TESTA B and MORELLI G: Rilievi oto-neurologici in un gruppo di operai esposti cronicamente all'azione dell'ossido di carbonio. (Otoneurological observations in a group of workmen chronically exposed to the action of carbon monoxide.) Arch Ital Laring 78: 47-60, 1970. 468
- MORRIS T M O: Deafness following acute carbon monoxide poisoning. Laryng 83: 1219-25, 1969. 469
- MOUNIER-KUHN P, ROCHE L, MORGON A and BERNARD P: Les atteintes vestibulaires au décours immédiat des intoxications aiguës par l'oxyde de carbone. (Vestibular involvement immediately following acute carbon monoxide poisoning.) J Franc Otorhinolaryng 17: 512-5, 1968. 470
- SATO T: (Hearing disturbances in monoxide-gas toxicosis.) Otolaryngology / (Tokyo) 38: 805-16, 1966. 471
- STANKOVIĆ D, KANTA F, FOCO S and ALJINOVIC M: Oštećenja sluha posle ponovljenih akutnih trovanja ugljen-monoksidom. (Hearing disorders following repeated carbon monoxide poisoning.) Acta Med Jugosl 18: 95-106, 1964. 472

1005051150

- STRZELCZYK P and ZENK H : Permanente subtoxische CO-Einwirkungen auf das Hör- und Gleichgewichtsorgan bei Gaswerkarbeitern. (Permanent subtoxic CO effects on the hearing and equilibrium apparatus in gas plant workers.) Arch Ohr Nas Kehlk 184: 81-92, 1964. 473
- TANIEWSKI J and KUGLER R : Uposledzenie sluchu w zatruciu tlenkiem wegla. (Hearing disorders in carbon monoxide poisoning.) Otolaryng Pol 18: 493-7, 1964a. 474
- TANIEWSKI J and KUGLER R : Audiometrische Kurve in U-form bei Kohlenoxydvergiftungen. (Audiometric curve in U form in carbon monoxide poisoning.) Msschr Ohrenheilk 98: 298-301, 1964b. 475
- TIBBLING L : The influence of tobacco smoking, nicotine, CO and CO<sub>2</sub> on vestibular nystagmus. Acta Otolaryng 68: 118-26, 1969. 476
- ZENK H : CO-Intoxikationen in der otologisch-arbeitsmedizinischen Gutachterpraxis. (CO poisoning in otological industrial medical expert testimony practice.) Int Arch Gewerbepath 20: 432-42, 1964. 477
- ZENK H : Die Auswirkungen berufsbedingter CO-Intoxikationen auf Geruchs-, Gehör- und Gleichgewichtsorgan. (Effects of occupationally caused CO intoxications on the organs of smell, hearing and equilibrium.) Z Laryng Rhinol Otol 44: 821-8, 1965. 478

1005051151



VI D. Behavior of Man

The effects of carbon monoxide on human behavior have been summarized by McFarland (1952), Beard and Grandstaff (1970) and LaVerne (1970). Several reports have appeared which determine the blood level of carboxyhemoglobin that would influence functions of the higher centers of the central nervous system in man. The results are as follows:

1. Sayers et al, (1929) summarized the findings of an investigation in which 6 men were exposed for 4 to 7 hours daily over a period of 68 days to gasoline engine exhaust gas-air mixtures. Exposure to 200 ppm carbon monoxide, resulting in blood levels of 20% carboxyhemoglobin, caused frontal headache in 3-1/2 to 4 hours. Exposure to 300 ppm, producing a blood level of 30%, caused frontal and occipital headaches and vertigo after 3 hours. Nielsen (1971) in another group of subjects with 25% to 33% carboxyhemoglobin noted an elevation of body temperature.

2. The control precision and multiple limb coordination are impaired when the concentration of carboxyhemoglobin in the blood exceeds 5% (Trouton and Eysenck, 1961).

3. Psychomotor abilities are sensitive to the presence of carboxyhemoglobin. Ability to discriminate is altered at levels below 5%, while reaction time, static steadiness and muscle persistence are measurably altered by concentrations of up to 20% (Schulte, 1963). There was no difference between the test results in nonsmokers and those in smokers.

4. The ability of subjects to estimate time intervals of 1, 3 and 5 seconds was not impaired by exposures to as much as 20% carboxyhemoglobin.

1005051152

Levels between 20 and 30 % caused interference with visual evoked response and impairment of manual dexterity (Stewart et al., 1970).

5. Estimation of time and tracking performance was not influenced by exposure to 125 ppm carbon monoxide with a mean of 6.64 % carboxyhemoglobin (Hanks, 1970; Mikulka et al., 1970; O'Donnell et al., 1970, 1971 a and b).

6. Levels of carboxyhemoglobin below 10 % influence the results of tests for automobile driving performance (Ray and Rockwell, 1970).

7. Levels of 6.6 % carboxyhemoglobin caused a reduction in the ability of the subjects to identify signals. This effect on vigilance did not appear with 2.3 % carboxyhemoglobin (Horvath et al., 1971).

8. Exposure of students to 100 ppm carbon monoxide for 2 hr, with mean levels of 7.2 % carboxyhemoglobin, resulted in diminution of manual dexterity, visual perception and ability to learn and perform certain intellectual tasks (Bender et al., 1971). There was no discernible difference between smokers and nonsmokers with regard to their psychological susceptibility.

9. Ramsey (1972) exposed 20 normal subjects and 20 patients with pulmonary emphysema and 20 with anemia to inhalation of 0.03 % carbon monoxide in air for 40 min. The mean increase of 4.5 % in carboxyhemoglobin caused a diminution in the speed of reaction to a visual stimulus but no significant change in tests for depth perception and for visual discrimination of brightness.

10. The only observations relating to cigarette smoking pertain to high altitude. Kratochvil et al. (1957) tested smoking at a level of 18,000 feet. There was only a slight decrement in performance of subjects when smoking/at high altitude compared with that of the hypoxic controls. Astrup et al. (1971)

1005051153

reported on a patient who suffered from mountain sickness over many years.

The attacks were provoked either by cigarette smoking or by inhalation of 0.5% carbon monoxide. There are no other reports of such patients with mountain sickness induced by tobacco smoking. It is possible that these responses may relate to changes in cerebral blood flow described by Demange and Auzas (1969).

1005051154

## VI. NERVOUS SYSTEM

## D. Behavior of Man

Reprint

- ASTRUP P, KJELDSEN K and SIGGAARD-ANDERSEN J: Carbon monoxide-induced mountain sickness provoked by tobacco smoking. Lancet 1: 781-2, 1971. 479
- BEARD R and GRANDSTAFF N: Carbon monoxide exposure and cerebral function. Ann NY Acad Sci 174: 385-95, 1970. 480
- BENDER W, GOTHERT M, MALORNY G and SEBBESSE P: Wirkungsbild niedriger Kohlenoxid-Konzentrationen beim Menschen. (Effects of low carbon monoxide concentration in man.) Arch Toxikol 27: 142-58, 1971. 481
- DEMANGE J and AUZAS A: Action de l'altitude chez le fumeur et le non fumeur sur un index de débit sanguin cérébral. Rev Med Aeron Spat 8: 125-7, 1969. 482
- HANKS T G: Human performance of a psychomotor test as a function of exposure to carbon monoxide. Ann NY Acad Sci 174: 421-4, 1970. 483
- HORVATH S M, DAHMS T E and O'HANLON J F: Carbon monoxide and human vigilance. Arch Environ Health 23: 343-7, 1971. 484
- KRATOCHVIL C H, WILKS S S and GERRARD W A III: Cigarette smoking at altitude. Fed Proc Balt 16: 75, 1957. 485
- LAVERNE A A: Air pollution, healing, and civilization. Behav Neuropsychiat 2: 26-37, 1970. 486
- McFARLAND R A: Anoxia: Its effects on the physiology and biochemistry of the brain and on behavior. The Biology of Mental Health and Disease, Paul B. Hoeber, Inc., New York, 1952, pp 335-55. 487
- O'DONNELL R, HEINIG P and THEODORE J: The effect of carbon monoxide on human performance. Ann NY Acad Sci 174: 409-20, 1970. 488
- NIELSEN B: Thermoregulation during work in carbon monoxide poisoning. Acta Physiol Scand 82: 98-106, 1971. 489
- O'DONNELL R D, CHIKOS P and THEODORE J: Effect of carbon monoxide exposure on human sleep and psychomotor performance. J Appl Physiol 31: 513-8, 1971. 490
- O'DONNELL R D, MIKULKA P, HEINIG P and THEODORE J: Low level carbon monoxide exposure and human psychomotor performance. Toxicol Appl Pharmacol 18: 593-602, 1971b. 491
- HEINIG P and THEODORE J
- O'DONNELL R D, MIKULKA P: Effects of short-term low level carbon monoxide exposure on human performance. US Department of Defense, Aerospace Medical Research Laboratory, Ohio, AD-717716, 52 pages, 1970. 492
- RAMSEY J M: Carbon monoxide, tissue hypoxia and sensory psychomotor response in hypoxaemic subjects. Clin Sci 42: 619-25, 1972. 493
- RAY A M and ROCKWELL T H: An exploratory study of automobile driving performance under the influence of low levels of carboxyhemoglobin. Ann NY Acad Sci 174: 396-408, 1970. 494
- SAYERS R R, YANT W P, LEVY E and FULTON W B: Effect of repeated daily exposure of several hours to small amount of automobile exhaust gas. Public Health Bulletin No. 186 Washington, D. C., 1-57, 1929. 495

1005051155

SCHULTE J H : Effects of mild carbon monoxide intoxication. Arch Environ Health 7:  
524-30, 1963.

496

Q STEWART R D, PETERSON J E, BARETTA E D, BACHAND R T, HOSKO M J and HERR-  
MANN A A : Experimental human exposure to carbon monoxide. Arch Environ Health  
21: 154-64, 1970.

497

TROUTON D and EYSENCK H J : The effects of drugs on behavior. Handbook of Abnormal  
Psychology, H. J. Eysenck, Editor, Basic Books Inc., New York: 634-96, 1961.

1005051156

VI. E. Behavior of Animals

The effects of carbon monoxide on the behavior of certain animals have been described, viz., on that of rats (Beard and Wertheim, 1967; Goldberg and Chappell, 1967; Teichner, 1967; Stupfel et al., 1968; Khachaturiyan et al., 1969; Rose et al., 1970; Stupfel and Bouley, 1970; Nečas and Neuwirt, 1971), mice (Hirata et al., 1969; Gaume et al., 1971), dogs (Carding, 1968; Preziosi et al., 1970), monkeys (Back, 1969), and laboratory animals (Fazekas, 1967). The morphologic and functional features of the brain have been described, resulting from exposure of various animals to carbon monoxide, viz., rats (Thomas and Pearse, 1964; Miyagishi and Hiyashi, 1968; Miyagishi and Suwa, 1969; Takahata and Miyagishi, 1969), mice (Estler et al., 1971A), guinea pigs (Kupfer and Wunscher, 1968), cats (Ando et al., 1969; Barrios et al., 1969), birds (Dugnat, 1965), and frogs (Segal, 1970). The brain exposed to carbon monoxide shows a reduction in content of gangliosides (Mawatari, 1970), accumulation of ammonia (Okunyeu and Prokhorenko, 1965; Mishchenko and Frenkel, 1966), decrease in catecholamines (Marks and Swiecicki, 1971); decrease in serotonin (Pare et al., 1969), and decrease in glycogen (Estler et al., 1969, 1971b). The relationship of these results to the behavior pattern has not been elucidated.

1005051157

## BIBLIOGRAPHY

Page 94

## VI. NERVOUS SYSTEM

## E. Behavior of Animals

- ANDO S, SEINO S and HAGIWARA I: Experimental studies on cat's brains with lesions induced by carbon monoxide poisoning after the ligation of unilateral common carotid artery. Adv Neurol Sci (Tokyo) 13: 56-62, 1969. Reprint 498
- BACK, K C: Effects of carbon monoxide on the performance of monkeys. Aerospace Med Div Bact AMRL-TR-69-130 Paper No. 3:41-51, 1969. 499
- BARRIOS P, KOLL W and MALORNY G: Rückenmarksreflexe und afferente Nervenleitung der Katze unter dem Einfluß von Kohlenmonoxyd. (Effect of carbon monoxide on spinal cord reflexes and afferent nerve conduction in cats.) Naunyn Schm Arch Pharm 264: 1-17, 1969. 500
- BEARD R R and WERTHEIM G A: Behavioral impairment associated with small doses of carbon monoxide. Am J Pub Health 57: 2012-22, 1967. 501
- CARDING A H: Mass euthanasia of dogs with carbon monoxide and/or carbon dioxide; preliminary trials. J Small Anim Pract 9: 245-59, 1968. 502
- DUGNAT J M: Action du gaz d'éclairage et du gaz de Lacq sur le système nerveux de l'embryon d'Oiseau. (Action of illuminating gas on the nervous system of the bird embryo.) C R Soc Biol Paris 159: 1545-7, 1965. 503
- ESTLER C J, AMMON H P T and ZIMMERMANN V: Katecholaminunabhängige Abnahme des Glykogengehalts im Gehirn bei Vergiftung mit Kohlenmonoxyd. (Reduction in cerebral glycogen content after carbon monoxide poisoning independent of catecholamines. Naunyn Schm Arch Pharm 263: 204-5, 1969. 504
- ESTLER C J, AMMON H P T, and HEIM F: Metaboliten des Kohlenhydratstoffwechsels und energiereiche Phosphate im Gehirn weisser Mäuse nach wiederholter Kohlenmonoxydvergiftung. (Metabolites of carbohydrate metabolism and high-energy phosphates in the brain of mice after repeated carbon monoxide poisoning.) Arch Toxikol 27:24-8, 1971a. 505
- ESTLER C J, HEIM F, AMMON H P T and ZIMMERMANN V: Catecholamine-independent Glycogenolysis in brain during carbon monoxide poisoning. Pharmacology 5: 55-63, 1971b. 506
- FAZEKAS I G: Aethylalkohol és szénmonoxidmérgezés valamint ezek együttes hatása emberekre és állat-kísérletekben. (Poisoning by ethyl alcohol and carbon monoxide, their combined effect on humans and experimental animals.) Orv Hetil 108: 1503-6, 1967. 507
- GAUME J G, BARTEK P and ROSTAMI H J: Experimental results on time of useful function (TUF) after exposure to mixtures of serious contaminants. Aerospace Med 42: 987-90, 1971. 508
- GOLDBERG H D and CHAPPELL M N: Behavioral measure of effect of carbon monoxide on rats. Arch Environ Health 14: 671-7, 1967. 509
- HIRATA M, HIOKI A and HASHIMOTO K: Distribution of death rate in acute carbon monoxide intoxication in mice. Tohoku J Exp Med 97: 67-73, 1969. 510
- KHACHATURYAN M K, MITAREVSKAYA E V and EGORENKOVA G A: (The effect of chronic inhalation poisoning of experimental animals with phenol in combination with carbon monoxide on functioning of the central nervous system.) Gig Sanit 34: 22-5, 1969. 511
- KUPFER M and WÜNSCHER W: Experimentelle allergische Encephalomyelitis bei CO-geschädigten Meerschweinchen. (Experimental allergic encephalomyelitis in the carbon monoxide exposed guinea pig.) Zbl Allg Path 111: 313-24, 1968. 512

1005051158

- MARKS E and SWIECICKI W : Wplyw tlenu wegla, wibracji i obnizonego cisnienia atmosferycznego na mozgowy uklad neurosekrecyjny i poziom aminokatecholil (Effect of carbon monoxide, vibration and lowered atmospheric pressure on the cerebral neuro-secretory system and catecholamine levels.) Med Pracy 22: 335-42, 1971. 513
- MAWATARI S : Biochemical study on rat brain in acute carbon monoxide poisoning. Fol Psychiat Neurol Jap 24: 123-9, 1970. 514
- MISHCHENKO L I and FRENKEL S R : (Some conversions of glutaminic acid in rat brain tissue on disturbance of its function owing to carbon monoxide intoxication.) Ukr Biokhim Zh 38: 555-9, 1966. 515
- MIYAGISHI T and SUWA N : Electron microscopic studies on the cerebral lesions of rats in experimental carbon monoxide poisoning. Acta Neuropath Berlin 14: 118-25, 1969. 516
- MIYAGISHI T and HAYASHI H : (Electron microscopic studies on the cerebral lesions of rats with experimental carbon monoxide poisoning.) Psychiat Neurol Jap 70: 812-24, 1968. 517
- and
- NEČAS E/ NEUWIRT F : Jednoduché zarizení na vyvolání otravy kyslíčnikem uhelnatým u malých laboratorních zvířat. (Simple technic for inducing carbon monoxide poisoning in small laboratory animals.) Cesk Fysiol 20: 279-81, 1971. 518
- OKUNYEV V N and PROKHORENKO L G : (Some peculiarities of nitrogen metabolism in the rat brain on intoxication with detonating gases.) Ukr Biokhim Zh 38: 469-71, 1966. 519
- PRICE K and STACEY R S :
- PARE C M B and YEUNG D P H/ 5-Hydroxytryptamine, noradrenaline, and dopamine in brainstem, hypothalamus, and caudate nucleus of controls and of patients committing suicide by coal-gas poisoning. Lancet 2: 133-5, 1969. 520
- PREZIOSI T J, LINDENBERG R, LEVY D and CHRISTENSON M : An experimental investigation in animals of the functional and morphologic effects of single and repeated exposures to high and low concentrations of carbon monoxide. Ann NY Acad Sci 174: 369-84, 1970. 521
- ROSE C S, JONES R A, JENKINS L J Jr and SIEGEL J : The acute hyperbaric toxicity of carbon monoxide. Toxicol Appl Pharmacol 17: 752-60, 1970. 522
- SEGAL J R : Metabolic dependence of resting and action potentials of frog nerve. Am J Physiol 219: 1216-25, 1970. 523
- STUPFEL M, BOULEY G, DEKOV S, BOURGEOIS M and ROUSSEL A : Étude expérimentale sur le rat et la souris des effets d'une exposition pendant un à trois mois à 15 et 50 ppm d'oxyde de carbone. (An experimental study in the rat and the mouse on the effects of a 3-month exposure to carbon monoxide at 15 and 50 ppm.) Bull Inst Nat Sante 23: 309-51, 1968. 524
- STUPFEL M and BOULEY G : Physiological and biochemical effects on rats and mice exposed to small concentrations of carbon monoxide for long periods. Ann NY Acad Sci 174: 342-68, 1970. 525
- TAKAHATA N and MIYAGISHI T : (An electromicroscopic study of carbon monoxide poisoning.) Adv Neurol Sci Tokyo 13: 63-8, 1969. 527
- TEICHNER W H : An exploration of some behavioral techniques for toxicity testing. J Psychol 65: 69-90, 1967. 528
- THOMAS E and PEARSE A G E : The solitary active cells. Histochemical demonstration of damage-resistant nerve cells with a TPN-Diaphorase reaction. Acta Neuropath 3: 238-49, 1964. 529

1005051159



## VI F. Electroencephalogram in Man and Animals

All of the available information regarding the effects of carbon monoxide on the electroencephalogram relates to high concentration of the gas in the blood. In rabbits, levels of 45% and 74% carboxyhemoglobin were used respectively by Komura (1967) and De Valois and Schadé (1967). With these high concentrations there are various effects: shortening of the hippocampal electroencephalogram, appearance of slow waves in the neocortex, hypothalamus and caudate nucleus, and increase of the threshold of the arousal reaction in the neocortex. In rats, a decrease occurs in the amplitude and frequency of electrical cortical activity (Fodor et al, 1964).

The electroencephalogram of patients who become unconscious as a result of carbon monoxide poisoning shows the following changes: appearance of slow waves (Faure et al, 1965; Revol et al, 1966; Sasaki et al, 1966); low voltage fast waves (Tatetsu et al, 1968); delta waves (Mann, 1965; Tatetsu et al, 1967a); reduction in fast waves accompanied by rapid eye movement (Karacan et al, 1971); and a decrease in cortical evoked potential to photic, somatosensory and auditory stimulation (Kuroiwa et al, 1968). Most of these abnormalities remain even for two to five years after the acute exposure to carbon monoxide (Orioli and Cattania, 1965; Geier, 1966; Inanaga et al, 1966a and b; Revol et al, 1966; Sasaki et al, 1966; De Valois and Schadé, 1967; Tatetsu et al, 1967b; Harada and Kozuma, 1968; Tomonari, 1968; Grohme et al, 1969; Yasuoka, 1970; Harada et al, 1971; Totsuka et al, 1971).

1005051160

- LAMY M and HANQUET M : Cinquante cas d'intoxication oxycarbonée traités par l'oxygénothérapie hyperbare. (Fifty cases of carbon monoxide poisoning treated by hyperbaric oxygenation). Acta Anaesthesiol Belg 20: 49-64, 1969. A 821
- LARCAN A, ROBERT J, CALAMAI M and FREJAVILLE J P : Le traitement de l'intoxication oxycarbonée par le caisson mobile de l'oxygénation hyperbare. Premiers résultats. (Treatment of carbon monoxide poisoning using a mobile hyperbaric oxygenation unit. 1st results) Presse Med 75: 1325-9, 1967. A 822
- LEDINGHAM I M: New approaches to poisoning. Hyperbaric oxygenation. Proc Roy Soc Med 57: 807-9, 1964. A 823
- LEE K : The use of portable oxygen sets in carbon monoxide poisoning. Trans Soc Occup Med 16: 85-6, 1966. A 824
- MANTZ J M and TEMPE J D : L'ossigenoterapia iperbarica. (Hyperbaric oxygen therapy). Minerva Med 59: 3137-49, 1968. A 825
- MANTZ J M and TEMPE J D : Die Sauerstoff-Überdruckbehandlung. (Treatment with hyperbaric oxygen). Munch Med Wochensh 110: 2186-97, 1968. A 826
- MATTHEW H : Mobile pressure chamber. Brit Med J 600, 1970. A 827  
and BAYLISS G J A
- MILLER J N : Clinical applications of hyperbaric oxygen therapy in Sydney - a review of recent cases. Med J Aust 1: 835-8, 1966. A 828
- MOLFINO F and ZANNINI D : Sulla cura dell'ossicarbonismo acuto mediante ossigenoterapia in camera pressurizzata. (On the treatment of acute carbon monoxide poisoning by oxygen therapy in a pressure chamber. Rass Med Industr 33: 334-6, 1964. A 829
- NORMAN J N, MacINTYRE J, SHEARER J R and SMITH G : Use of a one-man mobile pressure chamber in the treatment of carbon monoxide poisoning. Brit Med J 2: 333-4, 1970. A 830
- OHRESSER P, CHASSON J, JOUGLARD J, GOUIN F, DUBOULOZ F and TASSY J : Traitement hyperbare des intoxications oxycarbonées. (Hyperbaric treatment of carbon monoxide poisoning). Marseille Med 105: 555-7, 1968. A 831
- PETTY T L : Oxygen therapy. Ann Intern Med 71: 666-7, 1969. A 832
- ROCHE L, BERTOYE A, VINCENT P, MOTIN J, GARIN J P, BOLOT J F and CHADENSON O : Comparaison de deux groupes de vingt intoxications oxycarbonées traitées par oxygène normobare et hyperbare. (Comparison of 2 groups of 20 cases of carbon monoxide poisoning treated with normobaric and hyperbaric oxygen. Lyon Med 220: 1483-99, 1968. A 833
- SAVATEEV N V, TONKOPIJ V D and FROLOV S F : (Hyperbaric oxygenation in certain types of acute poisoning. Review of the literature). Voen. Med Zh 2: 23-8, 1970. A 834
- SCHULTE J H : The use of hyperbaric oxygen in clinical medicine. J Occup Med 11: 462-5, 1965. A 835
- SIMPSON B R and RITCHIE H D : Expérience personnelle du traitement à l'oxygène à haute pression. (Personal experience with high pressure oxygen therapy). Lyon Chir 64: 5-12, 1968. A 836
- SLUJTER M E : The treatment of carbon monoxide poisoning by administration of oxygen at high atmospheric pressure. Progr Brain Res 24: 123-82, 1967. A 837
- SMITH G : Carbon monoxide poisoning. Ann NY Acad Sci 117: 684-7, 1965. A 838
- SUGIMOTO T and YASUMITSU T : (Hyperbaric oxygen therapy of carbon monoxide poisoning). Jap J Clin Med 27: 2157-64, 1969. A 839

1005051295

Bibliography VI F

- SASAKI K, HIRANO H, NAGAHARA T, USUI H/ and TSJKADA J: (Electroencephalograms in acute carbon monoxide poisonings.) Naika 17: 969-72, 1966. 544
- TATETSU S, Kiyota K, TOYA G, TERAOKA H, FUJITA E, INOUE T, MIMURA K, HARADA M, TAKAGIM, YAMAGATA M, KOZUMA Z, KOZUMA Y, MIYAGAWA T, TOMONARI H, TERAOKA A, MURAYAMA E, YASUOKA F, MIYOSHI K and KASAGI Y: (EEG of carbon monoxide poisoning in mass accident due to an explosion of coal dust.) Psychiatr Neurol Jap 69: 71-97, 1967a. 545
- TATETSU S, TOYA G, FUJITA E, INOUE T, HARADA M, TOMONARI H, MURAYAMA E, YASUOKA H, MIYOSHI K and KASAGI Y: (The EEG and prognosis in carbon monoxide poisoning - serial observation for two years.) Brain Nerve(Tokyo) 19: 210-7, 1967b. 546
- TATETSU S, HARADA S, NAKAMURA K, KASAGI Y, ISIKAWA T and KAMANO Y: (An electroencephalographical study of neuropsychiatric disturbances due to intoxication.) J Kumamoto Med Soc 42: 371-8, 1968. 547
- TOMONARI H: (A clinical study on the sequelae of acute carbon monoxide poisoning.) Psychiat Neurol Jap 70: 1029-48, 1968. 548
- TOTSUKA S, MORO K, HORIE T and YAZAKI M: (Two autopsied cases of gas poisoning (propane and carbon monoxide) - with special reference to the non-interval and most-prolonged cases.) Adv Neurol Sci(Tokyo) 15: 592-605, 1971. 549
- YASUOKA F: (Clinical studies on the sequelae of acute carbon monoxide poisoning.) Psychiat Neurol Jap 72: 938-65, 1970. 550

1005051162

## VII. OTHER ORGAN SYSTEMS

The investigations of the effects of carbon monoxide on organs other than the respiratory, circulatory and nervous systems are reviewed in this section. Two topics are remotely related to cigarette smoking and the coverage of each is confined to a bibliographic list: the effects of carbon monoxide on hepatic cells (see additional bibliographic list no. 12) and on unicellular organisms and lower vertebrates (see additional bibliographic list no. 13).

### VII A. Renal System

Some patients with severe carbon monoxide poisoning develop acute renal failure (Burck and Portwich, 1964; Nicolas and Nicolas, 1964; Kobulniczky and Koncz, 1966; Wijdeveld, 1968; Leonowicz, 1967; Linton et al., 1968; Quintana et al., 1969; Grosse and Neuhaus, 1970; Wojczuk and Chylak, 1971). The mechanism for the development of acute renal failure is believed to be other than hypoxia alone. Pauli et al. (1968a) noted a difference in the renal function of normal subjects at a high altitude (3,454 meters above sea level) for 10 days and that of 6 subjects exposed to carbon monoxide for 10 days with a carboxyhemoglobin concentration of 15%. In the presence of carboxyhemoglobin there is an increase of glomerular filtration rate with renal plasma flow remaining unchanged. This combination was interpreted to mean constriction of the efferent vessels with dilatation of the afferent arterioles. On the other hand, hypoxia caused an increase in renal plasma flow with an accompanying change in glomerular filtration rate in the same direction. In the same group of subjects, renal tubular electrolyte handling was consistent with the primary

1005051163

elevation of glomerular filtration rate during exposure to carboxyhemoglobin (Pauli et al., 1968b). There was no increase in urinary elimination of protein (Steiner et al., 1971).

1005051164

## VII. OTHER ORGAN SYSTEMS

## A. Renal System

- BURCK H C and PORTWICH F: Akute Niereninsuffizienz nach schwerer Kohlenmonoxyd-Intoxikation. (Acute renal failure after severe carbon monoxide poisoning.) Frankfurt Z Path 73: 520-34, 1964. 551
- GROSSE G and NEUHAUS G A: Amino-peptidase-Aktivitat im Urin bei Patienten mit Hypnotica-Intoxikationen und unter den Bedingungen der Vita reducta. (Amino-peptidase activity in urine in patients with sedative poisoning and under conditions of vita reducta.) KlinWochenschr 48: 978-85, 1970. 552
- and KONGZ E
- KOBULNICZKY E: Szénmonoxid-mérgezéshez társult myoglobiniuriás vesételelésig. (Carbon monoxide poisoning associated with myoglobinuria and kidney failure.) Orv Hetil 107: 2377-9, 1966. 553
- LEONOWICZ K: (Acute renal failure due to carbon monoxide poisoning.) Wiad Lek 20: 1805-7, 1967. 554
- LINTON A L, ADAMS J H and LAWSON D H: Muscle necrosis and acute renal failure in carbon monoxide poisoning. Postgrad Med J 44: 338-41, 1968. 555
- NICOLAS F and NICOLAS G: Insuffisance rénale aiguë complication d'une intoxication grave par l'oxyde de carbone. (Acute renal failure, a complication of severe carbon monoxide poisoning.) Anesth Analg(Paris) 21: 659-76, 1964. 556
- PAULI H G, TRUNIGER B, LARSEN J K and MULHAUSEN R O: Renal function during prolonged exposure to hypoxia and carbon monoxide. I. Glomerular filtration and plasma flow. Scand J Clin Lab Invest 22 Suppl 103: 55-60, 1968A. 557
- PAULI H G, TRUNIGER B, LARSEN J K and MULHAUSEN R O: Renal function during prolonged exposure to hypoxia and carbon monoxide. II. Electrolyte handling. Scand J Clin Lab Invest 22 Suppl 103: 61-7, 1968B. 558
- QUINTANA F V, MIRETE J O and GARCIA J B: Intoxicacion por monoxido de carbono con fracaso renal agudo. Rev Clin Esp 137-42, 1969. 559
- STEINER S, LARSEN J K, DONATH A and PAULI H G: Renal function and protein elimination of human subjects during carbonmonoxide exposure. Helv Med Acta 36: 39-42, 1971. 560
- WIJDEVELD P G A B: Acute tubulaire nierinsufficiëntie bij vergiftiging door koolmonoxyde. (Acute tubular kidney insufficiency caused by carbon monoxide poisoning.) Nederl T Geneesk 112: 497-503, 1968. 561
- WOJCZUK J and CHYLAK I: Przypadek ostrej niewydolności nerek po zatruciu tlenkiem węgla. (Case of acute renal failure following carbon monoxide poisoning.) Wiad Lek 24: 65-7 1971. 562

1005051165

### VII B. Pregnancy

Accidental carbon monoxide poisoning in a pregnant mother has been known to cause fetal death (Goldstein, 1965; Kells, 1969; Stokowski and Kęsiak, 1969). There is a report of a case without fetal abnormality (Larcan et al., 1970). Most other reports describe brain lesions in babies born of mothers who have suffered from carbon monoxide poisoning (Bankl and Jellinger, 1967; Kamraj-Mazurkiewicz, 1967; Szilágyi, 1967; Beaudoin et al., 1969; Matsuyama, 1969).

The diffusion of carbon monoxide across the placenta has been investigated in the sheep (Metcalf et al., 1965; Longo et al., 1967, 1969; Longo, 1970; and Piédelièvre et al., 1969) and in humans (Gemzell et al., 1958; Friberg et al., 1959). Small quantities of carbon monoxide have been administered to pregnant women for measurement of the diffusing capacity of the placenta (Delivoria-Papadopoulos and Coburn, 1972). The radioactive form of carbon monoxide administered by inhalation has been used to localize the placenta in a mother close to parturition (Glass et al., 1968; Hakim, 1970).

Carboxyhemoglobin levels in maternal and fetal blood have been measured by several investigators (Table 7). The levels in mothers who smoked were higher than in those who were nonsmokers.

**1005051166**

Respiration in infants of mothers who smoked took longer to establish (Tanaka, 1967; Scoppetta, 1968). There was a higher incidence of prematurity among mothers who smoked (Girond, 1967) and a higher neonatal mortality rate (Comstock et al., 1971). The period of gestation is 29 to 34 hours

shorter for smokers than for nonsmokers (Buncher, 1969). The birth weight of babies born of mothers who smoked was lower (Mantell, 1964; Baribaud et al., 1970; Murphy and Mulcahy, 1971; Astrup et al., 1972).

There is a rise in carboxyhemoglobin level in the blood of the newborn. Behrman et al. (1971) have noted a mean rise of 6.98% in 16 newborn infants, which related to the level of pollution in the nursery. When the concentration of carbon monoxide exceeded 20 ppm, there was a decrease of 11.4% in the oxygen-carrying capacity during the first 24 hours and a decrease of 13.9% during the 25 to 85 hours after birth. These changes do not correlate with the history of maternal smoking and are caused by the concentration of carbon monoxide in the atmosphere.

The effect of carbon monoxide alone on the birth weight of rabbits has been investigated by Astrup et al. (1972). An exposure to 90 ppm carbon monoxide in the blood and in rats by Younoszai et al. (1969). resulting in 9 to 10% carboxyhemoglobin/caused a reduction in birth weight and an increase in neonatal mortality. The influence of a level of 5% carboxyhemoglobin was not investigated.

(Table 7 appears on the following page.)

1005051167



Table 7. Carboxyhemoglobin levels of fetal and maternal blood.

Reference (year)	Nonsmoking mothers			Smoking mothers		
	No of mothers	Maternal Mean $\pm$ SD (Range)	Fetal Mean $\pm$ SD (Range)	No of mothers	Maternal Mean $\pm$ SD (Range)	Fetal Mean $\pm$ SD (Range)
Haddon, Nesbitt and Garcia (1961)	7	(0.1-3.1)	(0.2-2.8)	12	(0.1-8.4)	(0.1-9.8)
Heron (1962)	27	2.6 (0.4-4.4)	2.5 (0.2-3.6)	21	6.7 (1.6-14)	5.0 (1.1-9.2)
Young and Pugh (1963)	9	1.6 $\pm$ 0.42	1.61 $\pm$ 0.37	6	2.0 $\pm$ 0.77	2.4 $\pm$ 0.74
Bjure and Fallström (1963)	8	1.0 (0.7-1.43)	1.15 (0.77-1.64)			
Younaszai, Kacic and Haworth (1968)	11	1.2 (0-3)	0.7 (0-1.5)	10	8.3 (2-12)	7.3 (2-10)
Cole, Hawkins and Roberts (1972)	129	1.2 (0-2.4)	2.2	93	4.1 (0.5-14)	7.5
Astrup, Trolle, Olsen and Kieldsen (1972)	884	0.87		824	1.92	

1005051168

## VII. OTHER ORGAN SYSTEMS

## B. Pregnancy

- |   | Reprint |
|---|---------|
| ASTRUP P, TROLLE D, OLSEN H M and KJELDSEN K: Effect of moderate carbon monoxide exposure on fetal development. <u>Lancet</u> 2: 1220-22, 1972.   | 563     |
| BANKL H and JELLINGER K: Zentralnervöse Schaden nach fetaler Kohlenoxydvergiftung. (Central nervous system injuries following fetal carbon monoxide poisoning.) <u>Beitr Path Anat</u> 135: 350-76, 1967.           | 564     |
| BARIBAUD L, YACOB M, FAURE J, MALINAS Y and CAU G: L'oxycarbonémie de l'enfant né de mère fumeuse. (Blood carbon monoxide in children born of smoking mothers.) <u>Med Leg Donn Corp Paris</u> 3: 272-4, 1970.      | 565     |
| BEAUDOING A, GACHON J, BUTIN L P and BOST M: Les conséquences fœtales de l'intoxication oxycarbonée de la mère. (Fetal consequences of carbon monoxide poisoning of the mother.) <u>Pediatric</u> 24: 539-53, 1969. | 566     |
| BEHRMAN R E, FISHER D E and PATON J: Air pollution in nurseries: Correlation with a decrease in oxygen-carrying capacity of hemoglobin. <u>J Pediatr</u> 78: 1050-4, 1971.  | 567     |
| BJURE J and FALLSTRÖM S P: Endogenous formation of carbon monoxide in newborn infants. I. Non-icteric and icteric infants without blood group incompatibility. <u>Acta Pediatr</u> 52: 361-6, 1963.                 | 568     |
| BUNCHER C R: Cigarette smoking and duration of pregnancy. <u>Am J Obstet Gyn</u> 103/7: 942-6, 1969.  | 569     |
| COLE P V, HAWKINS L H and ROBERTS D: Smoking during pregnancy and its effects on the fetus. <u>J Obstet Gyn</u> 79: 782-7, 1972.  | 570     |
| COMSTOCK G W, SHAH F K, MEYER M B and ABBEY H: Low birth weight and neonatal mortality rate related to maternal smoking and socioeconomic status. <u>Am J Obstet Gyn</u> 111: 53-9, 1971.                           | 571     |
| DELIVORIA-PAPADOPOULOS M and COBURN R F: Placental carbon monoxide diffusing capacity ( $D_{CO}$ ) in pregnant women at term. <u>Fed Proc</u> 31(2): 237, 1972.   | 572     |
| FRIBERG L, NYSTROM A and SWANBERG H: Transplacental diffusion of carbon monoxide in human subjects. <u>Acta Physiol Scand</u> 45: 363-68, 1959.   | 573     |
| GEMZELL C A, ROBBE H and STROM G: On the equilibration of carbon monoxide between human maternal and fetal circulation in vivo. <u>Scand J Clin Lab Invest</u> 10: 372-8, 1958.                                     | 574     |
| GIROND J: Le délit d'omission. <u>Sem Med</u> 43: 69-70, 1967.  | 575     |
| GLASS H I, JACOBY J, WESTERMAN B, CLARK J C, ARNOT R N and DIXON H G: Placental localization by inhalation of radioactive carbon monoxide. <u>J Nucl Med</u> 9: 468-70, 1968.                                       | 576     |
| GOLDSTEIN D P: Carbon monoxide poisoning in pregnancy. <u>Am J Obstet Gyn</u> 92: 526-8, 1965.  | 577     |
| HADDON W Jr, NESBITT R E L and GARCIA R: Smoking and pregnancy: carbon monoxide in blood during gestation and at term. <u>Am J Obstet Gyn</u> 18: 262-7, 1961.  | 578     |

1005051169

- HAKIM C A : Placental localization. A comparison of the doppler ultrasound method with that of radioactive carbon monoxide and the gamma camera. J Obstet Gynecol Commonw 77: 625-6, 1970. 580
- HERON H J : The effects of smoking during pregnancy: a review with a preview. New Zealand Med J 61: 545-8, 1962. 581
- KAMRAJ-MAZURKIEWICZ K : Wpływ zatrucia tlenkiem węgla w okresie ciąży na ośrodkowy układ nerwowy płodu na podstawie obserwowanego przypadku. (Effect of carbon monoxide poisoning in the course of pregnancy on the fetal central nervous system according to our case.) Ginek Pol 38: 291-4, 1967. 581
- KELLS D U : Carbon monoxide poisoning during pregnancy. Medicoleg Bull 177: 1-11, 1968. 582
- LARCANA A, LANDES P and VERT P : Intoxication oxycarbonée au 2<sup>e</sup> mois de grossesse sans anomalie néonatale. (Carbon monoxide poisoning during the 2nd month of pregnancy without neonatal abnormality.) Bull Fed Soc Gyn Obstet Lang Fr 22: 338-9, 1970. 583
- LONGO L D : Carbon monoxide in the pregnant mother and fetus and its exchange across the placenta. Ann NY Acad Sci 174: 312-41, 1970. 584
- LONGO L D, POWER G G and FORSTER R E II : Respiratory function of the placenta as determined with carbon monoxide in sheep and dogs. J Clin Invest 46: 812-28, 1967. 585
- LONGO L D, POWER G G and FORSTER R E II : Placental diffusing capacity for carbon monoxide at varying partial pressures of oxygen. J Appl Physiol 26: 360-70, 1969. 586
- MANTELL, C D : Smoking in pregnancy: The role played by carbonic anhydrase. New Zealand Med J 63: 601-3, 1964. 587
- MATSUYAMA H : (Effects of maternal fuel-gas poisoning on / fetal brain.) Adv Neurol Sci Tokyo 34-8, 1969. 588
- METCALFE J, MALL W, BARTELS H, HILPERT P and PARER J T : Transfer of carbon monoxide and nitrous oxide in the artificially perfused sheep placenta. Circ Res 16: 95-101 1965. 589
- MURPHY J F and MULCAHY R : The effect of age, parity, and cigarette smoking on baby weight. Am J Obstet Gyn 111: 22-5, 1971. 590
- PIÉDELÈVRE C, BRETON R and DÉROBERT L : Intoxication par l'oxyde de carbone et passage du toxique au travers du filtre placentaire. (Carbon monoxide poisoning and passage of the poison across the placental filter.) Med Leg Domm Corpor (Paris) 2: 389-92, 1969. 591
- SCOPPETTA V : Sul contenuto di ossido di carbonio nel sangue circolante di gestanti fumatrici. (On the carbon monoxide content of the circulating blood of pregnant smokers. Arch Ostet Ginec 73: 369-75, 1968. 592
- STOKOWSKI C and KESIAK J : Zatrucie tlenkiem węgla u ciężarnej. (Carbon monoxide poisoning in pregnancy.) Ginek Pol 40: 801-4, 1969. 593
- SZILÁGYI L : Magzati CO mérgezés idegrendszeri maradandó ártalma. (Chronic neurologic damage from fetal CO poisoning.) Orv Hetil 108: 1559-61, 1967. 594
- TANAKA M : Studies on the mechanism of retardation of foetal growth by smoking during pregnancy. J Jap Obstet Gyn Soc 14: 45, 1967. 595
- YOUNOSZAI M K, KACIC A and HAWORTH J C : Cigarette smoking during pregnancy: The effect upon the hematocrit and acid-base balance of the newborn infant. Canad Med Ass J 99: 197-200, 1968. 596

1005051170

Bibliography VII B

YOUNOSZAI M K, PELOSO J and HAWORTH J C : Fetal growth retardation in rats exposed to cigarette smoke during pregnancy. Ain J Obstet Gyn 104: 1207-13, 1969.

597

YOUNG I M and PUGH L G C E : The carbon monoxide content of foetal and maternal blood. J Obs Gyn Brit Commonw 70:681-3, 1963.

598

1005051171

### VII C. Endocrine System

The sex organs in animals influence the survival rate in carbon monoxide exposure. Stupfel et al. (1971) noted a higher mortality in male than in female mice and that castration increased the mortality in the female. The mechanism by which estrogens and progestogens increase the tolerance to carbon monoxide poisoning is not known. On the other hand, carbon monoxide in vitro depresses some of the enzymes that are involved in the biosynthesis of estrogen (Meigs and Ryan, 1971). The structure of the ovary was unaffected and all the stages in the development of the ova were noted in rats exposed to carbon monoxide in vivo (Marnatsashvili, 1970). The effects of carbon monoxide on male fertility are uncertain (Wittgens, 1966).

Adrenalectomy increased the mortality of rats and mice in response to carbon monoxide (Pukhov, 1964). In animals with adrenals intact, adrenal cortical activity is stimulated by exposure to carbon monoxide (Katsuki et al., 1966; Hanke and Kiereś, 1967). Stimulation of the adrenal medulla also occurred, which contributed to the hyperglycemia (Bour et al., 1968; Hirano et al., 1968). There are substances in the rat adrenal gland which combine with carbon monoxide (Harding et al., 1964; Wilson and Harding, 1970). The role of these substances in mediating the stimulation of the adrenal gland and in reducing the lethality of carbon monoxide has not been considered. The reduction in rate of growth of rats exposed to carbon monoxide is associated with a decrease in activity of the thyroid gland (Truhaut et al., 1965; Pogrand, 1969).

The inhalation of carbon monoxide in human subjects causes an

1005051172

elevation of blood sugar (Hirano et al, 1967; Nakao, 1969). The hyperglycemia induced by cigarette smoking is believed to be due to nicotine stimulation of the adrenal medulla (Haggard and Greenberg, 1934). The contribution of carboxyhemoglobin derived from cigarette smoking has not been estimated.

1005051173

## C. Endocrine System

- BOUR H, GUY-GRAND B, ROGER M, TUTIN M and DORF G: Étude de 41 cas d'intoxication oxycarbonée aigue par les épreuves dynamiques de la glycorégulation. (Study of 41 cases of acute carbon monoxide poisoning with dynamic tests of glycoregulation.) Presse Med 76: 1051-4, 1968. Reprint 599
- HAGGARD H W and GREENBERG L A: The effects of cigarette smoking upon the blood sugar. Science 79: 165-6, 1934. 601
- HANKE J and KIERES H: Wydalanie 17-keto- i 17 ketogennych sterydów z moczem w ostrych zatruciach niektórymi substancjami chemicznymi. (Excretion of 17-keto and 17-ketogenic steroids in the urine in acute poisoning with some chemical substances.) Med Pracy 12: 109-16, 1967. 602
- HARDING B W, WONG S H and NELSON D H: Carbon monoxide-combining substances in rat adrenal. Biochim Biophys Acta 92: 415-7, 1964. 603
- HIRANO H, INOUE Y and TANAMI J: (Studies on the mode of action of carbon monoxide. 1st Report: Changes of percentage saturation of carboxyhemoglobin and of blood sugar levels during inhalation of carbon monoxide.) Jap J Hyg 22: 559-62, 1967. 604
- HIRANO H, INOUE Y and TANAMI J: (Studies on the mode of action of carbon monoxide. 2nd Report: Pattern of action of carbon monoxide. Jap J Hyg 23: 286-92, 1968. 605
- KATSUKI S, ANDO and NISHIDA Y: (Diurnal variations in blood 11-OHCS level in acute carbon monoxide poisoning.) Clin Endocr Tokyo 14: 475-8, 1966. 606
- MAMATSASHVILI M J: On the detrimental effect of carbon monoxide and sulfur dioxide on fertility of female rats. Hyg Sanit 76: 277-9, 1970. 607
- MEIGS R A and RYAN K J: Enzymatic aromatization of steroids. I. Effects of oxygen and carbon monoxide on the intermediate steps of estrogen biosynthesis. J Biol Chem 246: 83-7, 1971. 607
- NAKAO K: Carbon monoxide poisoning, especially from view point of Hematology. Adv Neurol Sci(Tokyo) 13: 21-4, 1969. 608
- POGRUND R S: Biologic synergisms in rats produced by carbon monoxide and positive ions. Int J Biometeor 13: 123-34, 1969. 609
- PUKHOV V A: The influence of adrenal insufficiency and hyperfunction upon the sensitivity of albino rats and mice to carbonic oxide poisoning.) Framakol Toksik 27: 343-5, 1964. 610
- STUPFEL M, BOULEY G and POLIANSKI J: Castration et mortalité de la Souris par oxyde de carbone. (Castration and mortality of mice due to carbon monoxide.) J Physiol (Paris) 63: 99A-100A, 1971. 611
- TRUHAUT R, BOUDENE C and CLAUDE J R: Fixation thyroïdienne de l'iode chez le rat intoxiqué chroniquement par l'oxyde de carbone. (Thyroid fixation of iodine-131 in the rat chronically poisoned with carbon monoxide.) Ann Biol Clin(Paris) 23: 73-82, 1965. 612
- WILSON L D and HARDING B W: Studies on adrenal cortical cytochrome P-450. IV. Effects of carbon monoxide and light on cholesterol side chain cleavage. Biochemistry 9: 1621-5, 1970. 613
- WITTGENS H: Fertilitätsstörungen des Mannes und Beruf. (Fertility disorders in men and occupation. Berufsdermatosen 14: 105-8, 1966. 614

1005051174

#### VII D. Bone, Muscle and Skin

There is a deposition of bone in soft tissues in patients who suffer from carbon monoxide poisoning (Ri, 1966; Mouren et al., 1972). The mechanism for this deposition is not known. In the laboratory, carbon monoxide has been used as a tool to test the polarizing power of bone mineral (Nash and Beebe, 1969).

Some patients suffering from acute carbon monoxide poisoning develop contractures of the skeletal muscle (Howse and Seddon, 1966; Kolb, 1968). Three factors contribute to the development of contracture:

(1) ischemia due to reduction in blood oxygen content; (2) hyperexcitability of neuromuscular junction (Paris, 1964); and (3) binding of myoglobin with carbon monoxide (Rossi-Fanelli and Antonini, 1958; Wittenberg et al., 1965; Gladyshevskaja et al., 1966; Coburn and Mayers, 1971; Rudolph et al., 1972).

The cutaneous lesions in a patient with carbon monoxide poisoning are varied. They include bullae, subepidermal vesicles, intracellular edema and occlusion of the epidermal portion of sweat ducts (Jopkiewicz et al., 1965; Long, 1969; Leavell et al., 1969; Ippen and Goerz, 1969; Baden, 1970; Achten et al., 1971). Erythema of the face associated with a cherry-red color of the blood is a classical diagnostic sign of poisoning with carbon monoxide (Danto, 1964).

1005051175



## VII. OTHER ORGAN SYSTEMS

## D. Bone, Muscle and Skin

- |   | Reprint |
|---|---------|
| ACHTEN G, LEDOUX-CORREUSIER M and THYS J P.: Intoxication à l'oxyde de carbone et lésions cutanées. (Carbon monoxide poisoning and cutaneous lesions.) <u>Ann Derm Syphil Paris</u> 93: 421-8, 1971.  | 615     |
| BADEN M M: Bullous skin lesions in barbiturate overdosage and carbon monoxide poisoning. <u>JAMA</u> 213, 2271, 1970.   | 616     |
| COBURN R F and MAYERS L B: Myoglobin O <sub>2</sub> tension determined from measurements of carboxymyoglobin in skeletal muscle. <u>Am J Physiol</u> 220: 66-74, 1971.  | 617     |
| DANTO B L: The man with a red face. <u>Am J Psychiat</u> 121: 275-6, 1964.  | 618     |
| GLADYSHEVSKAI A T N, DOLOSHITSKY L M and SOBCHUK B A: (Myoglobin in intoxication of the organism with carbon monoxide.) <u>Ukr Biokhim Zh</u> 38: 9-13, 1966.   | 619     |
| HOWSE A J G and SEDDON H: Ischaemic contracture of muscle associated with carbon monoxide and barbiturate poisoning. <u>Brit Med J</u> 5483: 192-5, 1966.   | 620     |
| IPPEN H, and GOERZ G: Carbon monoxide and dermal changes. <u>JAMA</u> 207: 1718, 1969.  | 621     |
| JOPKIEWICZ R, KONECKI J and WENTKOWSKI A: Trwale zmiany rumieniowe skóry w następstwie ostrego zatrucia tlenkiem węgla. (Durable erythema of the skin due to acute carbon monoxide poisoning.) <u>Med Pracy</u> 16: 517-24, 1965.   | 622     |
| KOLB K: Lokale ischämische Kontraktur der Hand nach Suicidversuchen. (Local ischemic contracture of the hand following suicide attempts.) <u>Hefte Unfallheilk</u> 94: 208-9, 1968.   | 623     |
| KOLB K P: Lokale ischämische Kontrakturen der Hand nach Suizidversuchen. (Local ischemic contractures of the hand following suicide attempts.) <u>Münch Med Wochenschr</u> 110: 1873-4, 1968.   | 624     |
| FARLEY C H  |         |
| LEAVELL U W and MCINTYRE J S: Cutaneous changes in a patient with carbon monoxide poisoning. <u>Arch Derm</u> 99: 429-33, 1969.   | 625     |
| LONG P I: Carbon monoxide poisoning. <u>Arch Derm</u> 100: 385, 1969.   | 626     |
| MOUREN P, POINSON Y, JOUGLARD M, GIUDICELLI S, FRESCO R and D'OMEZON Y: Les para-ostéo-arthropathies neurogènes (a propos de deux observations au cours d'intoxication oxycarbonée grave). (Neurogenic paraosteoarthropathies (a propos of 2 cases during severe carbon monoxide intoxication.) <u>Mars Med</u> 109: 17-26, 1972. | 627     |
| NASH R J and BEEBE R A: Heats of adsorption of carbon monoxide on bone mineral and on thorium oxide by gas-solid chromatography. <u>J Colloid Interface Sci</u> 31: 343-52, 1969.   | 628     |
| PARIS J: Etude des courbes 'excitation-durée' apres intoxication aigue professionnelle par l'oxyde de carbone. (Study of the 'excitation-duration' curves after occupational acute carbon monoxide poisoning.) <u>Rass Med Industr</u> 33: 275-91, 1964.  | 629     |
| RI K: (Case of metaplastic ossification in carbon monoxide poisoning.) <u>Orthop Surg Tokyo</u> 17: 397-403, 1966.  | 630     |
| ROSSI-FANELLI A and ANTONINI E: Studies on the oxygen and carbon monoxide equilibria of human myoglobin. <u>Arch Biochem Biophys</u> 77: 478-92, 1958.  | 631     |
| RUDOLPH S A, BOYLE S O, DRESDEN C F and GILL S J: A calorimetric study of the binding of carbon monoxide to myoglobin. <u>Biochemistry</u> 11: 1098-101, 1972.  | 632     |
| WITTENBERG B A, BRUNORI M, ANTONINI E, WITTENBERG J and WYMAN J: Kinetics of the reactions of alysia myoglobin with oxygen and carbon monoxide. <u>Arch Biochem</u>   |         |

1005051176

## VIII. CARBON MONOXIDE POISONING

It is rather a remote possibility that cigarette smoking may cause death by carbon monoxide poisoning. There are publications on the diagnosis and treatment of carbon monoxide poisoning, both accidental and industrial. The additional bibliography lists relate to the following topics:

Accidental poisoning involving carbon monoxide and other causative factors (no. 14).

Accidental poisoning involving carbon monoxide (no. 15).

Industrial poisoning involving carbon monoxide (no. 16).

Therapy of carbon monoxide poisoning (no. 17).

Therapy of poisoning by use of normobaric and hyperbaric oxygen (no. 18).

1005051177

## IX. COMMENTARY OF SELECTED ARTICLES

Page 114

Page		Reprint
116	(1) <u>Smoking and Health: Report of the Advisory Committee to the Surgeon General of the Public Health Service, 344-345, 1964.</u>	634
117	(2) NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH: <u>The health consequences of smoking, 62-64, 1967.</u>	635
119	(3) NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH: <u>The health consequences of smoking, 1968 Supplement to the 1967 Public Health Service Review, 38-40, 1968.</u>	636
121	(4) NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH: <u>The health consequences of smoking, 1969 Supplement to the 1967 Public Health Service Review, 28-29, 1969.</u>	637
123	(5) NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH: <u>The health consequences of smoking. A report to the Surgeon General: 1971. 59-62, 1971.</u>	638
128	(6) NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH: <u>The health consequences of smoking. A report to the Surgeon General: 1972. 21-23, 121-135, 1972.</u>	639
141	(7) ABELSON P H: A damaging source of air pollution. <u>Science</u> 158: 1527, 1967.	640
142	(8) ANON: Carbon monoxide. <u>Am Industr Hyg Ass J</u> 26: 431-4, 1965.	641
143	(9) ANON: Carbon monoxide poisoning - a timely warning. <u>New England J Med</u> 276: 849-50, 1968.	642
144	(10) ANON: Warning: Cigarettes are dangerous to your health. American Cancer Society. <u>Med Bull Montgomery County Med Soc</u> 24: 45-7, 1968.	643
145	(11) ANON: World action on smoking. <u>Brit Med J</u> 4: 65, 1971.	644
146	(12) ANON: Cigarette smoking and carbon monoxide. <u>Med Letter Drug Ther</u> 13: 91-2, 1971.	645
147	(13) ARONOW W S, KAPLAN M A and JACOB D: Tobacco: A precipitating factor in angina pectoris. <u>Ann Int Med</u> 69: 529-36, 1968.	646
147	(14) ARONOW W S, DENDINGER J and ROKAW S N: Heart rate and carbon monoxide level after smoking high-, low-, and non-nicotine cigarettes. A study in male patients with angina pectoris. <u>Ann Int Med</u> 74: 697-702, 1971.	647
147	(15) ARONOW W S and ROKAW S N: Nonnicotine cigarettes.: Effects in angina pectoris. <u>Circulation</u> 44: 782-8, 1971.	648
147	(16) ARONOW W S, HARRIS C N, ISBELL M W, ROKAW S N and IMPARATO B: Effect of freeway travel on angina pectoris. <u>Ann Int Med</u> 77: 669-76, 1972.	649
148	(17) AYRES S M, GIANNELLI S Jr and MUELLER H: Myocardial and systemic responses to carboxyhemoglobin. <u>Ann NY Acad Sci</u> 174: 268-93, 1970.	650
148	(18) AYRES S M, MUELLER H S, GREGORY J J, GIANNELLI S Jr and PENNY J L: Systemic and myocardial hemodynamic responses to relatively small concentrations of carboxyhemoglobin (COHB). <u>Arch Environ Health</u> 18: 699-709, 1969.	651
149	(19) AYRES S M: Roles of carbon monoxide and nicotine in circulatory effects of cigarette smoke. <u>JAMA</u> 219: 520, 1972.	652

1005051178

- 151 (20) BANYAI A L : Ominous misalliance: Inhalation of carbon monoxide from motor vehicles and cigarettes. Chest 58: 532, 1970. 653
- 153 (21) BARTLETT D Jr : Pathophysiology of exposure to low concentrations of carbon monoxide. Arch Environ Health 16: 719-27, 1968. 654
- 154 (22) CAMM A J : The effects of smoking. Guy Hosp Gaz 81: 185-203, 1967. 655
- 155 (23) CONROY J P : Smoking and the anesthetic risk. Anest Anal 48: 388-400, 1969. 656
- 156 (24) CURPHEY T J : Carboxyhemoglobin in relation to smoking. Nat Cancer Inst Monogr 28: 231-5, 1968. 657
- 160 (25) DINMAN B D : Carbon monoxide and cigarette smoking. JAMA 212: 1785, 1970. 653
- 162 (26) DOYLE J T : Smoking and myocardial infarction. Circulation 39 and 40: Suppl 4: 136-43, 1969. 659
- 163 (27) GIEL B G : Air pollution and your lungs. Pub Health News 46: 246-53, 1965. 660
- 164 (28) GOLDSMITH J R : Carbon monoxide and coronary heart disease. Ann Int Med 71: 199-201, 1969. 661
- 165 (29) GOLDSMITH J R : Carbon monoxide and coronary heart disease: compelling evidence in angina pectoris. Ann Int Med 77: 808-10, 1972. 662
- 166 (30) GOLDSTEIN R E and EPSTEIN S E : Medical management of patients with angina pectoris. Prog Cardiovas Dis 14: 360-98, 1972. 663
- 167 (31) LINDQUIST V A Y : Carbon monoxide: Its relationship to air pollution and cigarette smoking. Publ Health London 86: 20-6, 1970. 664
- 168 (32) NAHUM L H : Smoking and thrombosis. Conn Med 29: 853-4, 1965. 665
- 169 (33) NAHUM L H : Toxic products in cigarette smoke: pleasure or poison. Conn Med 32: 154-5, 1968. 666
- 170 (34) NAHUM L H : The effects of carbon monoxide on human health. Conn Med 33: 90-2, 1969. 667
- 172 (35) ROSE E F and ROSE M : Carbon monoxide: A challenge to the physician. Clin Med 78: 12-21, 1971. 668
- 174 (36) SELTZER C C : The effect of cigarette smoking on coronary heart disease. Arch Environ Health 20: 418-23, 1970. 669

1005051179

- (1) SMOKING AND HEALTH: Report of the Advisory Committee to the Surgeon General of the Public Health Service, 344-5, 1964.

## SMOKING AND ACCIDENTS

Smoking has been associated with a variety of accidents. Among these, fires have the most obvious and important consequences.

In a special study of home accident fatalities in 1952 through 1953, the Public Health Service and the National Safety Council reported that 231 (18%) of 1,274 deaths from fires of known origin were due to cigarettes, cigars or pipes (1).

The Metropolitan Life Insurance Company reported that of 352 deaths in 1956 and 1957 among their policyholders from fires and burns with known causes in and about the home, 57 (16%) were due to smoking (2).

Of physiological responses related to driving, smoking degrades detectably only the differential brightness threshold and this effect increases with amount of smoking (4). The epidemiological data available on the effects of smoking on traffic accidents are inconclusive.

It has been shown that a level of carboxyhemoglobin of 5 percent—a level which is not uncommon among heavy cigarette smokers (3, 6)—depresses visual perception to as great an extent as anoxia at 8,000 to 10,000 feet altitude (4, 5).

1. Home Accident Fatalities: 1952-1953. National Office of Vital Statistics, U.S. Public Health Service, 1956. Mimeographed report. Table 12.
2. How fatal accidents occur in the home. Metrop Life Insur Statist Bull 40: 6-8, November-December, 1959.
3. Larson, P. S., Haeg, H. B., Silvette, H. Carboxy-hemoglobin, p. 107-110. Tobacco: Experimental and Clinical studies. Baltimore, Williams and Wilkins, 1961.
4. McFarland, R. A., Moseley, A. L. Carbon monoxide in trucks and buses and information from other areas of research on carbon monoxide, altitude and cigarette smoking. In: Conference proceedings: Health, medical and drug factors in highway safety. National Academy of Sciences—National Research Council Publication 328, 1954. Sect. 4.17-4.33.
5. McFarland, R. A., Roughton, F. J. W., Halperin, M. H., Niven, J. I. The effects of carbon monoxide and altitude on visual thresholds. J Aviat Med 15: 6, 381-94, 1944.
6. Schrenk, H. H. Results of laboratory tests. Determination of concentration of carbon monoxide in blood. Pub Health Bull 278: 36-49, 1942.

(a) The epidemiological data reported by Boeck (1948) indicate that cigarette smoking is not responsible for the occurrence of vehicular accidents and poor driving (see page 37).

(b) A more recent report by Bender et al (1971) states that exposure to 100 ppm carbon monoxide for 2-1/2 hours, with mean blood levels of 7.2% carboxy-hemoglobin, diminished visual perception. There was no difference between smokers and nonsmokers with regard to their susceptibility (see page 88).

1005051180

(2) NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH: The health consequences of smoking, 62-4, 1967.

*Carbon Monoxide Effect*

The gaseous phase of cigarette smoke contains about 4 percent carbon monoxide. This quantity can increase the levels of carboxyhemoglobin saturation of cigarette smokers from 2 percent to 10 percent (21). The average nonsmoker, depending on environmental exposure, usually has less than 2 percent carboxyhemoglobin saturation (10). Since smokers of one pack or more a day may have chronically elevated carboxyhemoglobin levels of more than 4 percent (9), there may be appreciable differences in the carboxyhemoglobin levels between some heavy cigarette smokers and nonsmokers.

(a) The carboxyhemoglobin blood levels of cigarette smokers expressed as ranging from 2% to 10% is not an accurate statement. A survey of the literature reveals 30 investigations with an overall mean level for 2,054 smokers of 3.76% in the blood taken 4 to 12 hours after smoking. Only 30% of subjects have levels exceeding 5% (see pages 12 to 14). The peak levels after smoking one or more cigarettes had a mean of 5.26% (see page 20). A more accurate statement would be based on integration of levels taken at hourly intervals but such information is not available.

In addition to displacing oxyhemoglobin, carbon monoxide effects a shift in the oxygen-hemoglobin dissociation curve (2, 3, 4, 5, 6). This may result in a decreased release of oxygen at the tissue level. A series of studies (61, 62) has been performed on young adults to analyze the effect of cigarette smoking on carboxyhemoglobin levels, and the consequent effect on some parameters of cardiopulmonary function. An increased post exercise oxygen debt was observed after cigarette smoking as compared to controls. This, in part, may reflect not only ventilatory disturbances but also a decreased supply of oxygen in the blood due to the carbon monoxide effect, resulting in less available oxygen to meet the increased tissue demand. Similar post-exercise oxygen debts have been noted after inhalation of enough carbon monoxide to produce comparable blood levels of carboxyhemoglobin (21).

(b) Inhalation of carbon monoxide in man, with blood levels of 5% to 10% carboxyhemoglobin does not influence ventilation. Ayres et al (1965) used short-term exposure and Astrup et al (1968) exposure for 8 days (see pages 42 and 45).

1005051181

## Commentary (2) National Clearinghouse, Cont.

The consequence of the smoking/carbon monoxide effect appears to be especially important in the myocardium where relatively more oxygen is normally extracted from the coronary circulation as compared to other organ systems. (Coronary venous blood usually has an oxygen saturation of less than 25 percent, whereas blood leaving some other organs is about 75 percent saturated with oxygen (45).)

Dogs were exposed to carbon monoxide to elevate their carboxyhemoglobin saturation levels (9). In response to inhalation of carbon monoxide there was an increase in coronary blood flow but a decrease in coronary arterial-venous oxygen differences. Patients with coronary heart disease were also studied following inhalation of enough carbon monoxide to elevate their carboxyhemoglobin saturation levels to the range of 5 to 12 percent (9). In response to carbon monoxide there was generally an increase in the cardiac output and the coronary blood flow in most of the patients. While the systemic arterial-venous oxygen differences varied, either increasing or decreasing, the coronary arterial-venous oxygen differences decreased, indicating a decreased oxygen extraction by the myocardial tissue despite the myocardium's increased demand for oxygen. These decreases in myocardial oxygen extraction are related to increases in the carboxyhemoglobin saturation levels. It was observed that some patients evidently could compensate by increasing their coronary blood flows adequately to supply the myocardial tissue with sufficient oxygen, as indicated by a rise in myocardial oxygen uptake in these individuals. However, the other patients with coronary heart disease, evidently more severe, could not increase their coronary blood flow rate enough to compensate for the decreased oxygen carried by the blood. This latter group of patients, even though they had increased cardiac output, had less significant increases of coronary blood flow than those noted in the first group of patients. The coronary arterial-venous oxygen differences and the myocardial tissue oxygen uptakes both decreased, indicating that the myocardial tissue oxygen demand was not being met entirely.

(c) The experiment cited in the above paragraph specifies an increase in cardiac output with a carboxyhemoglobin value of 5% to 12%. Experiments by Brody and Coburn (1969, 1970) and Klausen et al (1968) did not show an increase in cardiac output following either acute or chronic exposure of human subjects (see page 56). Experiments on animals show myocardial lesions with exposure exceeding 50% carboxyhemoglobin (see page 57).

The reduction in the amount of oxygen available to the myocardial tissue caused by the absorption of carbon monoxide from tobacco smoke may be especially critical in persons with pre-existing coronary heart disease, especially when they cannot significantly increase coronary blood flow to compensate for increased myocardial tissue oxygen demand. The carbon monoxide effect may, in part, contribute to the increased incidence of myocardial infarctions that occur in cigarette smokers. Additional research is needed.

(d) Carbon monoxide does not contribute to the increased incidence of myocardial infarction that occur in cigarette smokers. Haywood et al (1972) could not find a clear-cut relationship between carbon monoxide levels and incidence of acute infarction. De Bias et al (1972) exposed dogs with experimental myocardial infarction to 100 ppm carbon monoxide (14% carboxyhemoglobin) for 14 weeks and did not observe an increase in severity of myocardial hypoxia (see page 61 and 62).

1005051182

SCHULTE J H : Effects of mild carbon monoxide intoxication. Arch Environ Health 7: 524-30, 1963.

496

STEWART R D, PETERSON J E, BARETTA E D, EACHAND R T, HOSKO M J and HERR-MANN A A : Experimental human exposure to carbon monoxide. Arch Environ Health 21: 154-64, 1970.

497

TROUTON D and EYSENCK H J : The effects of drugs on behavior. Handbook of Abnormal Psychology, H. J. Eysenck, Editor, Basic Books Inc., New York: 634-96, 1961.

1005051156



Commentary (3) National Clearinghouse, Cont.

Anomalous hemoglobin-oxygen dissociation was noted in "heavy" cigarette smokers (more than one pack per day) without known coronary heart disease. In experiments where the amount of cigarette smoking was controlled, there appeared to be a threshold effect: more than 12 cigarettes per day caused this anomalous dissociation to occur (53). Birnstingl (9) reports finding an increased hemoglobin affinity for oxygen in smokers, which does not appear to be explained solely by the increased carboxyhemoglobin levels in smokers.

Research to further study the interrelationships of carbon monoxide to the myoglobin of heart muscle should be performed because it is possible that carbon monoxide may replace oxymyoglobin with carboxymyoglobin and disturb the oxygen-dissociation phenomena of myoglobin (88, 140, 159). The limitations of blood supply and the high energy output of heart muscle as compared to skeletal muscle may make the myoglobin impairments by carbon monoxide of possible etiologic importance in cigarette smoking and heart disease.

Hydrogen cyanide appears to be rapidly converted to thiocyanates by the body, but the absorption by the lung of cyanide from cigarette smoke might possibly result in higher serum cyanide levels in the coronary arteries than in the systemic circulation. As noted in the 1964 Report, the cyanide ion is capable of stopping cellular respiration abruptly through inactivation of cytochrome oxidase. In sublethal exposures, the cyanide ion is gradually released from its combination with the ferric ion of cytochrome oxidase, converted to thiocyanate ion and excreted in the urine. Thiocyanate blood levels in smokers are three times higher than in nonsmokers and relative differences in urinary excretion are even more pronounced. Cytochrome oxidase is very important in cellular respiration of all body cells. In view of the extremely high myocardial cellular needs for aerobic metabolism, it is possible that the cyanide ion inactivation of cytochrome oxidase also can occur in myocardial cells and be of critical importance, especially in light of other risk factors such as impaired coronary blood flow, the carbon monoxide effect, and the known increases in myocardial tissue oxygen demand caused by the smoking/nicotine-induced catecholamine release. Further research is needed to determine whether or not cyanide ions in concentrations equivalent to those found in cigarette smokers, have a harmful effect on the myocardium, in terms of both acute and chronic exposures.

(a) The statement that \_\_\_\_\_ relates directly to cigarette smoking is that of Birnstingl is entirely wrong and has completely reversed the meaning intended by that author. In the original publication of Birnstingl, Cole and Hawkins (1967), the increased oxygen affinity of hemoglobin in the blood of smokers could be "almost entirely due to a raised carbon monoxide hemoglobin level". A copy of a summary of the article which appeared in *Brit J Surg* 54: 615-8; 1967, follows:

Why male subjects over 40 years of age have a higher oxygen affinity which is greater than that of a similar group under this age. This would result in a shift to the left of the oxyhaemoglobin dissociation curve.

Cigarette smokers also have an increased oxygen affinity, but this is almost entirely due to a raised carbon monoxide haemoglobin level.

Patients with Buerger's disease have a normal oxyhaemoglobin dissociation curve, if age and smoking habits are taken into account. Furthermore, their oxyhaemoglobin dissociation does not appear to be more sensitive to the effects of cigarette smoke than normal controls.

1005051184

(27) GIEL B G: Air pollution and your lungs. Public Health News 46: 246-53, 1965.

#### FACTS ABOUT SPECIFIC AIR POLLUTANTS

##### *Carbon Monoxide*

Carbon monoxide is well known to all of us. Yet many smokers are unaware that approximately seven to eight percent of their hemoglobin may be bound as carboxyhemoglobin. If in the mean time, such an individual should develop vascular insufficiency to vital organs, and then be forced to breath ambient air containing 30 ppm of CO for four to six hours or gotten into an atmosphere where he would be exposed to 120 ppm CO for one hour, he would bind an additional five percent of his hemoglobin and could suffer tragic results.

The figure of 7% to 8% carboxyhemoglobin among smokers is an over-estimation. A review of the literature shows an overall mean of 3.76% for 2,054 smokers 4 to 12 hours after smoking, and a peak level of 5.26% after smoking (see pages 12 to 14 and 20 to 21).

1005051227

Commentary (4) National Clearinghouse, Cont.

Most of the references cited are review articles which are commented upon under Bartlett (page 153 ), Goldsmith (page 165 ), and Dinman (page 160 ). The only original work referred to is by Eliot et al, which is a personal communication to the National Clearinghouse.

1005051186

- (5) NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH: The health consequences of smoking. A report to the Surgeon General: 1971. 59-62, 1971.

#### CARDIOVASCULAR EFFECTS OF CARBON MONOXIDE

Carbon monoxide (CO) is a colorless and odorless gas, low levels of which have significant effects on human and animal physiology which are just now beginning to be understood. According to Wynder and Hoffmann (1961), it is present in cigarette smoke in concentrations of approximately 2.9 to 5.1 percent. The concentration of CO in smoke is subject to many factors, among them the type of tobacco and the porosity of cigarette paper. The concentration of CO in smoke has been found to increase significantly toward the last puffs of the cigarette.

(a) This paragraph states the concentration of carbon monoxide in cigarette smoke but omits mentioning the fact that the smoke is diluted with atmospheric air, dead space air and alveolar air.

According to Chevalier, et al. (1966), a concentration of approximately 4 percent CO in cigarette smoke will produce alveolar levels of around 0.04 percent which, equilibrated with hemoglobin, result in carboxyhemoglobin (COHb) concentrations of from 3 to 10 percent. A number of investigators have compared COHb levels in smokers and nonsmokers. Goldsmith and Landaw (1969) reported the analysis of expired air samples obtained from 3,311 longshoremen. Using a regression analysis, they calculated the concentration of COHb and found that nonsmokers showed levels of 1.2 percent while those smoking over 2 packs per day had levels of 6.8 percent and that smokers of lesser amounts had intermediate levels. Occupational exposure accounted for the mean nonsmokers' level being over 1.0 percent, an unusual finding in comparison with other studies. Kjeldsen (1969) interviewed and obtained blood samples from 934 CHD-free smokers and nonsmokers. The mean COHb level for 196 nonsmokers was 0.4 percent while all inhaling smokers had a mean level of 7.3 percent. All 416 cigarette smokers, regardless of inhalation or amount smoked, showed a mean level of 4.0 percent.

1005051187

(b) Chevalier et al (1966), cited in the opening paragraph, did not analyze alveolar air. They expressed the opinion that 4% carbon monoxide in cigarette smoke would result in a concentration of 0.04% after dilution. Jongbloed (1939) analyzed the alveolar air of a smoker after the 4th cigarette; the peak level was 0.0031% (see page 19).

The results of Kjeldsen (1969) have been misquoted. The blood level of carboxyhemoglobin in 934 subjects free from coronary heart disease had the following mean value: 0.4% for 196 nonsmokers and 4.2% for 738 smokers. Of the latter group the 121 light smokers had a mean of 2.5%; the moderate smokers one of 4.1% and the heavy smokers one of 5.7%. The figure 7.3% appearing in the paragraph is for atherosclerotic subjects. (Tables from Kjeldsen's paper are reproduced below.)

## Commentary (5) National Clearinghouse, Cont.

Table VI-11

Smoking category	Carboxyhaemoglobin (sat. per cent)			Serum cholesterol (mg/100 ml)		
	Controls	Atherosclerotic subjects	Significance	Controls	Atherosclerotic subjects	Significance
	M ± SD	M ± SD		M ± SD	M ± SD	
Smokers	4,2 ± 3,1 (738)	7,0 ± 3,7 (57)	p < 0,001 (t = 5,52)	247 ± 44 (738)	290 ± 33 (57)	p < 0,001 (t = 4,89)
Non-smokers	0,4 ± 0,9 (196)	0,5 ± 0,7 (2)	n.s. (t = 0,16)	236 ± 49 (196)	284 ± 56 (2)	p = 0,02 (t = 2,32)

Table VI-11: Average values of carboxyhaemoglobin and serum cholesterol in smokers and non-smokers in the control group and in the group with atherosclerotic cardiovascular diseases.

n.s. = not significant at the 0,05 level.

The number of subjects in each smoking category is given below the means and standard deviations.

Table VI-12

Smoking category	Carboxyhaemoglobin (sat. per cent)			Serum cholesterol (mg/100 ml)		
	Controls	Atherosclerotic subjects	Significance	Controls	Atherosclerotic subjects	Significance
	M ± SD	M ± SD		M ± SD	M ± SD	
Light smokers	2,5 ± 2,5 (121)	3,7 ± 2,5 (3)	n.s. (t = 0,76)	245 ± 38 (121)	279 ± 67 (3)	n.s. (t = 1,45)
Moderate smokers	4,1 ± 3,0 (463)	7,3 ± 3,6 (34)	p < 0,001 (t = 4,95)	246 ± 45 (463)	286 ± 50 (34)	p < 0,001 (t = 4,52)
Heavy smokers	5,7 ± 3,0 (154)	7,0 ± 4,0 (20)	n.s. (t = 1,45)	253 ± 45 (154)	298 ± 53 (20)	p < 0,05 (t = 2,18)

Table VI-12: Average values of carboxyhaemoglobin and serum cholesterol related to intensity of smoking in normal smokers and in smokers with atherosclerotic cardiovascular diseases.

n.s. = not significant at the 0,05 level.

The number of subjects in each smoking category is given below the mean values and standard deviations.

Commentary (5) National Clearinghouse, Cont.

c Carbon monoxide has many varied and significant effects on human physiology. An overall review of these effects may be found in a discussion by Lilienthal (127), or more recently in an extensive review by the United States Public Health Service National Air Pollution Control Administration (194). Apart from its effects on respiratory and circulatory function, CO has been found to affect certain central nervous system functions adversely. These effects are probably due to interference by CO with the proper oxygenation and oxidative metabolism of the tissue in question.

CO interferes with oxygen transport in a variety of ways. First, the affinity of hemoglobin for CO is approximately 200 times greater than its affinity for oxygen, and thus CO can easily displace oxygen from hemoglobin. Second, CO shifts the oxyhemoglobin dissociation curve. By increasing the avidity with which oxygen is bound by hemoglobin, CO interferes with O<sub>2</sub> release at the tissue level. This is of greatest importance at the tissue level where the oxygen content of the capillary blood has been reduced to approximately 40 percent saturation. Here the shift can substantially decrease the oxygen tension supplying the tissues.

Third, and of more recent note, is the possible interference by CO with the homeostatic mechanism by which 2, 3-diphosphoglycerate (2, 3-DPG) controls the affinity of hemoglobin for oxygen. Bunn and Jandl (34) have recently reviewed the various experiments concerning this glycolytic intermediate. The question of whether the low levels of CO present in the blood of smokers can affect this homeostasis is presently under investigation (29, 143), and firm conclusions cannot be drawn at this time.

(c) This is an incomplete list of review articles on carbon monoxide. Fifteen others are available (see pages 6 to 10).

d Apart from its effect on hemoglobin affinity, CO appears to induce arterial hypoxemia, and this may act as an additional cause of tissue hypoxia. Ayres, et al. (14, 15) observed unexpectedly that exposure of individuals to CO sufficient to raise their levels of COHb to between 5 and 10 percent was associated with a significant fall in arterial pO<sub>2</sub>. Greater fall in venous pO<sub>2</sub> was noted, but this was considered secondary to increased tissue extraction. In a recent article, Brody and Coburn (30) suggested that this COHb-induced arterial hypoxemia was due to the interaction of a number of factors. These authors noted that in the presence of veno-arterial shunts or of an imbalance in the ventilation-perfusion ratio, the shift in the oxyhemoglobin dissociation curve increased the alveolar-arterial O<sub>2</sub> gradient and resulted in arterial hypoxemia. The presence of shunts as small as 2 percent of cardiac output as well as of approximately 10 percent COHb was found to cause an increase in the gradient. Such ventilation-perfusion (V/Q) abnormalities have recently been noted even in asymptomatic smokers (see Chapter on Chronic Obstructive Bronchopulmonary Disease). The increased levels of COHb found in the blood of smokers may interact with these V/Q abnormalities to further decrease available oxygen.

1005051189

## Commentary (5) National Clearinghouse, Cont.

In normal individuals, coronary flow can increase to meet the increased oxygen demands of a stressed myocardium (as that under nicotine stimulation), while in individuals with severe CHD coronary flow cannot respond as readily. In such cases, myocardial oxygen extraction must be increased above the almost maximal extraction found at rest. Any interference with arterial oxygen levels or hemoglobin affinity could very well decrease available oxygen supplies below the level required for proper tissue function. That this occurs is suggested by the experiments discussed below.

(d) The studies of Ayres et al (1965, 1969) were performed in a non-steady state. These papers are discussed elsewhere in this commentary (page 148 ).

2 Chevalier, et al (17) exposed 10 young nonsmokers to CO concentrations sufficient to induce COHb levels of approximately 4 percent. Taking measurements from blood specimens obtained at cardiac catheterization under resting and exercise conditions, the authors noted that the ratio of oxygen debt to oxygen uptake increased significantly under conditions of increased COHb. According to the investigators this implied that the same work was being done at a greater metabolic cost. These same authors (121, 122) had previously noted similar findings among smokers and observed that cessation of smoking was associated with a significant improvement in oxygen debt accumulation.

(e) The observations of Chevalier et al were made not on blood but on alveolar air analysis. The subjects inhaled 0.5% carbon monoxide, resulting in an indirectly measured blood value of 4% carboxyhemoglobin. There is a serious error here, since other investigators have obtained 25% to 75% carboxyhemoglobin.

3 More recent work by Ayres, et al. (15) has focused on the difference in response to CO exposure between 7 normals and 4 patients suffering from CHD (proven arteriographically). The induction of a COHb concentration of approximately 9 percent in the normals was followed by an increase in coronary blood flow, a decrease in hemoglobin-oxygen percent extraction and no change in myocardial oxygen consumption, coronary sinus oxygen tension, and lactate and pyruvate extraction ratios. The induction of similar COHb levels in the CHD patients was followed by no change in coronary blood flow, a decrease in the hemoglobin-oxygen extraction ratio, and no change in myocardial oxygen consumption. However, these patients did manifest a decrease in coronary sinus pO<sub>2</sub> as well as a decrease in lactate and pyruvate extraction. The latter measures indicate that the myocardium was functioning under hypoxic conditions. Because the coronary flow could not increase and because the myocardium could not extract O<sub>2</sub> from HbO<sub>2</sub>, which was under the influence of CO, coronary sinus oxygen tension decreased to a point which could inactivate certain oxidative enzyme processes. Thus, the myocardial function of persons with CHD may be unable to compensate for the stresses induced by smoking.

1005051190

## Commentary (5) National Clearinghouse, Cont.

(f) The comments in paragraph (d) above apply to another publication by Ayres et al (1969). These measurements were taken in a non-steady state after administering 5% carbon monoxide.

g. Although COHb levels resulting from the CO present in the atmosphere during periods of high air pollution are much lower than those due to the inhalation of cigarette smoke, these concentrations of COHb might contribute to the manifestations of CHD. Cohen, et al. (1972) studied the case fatality rates for patients admitted to 35 Los Angeles area hospitals with myocardial infarction in relation to atmospheric CO pollution. The authors observed an increased MI case fatality rate in areas of increased pollution, and then only during periods of relatively increased CO pollution.

(g) The opening sentence is a poor comparison. Instead of atmospheric air and cigarette smoke a comparison between blood levels of nonsmokers and smokers would be pertinent. There is also an endogenous source of carbon monoxide.

Subsequent to a report by Cohen et al (1969), Haywood et al (1972) examined patients dying of acute myocardial infarction or diverse diseases. There was no clear-cut relationship between carbon monoxide levels and acute infarction (see page 61).

1005051191



- (6) NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH: The health consequences of smoking. A report to the Surgeon General: 1972. 21-3, 121-135, 1972.

### CARBON MONOXIDE

- c Because cigarette smoke contains from 2.7 to 6 percent carbon monoxide (CO), significantly higher carboxyhemoglobin (COHb) levels are found in smokers than nonsmokers (13, 20, 24, 63). COHb levels in nonsmokers are usually less than 1 percent, while those in smokers average around 4 percent and may exceed 15 percent (4, 20, 52). Heavy smokers and those who inhale show the highest carboxyhemoglobin levels (20).

(a) The blood levels of carboxyhemoglobin cited in this paragraph do not represent the values reported in the literature. For nonsmokers the level is not usually less than 1%. In the 26 investigations reported, the overall mean for 1,662 subjects was 1.45% (see pages 32 and 33). For smokers the average is not 4%. The 30 investigations reported in the literature have an overall mean of 3.76% for 2,054 subjects. Values above 15% are very rare and are probably the result of an error in the analysis (see pages 12-15).

- ✓ Haebisch (24) found that a smoker with a daily consumption of 35 to 40 cigarettes easily attains and maintains for hours an alveolar CO concentration of 50 p.p.m., which reaches or exceeds legally-established ambient air quality standards (11, 18, 23, 24).

Cohen, et al. (11) and Aronow, et al. (2) have shown that there is no significant difference in mean expired air carbon monoxide levels after patients have smoked tobacco or lettuce leaf cigarettes. Although pipe and cigar smokers in the United States are reported to have lower exposure to CO than cigarette smokers (20), CO intoxication has been reported in cigar smokers (23).

(b) The observation of Haebisch (1970) is an isolated one. It should be noted that his value of 50 ppm carbon monoxide in alveolar air is 2 to 3 times greater than that reported by Jongbloed (1939) and Ringold et al (1962) (see page 19).

- c CO exerts its adverse effects on the cardiovascular system of smokers through one or more of the following mechanisms: (a) reduction of the amount of hemoglobin available for oxygen transport; (b) shift of the oxygen-hemoglobin dissociation curve to the left with consequent interference in oxygen release at the tissue level; and (c) induction of arterial hypoxemia. CO may interfere with the homeostatic mechanism by which 2,3-DPG controls the affinity of hemoglobin for oxygen (56). CO has also been implicated in experimental atherogenesis in animals (56).

(c) The list of 3 mechanisms actually represents one, which causes the formation of carboxyhemoglobin. Item (c), hypoxemia, meaning low oxygen tension, is not correct. There is low oxygen content but normal tension.

1005051192

Commentary (6) National Clearinghouse, Cont.

Ayres, et al. (2) recently studied 41 patients during diagnostic cardiac catheterization, at which time they inhaled either 5 percent or .1 percent CO. Arterial and mixed venous oxygen tensions were decreased by administration of either concentration. In patients with CHD, coronary artery O<sub>2</sub> extraction decreased 7.9 percent after inhalation of .1 percent CO and 30.5 percent after inhalation of 5 percent CO. Some of the patients with CHD experienced changes in lactate and pyruvate metabolism indicative of inadequate myocardial oxygenation. The higher level of CO inhalation in this experiment is comparable to that experienced intermittently by cigarette smokers.

(d) The work of Ayres et al (1970) is based on a non-steady state and is commented upon elsewhere (see page 148).

Brewer and his colleagues (11) investigated cigarette smoking as a cause of hypoxemia in residents of Leadville, Colorado, at an altitude of 3,100 meters. The arterial pO<sub>2</sub> of 8 smokers was significantly lower ( $P < .05$ ) than that of 12 nonsmokers, but this was reversible upon cessation of smoking. They concluded that the adverse effect of cigarette smoking on O<sub>2</sub> transport may be especially pronounced at high altitude and may restrict an individual's ability to adapt to reduced O<sub>2</sub> tensions (11, 12).

(e) The statement of the results of Brewer et al (1970, 1971) is incomplete. It should be added that the difference in oxygen tension between smokers and non-smokers is small, amounting to 5 mm for pO<sub>2</sub>. More important is the fact that the oxygen saturation for smokers is 74% and is higher than that of the nonsmoking individual (68%). The shift of the oxygen dissociation curve to the left in the blood of smokers accounts for higher oxygen saturation. Therefore, although oxygen tension is reduced, the saturation is higher for smokers. A paragraph from Brewer et al (1971) is quoted below.

1005051193

In the course of our smoking experiments we determined the CO content of the air in our 15 cu m exposure chamber (with approximately one air change/hour) and we found that the CO rose to approximately 20 ppm following the smoking of seven cigarettes in an hour. Particulate matter was 3 mg/cum at the end of the hour. We also measured the concentrations of CO passing across the face of an observer who sat next to a subject for the ten minutes during which a cigarette was smoked. The air was sampled through a tube strapped to the observer's face, and it can be seen from Figure 9 that transient peaks of up to 90 ppm were measured.

7 7  
- 70 p. 134

Commentary (6) National Clearinghouse, Cont.

Kjeldsen (21, 22) examined 998 industrial workers, about one-half of whom were tobacco workers. Fifty-nine cases of arteriosclerosis were documented by such clinical symptoms as angina pectoris and intermittent claudication or by a previous history of myocardial infarction. While 20.9 percent of the 934 "control" individuals were nonsmokers, only 2 (3.4 percent) of the 59 patients with arteriosclerosis were nonsmokers. A significantly higher percentage of diseased workers were heavy smokers and inhaled the smoke.

The diseased smokers had significantly higher carboxyhemoglobin and serum cholesterol levels than either smoking or nonsmoking control patients. This was true after standardizing for differences in levels of smoking between controls and diseased patients. As expected, there was a gradient in carboxyhemoglobin levels from lower levels in light smokers to higher levels in heavy smokers (table 4).

TABLE 4.—Average values of carboxyhemoglobin and serum cholesterol in Danish smokers and nonsmokers in control group and group of patients with arteriosclerotic cardiovascular disease.

Smoking category	Carboxyhemoglobin (saturation percentage)			Serum cholesterol (mg/100 ml)		
	controls M±S.D.	patients M±S.D.	signifi- cance	controls M±S.D.	patients M±S.D.	signifi- cance
Smokers	4.2±3.1 (738)*	7.0±3.7 (57)	p<0.001 t=5.52	247±44 (738)	290±33 (57)	p<0.001 t=4.89
Nonsmokers	0.4±0.9 (196)	0.5±0.7 (2)	n.s. t=0.16	236±49 (196)	284±56 (2)	p<0.02 t=2.32
Light smokers	2.5±2.5 (121)	3.7±2.5 (3)	n.s. t=0.76	245±38 (121)	279±67 (3)	n.s. t=1.45
Moderate smokers	4.1±3.0 (463)	7.3±3.6 (34)	p<0.001 t=4.95	246±45 (463)	286±50 (34)	p<0.001 t=4.52
Heavy smokers	5.7±3.0 (154)	7.0±4.0 (20)	n.s. t=1.45	253±45 (154)	298±53 (20)	p<0.05 t=2.18

p = Probability that difference is not due to chance.

t = Student's t calculation.

n.s. = not significant.

\* The number of subjects in each category is enclosed in parentheses beneath the mean (M) and standard deviation (S.D.).

Source: Kjeldsen, K. (31).

1005051194

Kjeldsen also observed that the COHb levels of 8 to 19 percent seen in 40 percent of the patients with arteriosclerosis were of the same magnitude as those provoking experimental atherosclerosis and cardiac necrosis in animals.

(f) The observations of Kjeldsen (1969, 1970) have not been confirmed by others. It should be stated that Frerovska and Drdkova (1967, 1971) completed a retrospective examination of individuals who had been exposed to an environment of up to 1,000 ppm for an average duration of 10.5 years. There was no early development of arteriosclerosis.

Table 5. Carboxyhemoglobin blood levels of nonsmokers following exposure to vehicular traffic.

Reference (Year)	Nature of subjects	No of Subjects	Carboxyhemoglobin blood levels %		
			Before Mean	After Mean $\pm$ SD (Range)	$\Delta$
Goldsmith, Terzaghi and Hackney (1963)	Los Angeles drivers	1	1.8	2.5	+10
Moureu (1964)	Paris vehicular drivers	597		4.5	
DeBruin, Bult and Van Haeringen (1965)	Amsterdam policemen	10	1.43	1.74	+0.3
DeBruin, Vroeghe and Van Haeringen (1965)	Rotherdam policemen	36	0.93	1.11	+0.08
Morando and Rovida (1965)	Genoa policemen	4		(1.83-4.00)	
Alivisatos, Baza s, Alexopoulos and Verykokakis (1967)	Athens residents	27	0		
Desoille (1967)	Paris garagemen	2		(2.3-3.5)	
Srch (1967)	Prague vehicular passengers	2		(2-5)	
Chovin (1967)	Paris policemen	7		1.25	
Ramsey (1967)	Dayton parking attendant	14	1.5	7.3 $\pm$ 3.46	+5.8
DeBruin (1967)	Amsterdam vehicular drivers	23	1.9	2.15	+0.25
Buchwald (1969)	Alberta garage operators	122		5.0	
Breyse and Bovee (1969)	Seattle fork lift drivers	92		1.2 (0-6)	
Gothe, Fristedt, Sundell, Kolmodin Ehrner-Samuel and Göthe (1969)	Stockholm policemen Malmo policemen Oreho policemen	28 6 3		1.2 $\pm$ 0.39(0.5-2.0) 0.8 $\pm$ 0.14(0.7-1.1) 0.6 $\pm$ 0.38(0.4-1.1)	
Szadkowski, Mastall, Schaller and Lehnert (1970)	Nuhrenberg dustmen	138		4.04 $\pm$ 2.82	
Petrilli and Kanitz (1970)	Genoa vehicular drivers	20		(1.5-3.0)	
Cohen, Dorion, Golds- mith, Permutt(1971)	US-Mexican border inspectors	9	1.5	3.6	
Ayres and Buchler (1970)	New York pedestrians	1481		1.0	
uderitz (1971)	Berlin policemen	120		(4-7)	
Mean (overall for number of subjects)				2.24 (2567 subjects)	+1.12 (84 subjects)

1005051102

Commentary (6) National Clearinghouse, Cont.

Since pipe and cigar smokers inhale less commonly than do cigarette smokers, their contribution to the substances in the air breathed in exposure to smoke pollutants consists of a composite of sidestream smoke and relatively unfiltered mainstream smoke which has been held in the mouth and then expelled.

The actual effluents in the mainstream and sidestream cigarette smoke have been considered by Pascasio, et al. (45) and Scassellati Sforzolini and colleagues (50, 51). These authors stated that "tar" and nicotine levels in sidestream smoke may be significantly higher than those of mainstream smoke and may be harmful to the non-smoker. Actual volume measurements were not reported, however.

Actual measurements of the contamination due to cigarette smoking have been carried out by a number of research groups. A recent, well-controlled study by Harke (24) involved the smoking of 42 cigarettes in 16 to 18 minutes using German blend cigarettes of 85 mm. length, 18 mm. filter, and smoked to a 25 mm. butt length in a room with a volume of 57 cubic meters (approximately the equivalent of a room with a 10-foot ceiling and dimensions of 12 by 14 feet). The author observed that in the absence of ventilation the atmosphere contained up to 50 p.p.m. carbon monoxide and .57 mg./m.<sup>3</sup> nicotine. With substantial ventilation, these levels fell significantly (to approximately 10 p.p.m. carbon monoxide and .10 mg./m.<sup>3</sup> nicotine). He also found that cigar smoke (9 cigars of Clear Sumatra tobacco smoked in 30 to 35 minutes) produced similar amounts of contamination while pipe smoke (2 grams of Navy type medium cut tobacco smoked as eight pipefuls in 35 to 40 minutes) produced much less. Other authors have made similar measurements. Galuskinova (20) found that 3,4-benzpyrene levels in a smoky restaurant were from 2.82 to 14.4 mg./100 m.<sup>3</sup> as compared to outside atmospheric levels of 0.23 to 0.46 mg./100 m.<sup>3</sup>, although burning of food particles may have contributed to the presence of 3,4-benzpyrene in this setting. Kotin and Falk (33) have shown that sidestream cigarette smoke condensate may contain more than three times as much benzo(a) pyrene as mainstream smoke. Srch (55) observed that the smoking of 10 cigarettes to a 5 mm. butt length in an enclosed car of 2.09 m.<sup>3</sup> volume produced carbon monoxide levels up to 90 p.p.m. Lawther and Commins (34), working with a ventilated chamber, found levels of up to 20 p.p.m. of carbon monoxide after seven cigarettes were smoked in one hour; however, peaks of up to 90 p.p.m. were recorded at the seat next to the smoker. Coburn, et al. (9) recorded levels of 20 p.p.m. of carbon monoxide in a small conference room after 10 cigarettes were "burned." Harmsen and Effenberger (25) reported up to 80 p.p.m. of carbon monoxide in an enclosed 98 m.<sup>3</sup> room (approximately the equivalent of a room with a 10-foot ceiling and dimensions of 18 by 20 feet) in which 62 cigarettes had been smoked in two hours.

Another set of contaminants probably present in a tobacco smoke-polluted atmosphere are the oxides of nitrogen. These, specifically NO and NO<sub>2</sub>, have been shown to be present in tobacco smoke although the type most likely to be present in the atmosphere is NO<sub>2</sub>. No measurements have been reported of the amount of NO<sub>2</sub> in smoke-filled rooms. The importance of obtaining and evaluating this information is stressed by the results of Freeman and Haydon and

1005051196

Commentary (6) National Clearinghouse, Cont.

their colleagues (17, 18, 19, 27, 28) and of Blair, et al. (5) who observed bronchial and pulmonary parenchymal lesions in rodents continuously exposed to low levels of  $\text{NO}_2$ .

Other experimenters have measured carboxyhemoglobin (COHb) levels in nonsmokers exposed to cigarette smoke pollutants. Sneh (55) observed that the COHb level in two nonsmokers rose from 2 to 5 percent (that of smokers from 5 to 10 percent) when seated in the cigarette-smoke contaminated car mentioned above (exposure to 90 p.p.m.). Harke (24) reported that when seven nonsmokers were exposed for approximately 90 minutes to a "smoked" room containing 30 p.p.m. of CO there was a rise in COHb from a mean of 0.9 percent to 2.0 percent. In 11 smokers subjected to the same conditions, COHb rose from a mean of 3.3 percent to 7.5 percent. With improved ventilation of the experimental room, the COHb level decreased significantly.

The CO exposures and COHb levels reported above closely approximate the results obtained following experimental chamber exposure of humans to various levels of CO. The uptake of CO by the person depends on, among other parameters: CO concentration, previous COHb level, the level of activity, and the person's state of health. Equilibrium between CO concentration in the lung and in the blood requires over 12 hours exposure. However, as may be noted in table 1, reproduced from Stewart, et al. (56) and derived from measures of COHb in young sedentary males who were not smoking, over half of the equilibrium COHb level is reached within three to four hours of the onset of exposure. The equilibrium value associated with 100 p.p.m. is approximately 14 to 15 percent COHb. Exposure to 100 p.p.m. in the nonsmoker can lead to 3.0 percent of COHb within 60 minutes and 6.0 percent in two hours (16). Of equal significance is that COHb has a half-life of at least three to four hours in the body. As shown in table 1, the COHb level fell only to 2.7 percent in the two hours following cessation of exposure to 50 p.p.m. from the end exposure level of 3.7 percent. This lengthy half-life extends the period of effect of exposure to CO and provides for a buildup of COHb concentration from fresh exposures.

TABLE 1.—Percent of COHb during and following exposure to 50 p.p.m. of CO.

Time during exposure	Mean	Range	Number of subjects
Preexposure	0.7	0.4-1.5	11
30 minutes	1.3	1.3	3
1 hour	2.1	1.9-2.7	11
3 hours	3.8	3.6-4.2	10
6 hours	5.1	4.9-5.5	5
8 hours	5.9	5.4-6.2	5
12 hours	7.0	6.5-7.9	3
15 1/2 hours	7.6	7.2-8.2	3
22 hours	8.5	8.1-8.7	3
24 hours	7.9	7.6-8.2	3
Time without exposure after			
1 hour of exposure			
30 minutes	1.8	1.8	3
1 hour	1.7	1.6-1.8	3
2 hours	1.5	1.4-1.5	3
5 hours	1.1	1.0-1.1	2

1005051197

# Commentary (6) National Clearinghouse, Cont.

Time without exposure after 3 hours of exposure			
30 minutes	3.7	3.4-3.9	3
1 hour	3.3	2.7-3.8	3
2 hours	2.7	2.3-3.0	3
Time without exposure after 8 hours of exposure			
30 minutes	5.6	5.1-5.9	3
1 hour	5.1	4.8-5.4	3
1 1/2 hours	4.0	—	—
11 hours	1.5	1.4-1.7	3
Time without exposure after 24 hours of exposure			
30 minutes	7.5	7.2-7.8	3
1 hour	6.7	6.4-7.1	3
2 hours	5.8	5.6-6.2	3

(g) This section includes an incomplete citation of work by the various investigators. For instance, Harke's conclusion that "it is unlikely to find non-smokers in a room absorbing a significant amount of cigarette smoke" is not mentioned at all. The blood levels that show low absorption are mentioned two paragraphs later. The passive smoker had a blood level of 2% carboxyhemoglobin, an increase of 1.1% over the control level (see page 25).

(h) Srch (1967) not only smoked "in an enclosed car" but also had the motor running in a closed garage. The high levels of carbon monoxide arise largely from automobile exhaust and this is not stated in the quotation (see page 36).

(i) Lowther and Cummins (1970) worked with a ventilated chamber. Examination of the original article reveals that the chamber is 15 cu m, with approximately 1 air change per hour. The high values for carbon monoxide in the air are understandable (see paragraph from article).

Data from 2 representative oxygen tensions are shown in Figure 5 and Table 4. Although there is overlap, the mean saturation values of the normal smokers and the polycythemic smokers are significantly higher than that of non-smoking normals at one of the two oxygen tensions (Table 5). This means that, on the average, the oxygen dissociation curve of smokers in Leadville is left-shifted compared to non-smokers. There was no significant difference between the position of the curve in normal smokers and polycythemic smokers at either oxygen tension.

1005051198

THE EFFECTS OF LOW LEVELS OF  
CARBON MONOXIDE ON HUMAN HEALTH

The data on the effect of low levels of carbon monoxide on human psychological and physiological function have been summarized in two recent publications (3, 53).

There is presently much discussion as to the physiologic and psychophysiological effects of exposure to levels of CO approximating 50 to 100 p.p.m. Beard and Grandstaff (4) observed that exposure to 50 p.p.m. of CO for from 27 to 90 minutes altered auditory discrimination, visual acuity, and the ability to distinguish relative brightness. McFarland (2) observed that COHb levels of 4 to 5 percent caused visual threshold impairment. Ray and Rockwell (48), reporting on a study of the driving ability of three subjects under varying CO exposure, observed that the presence of 10 percent COHb was associated with increased response time for tail-light discrimination and increased variance in distance estimation. Schulte (52) observed that increased errors in cognitive and choice discrimination tests were manifest at levels of COHb as low as 3 percent. Chevalier, et al. (2) have also observed that levels of 4 percent COHb in nonsmokers are associated with an increase in oxygen debt formation with exercise similar to that seen in smokers.

On the other hand, other investigators utilizing complex psychomotor tasks in men and monkeys have observed no decrement in function upon exposures to CO at 50 to 250 p.p.m. (2, 3, 23, 41, 56).

(j) The investigations that concentration of 50 to 250 ppm carbon monoxide does not influence the central nervous system are not described in detail. There are many other reports which are also negative (see Pages 88-90).

Animals exposed to low levels of CO (50 to 100 p.p.m.) continuously for weeks have shown varying degrees of cardiac and cerebral damage similar to that produced by hypoxia (21, 47, 57).

Finally, the possible effects of exposure to 50-100 p.p.m. CO on patients with coronary heart disease (CHD) were investigated by Ayres, et al. (1) who observed a decrease in arterial and mixed venous oxygen tensions with COHb saturations of 5 percent. Certain patients with CHD developed altered lactate and pyruvate metabolism with COHb levels of 5 to 10 percent suggesting myocardial hypoxia.

(k) The cited work of Ayres et al (1970) was performed in a non-steady state with inhalation of 5% carbon monoxide. The validity of these observations is discussed elsewhere in this commentary (see page 143).

1005051199



## Commentary (6) National Clearinghouse, Cont.

¶ The evidence concerning the effect of low levels of carbon monoxide has recently been reviewed and evaluated by the National Air Quality Criteria Committee of the National Air Pollution Control Administration (53). The following is taken from the published conclusions of the Advisory Committee (also see table 2):

"Experimental exposure of nonsmokers to 58 mg/m<sup>3</sup> (50 ppm) for 90 minutes has been associated with impairment in time-interval discrimination. . . . This exposure will produce an increase of about 2 percent COHb in the blood. This same increase in blood COHb will occur with continuous exposure to 12 to 17 mg/m<sup>3</sup> (10 to 15 ppm) for 8 or more hours. . . .

"Experimental exposure to CO concentrations sufficient to produce blood COHb levels of about 5 percent (a level producible by exposure to about 35 mg/m<sup>3</sup> for 8 or more hours) has provided in some instances evidence of impaired performance on certain other psychomotor tests, and an impairment in visual discrimination. . . .

"Experimental exposure to CO concentrations sufficient to produce blood COHb levels above 5 percent (a level producible by exposure to 35 mg/m<sup>3</sup> or more for 8 or more hours) has provided evidence of physiologic stress in patients with heart disease. . . ."

The levels of carbon monoxide found to be present in "smoked" rooms (20 to 80 p.p.m.) are similar to the levels (30 to 50 p.p.m.) which the Advisory Committee has concluded are associated with adverse health effects:

"An exposure of 8 or more hours to a carbon monoxide concentration of 12 to 17 mg/m<sup>3</sup> (10 to 15 ppm) will produce a blood carboxyhemoglobin level of 2.0 to 2.5 percent in nonsmokers. This level of blood carboxyhemoglobin has been associated with adverse health effects as manifested by impaired time interval discrimination. Evidence also indicates that an exposure of 8 or more hours to a CO concentration of 35 mg/m<sup>3</sup> (30 ppm) will produce blood carboxyhemoglobin levels of about 5 percent in nonsmokers. Adverse health effects as manifested by impaired performance on certain other psychomotor tests have been associated with this blood carboxyhemoglobin level, and above this level there is evidence of physiologic stress in patients with heart disease."

These levels of CO are also similar to that set as the time-weighted occupational Threshold Limit Value of 50 p.p.m. for a 40-hour week (five 8-hour days) which has been in effect in the United States for the past several years (15). A further reduction in this limit to 25 p.p.m. is now under consideration. These levels of CO exceed those recently set by the Environmental Protection Agency as the national primary and secondary ambient air quality standards for CO (17). These standards are:

- (a) 10 milligrams per cubic meter (9 p.p.m.)—maximum 8-hours concentration not to be exceeded more than once per year.
- (b) 40 milligrams per cubic meter (35 p.p.m.)—maximum 1-hour concentration not to be exceeded more than once per year.

1005051200

## Commentary (6) National Clearinghouse, Cont.

TABLE 2.—*Effects of carbon monoxide.*

Environmental conditions	Effect	Comment
58 mg./m. <sup>3</sup> (50 p.p.m.) for 90 minutes	Impairment of time-interval discrimination in non-smokers.	Blood COHb levels not available, but anticipated to be about 2.5 percent. Similar blood COHb levels expected from exposure to 10 to 17 mg./m. <sup>3</sup> (10 to 15 p.p.m.) for 8 or more hours.
115 mg./m. <sup>3</sup> (100 p.p.m.) intermittently through a facial mask	Impairment in performance of some psychomotor tests at a COHb level of 5 percent.	Similar results may have been observed at lower COHb levels, but blood measurements were not accurate.
High concentrations of CO were administered for 30 to 120 seconds, and then 10 minutes was allowed for washout of alveolar CO before blood COHb was measured.	Exposure sufficient to produce blood COHb levels above 5 percent has been shown to place a physiologic stress on patients with heart disease.	Data rely on COHb levels produced rapidly after short exposure to high levels of CO; this is not necessarily comparable to exposure over a longer time period or under equilibrium conditions.

SOURCE: Adapted from U.S. Public Health Service, Air Quality Criteria for Carbon Monoxide, Washington, D.C., U.S. Department of Health, Education, and Welfare (53).

(1) The remainder of this section quotes the decision by the National Air Pollution Control Administration, which was drafted mainly for industrial exposures lasting 8 hours. The criteria used by this agency do not apply to brief exposures to cigarette smoke.

### ALLERGIC AND IRRITATIVE REACTIONS TO CIGARETTE SMOKE AMONG NONSMOKERS

(A more detailed discussion of this subject is presented in the Allergy chapter of this report.)

Several investigators have reported on the discomfort and symptoms experienced by both allergic and nonallergic individuals upon exposure to tobacco smoke. Johansson and Ronge (31, 32) in 1965 and 1966 have observed that the acute irritation experienced by nonsmokers in the presence of tobacco smoke is maximal in warm, dry air and that nonsmokers experience more nasal irritation than ocular irritation as compared with smokers exposed to similar amounts of smoke in the atmosphere. Speer (54) studied the reactions of 441 nonsmokers divided into two groups, one composed of individuals with a history of allergic reactions and the other of individuals without such a history. The allergic group underwent skin testing for the presence of sensitivity to tobacco extract while the "nonallergic" group was determined solely by questionnaire concerning subjective allergic responses. Approximately 70 percent of both groups experienced eye irritation while other symptoms differed in their frequency from group to group (nasal symptoms:

1005051201

Commentary (6) National Clearinghouse, Cont.

allergic 67 percent, "nonallergic" 23 percent; headache: allergic 46 percent, "nonallergic" 31 percent; cough: allergic 46 percent, "nonallergic" 25 percent; and wheezing: allergic 22 percent, "nonallergic" 4 percent). Thus, a significant proportion of nonsmoking individuals report discomfort and respiratory symptoms on exposure to tobacco smoke.

Other authors have attempted to separate out those patients who may have specific allergies to smoke. Zussman (61) found that in a random series of 200 atopic patients 16 percent were clinically sensitive to tobacco smoke, and that a majority of these were aided by desensitization therapy. In an earlier study, Pipes (46) observed that 13 percent of 229 patients with respiratory allergy showed positive skin tests to tobacco smoke. Savel (49) has recently reported on eight nonsmokers observed to be clinically hypersensitive to tobacco smoke. After *in vitro* incubation of their lymphocytes with cigarette smoke, increased incorporation of tritiated thymidine was recorded; similar exposure of the lymphocytes of those not sensitive resulted in depression of tritiated thymidine uptake.

Luquette, et al. (39) have recently reported on the immediate effects of exposure to cigarette smoke in school-age children. They observed that heart rate and blood pressure rose with such exposure, although questions remain about the adequacy of their controls and the manner in which the experimental situation may have excited the subjects. Finally, Cameron, et al. (6) observed that acute respiratory illnesses were more frequent among children from homes in which the parents smoked than among children of non-smoking parents. The meaning of these results is uncertain since smoking by the children was not considered and the level of exposure to cigarette smoke in their homes was not measured. Shy, et al. (53) in a study of second grade Chattanooga school children failed to demonstrate a relationship between parental smoking habits and the respiratory illness rates of their children.

(m) This section on allergic reactions refers to components of cigarette smoke other than carbon monoxide. There is no case of reported allergy to carbon monoxide.

#### THE KNOWN HARMFUL EFFECTS OF THE PASSIVE INHALATION OF CIGARETTE SMOKE IN ANIMALS

- 6 A number of investigators have studied the effects of the passive inhalation of high concentrations of cigarette smoke on the pulmonary parenchyma and tracheobronchial tree of animals. The results of these investigations are listed in detail in the recent report to Congress, "The Health Consequences of Smoking," (59) in table 9 of the Bronchopulmonary chapter, and table 16 of the Cancer chapter.

1005051202

Commentary (6) Nationalclearinghouse, Cont.

The pathologic changes observed in the respiratory tract of the animals included parenchymal disruption, bronchitis, tracheobronchial epithelial dysplasia and metaplasia, and pulmonary adenomatous tumor formation. Leuchtenberger, et al. (36) exposed 151 mice to the smoke of from 25 to 1,526 cigarettes over a period of 1 to 23 months and observed that 20 percent of the animals developed severe bronchitis with atypism. Working with 30 control rabbits exposed to up to 20 cigarettes per day for two to five years, Holland, et al. (30) observed increased focal and generalized hyperplasia of the bronchial epithelium and generalized emphysema in the exposed rabbits. Hernandez, et al. (29) observed significantly more pulmonary parenchymal disruption in adult greyhound dogs exposed to cigarette smoke 10 times per week for approximately one year than in nonexposed control animals.

Lorenz, et al. (38) observed no increase in respiratory tract tumor formation above that seen in controls in 97 Strain A mice exposed to cigarette smoke for up to 693 hours. Essenberg (15), however, exposed Strain A mice to cigarette smoke for 12 hours a day for up to one year and observed significantly more papillary adenocarcinomas in the exposed than in the control group. An increased percentage of hybrid mice were found by Mühlbock (42) to have alveolar carcinomas among the experimental group exposed to smoke for two hours a day for up to 684 days when compared with a nonexposed group. Similarly, Guerin (22) observed that 5.1 percent of rats exposed to cigarette smoke for 45 minutes a day for two to six months showed pulmonary tumors compared to 2.4 percent of the control mice.

Leuchtenberger, et al. (37), working with 400 female CF<sub>1</sub> mice, observed only a slight increase in the presence of pulmonary adenomatous tumors among those exposed to cigarette smoke compared with those in the control group. The authors commented that the presence of tumors showed an age relationship independent of smoking exposure. Otto (43) found that 11 percent of a group of albino mice exposed to 12 cigarettes a day for up to 24 months showed pulmonary adenomas as compared with five percent of the control non-exposed group. Dentenwill and Wiebecke (12) found that increasing the exposure of golden hamsters to up to four cigarettes a day for up to two years was associated with an increasing percentage of animals showing desquamative metaplasia and bronchial papillary metaplasia. Harris and Negroni (26) exposed 200 C57BL mice to cigarette smoke for 20 minutes a day every other day for life and found eight adenocarcinomas as compared to none in the control group.

Because the damage observed in these experiments was seen after prolonged exposure to high concentrations of cigarette smoke, and because the comparability of animal exposure to smoke with that of human exposure in smoke-filled rooms is unknown, it is presently impossible to be certain from animal experimentation about the extent of the damage that may occur during long-term intermittent exposure to lower concentrations.

1005051203

Commentary (6) National Clearinghouse, Cont.

## SUMMARY

1. An atmosphere contaminated with tobacco smoke can contribute to the discomfort of many individuals.

2. The level of carbon monoxide attained in experiments using rooms filled with tobacco smoke has been shown to equal, and at times to exceed, the legal limits for maximum air pollution permitted for ambient air quality in several localities and can also exceed the occupational Threshold Limit Value for a normal work period presently in effect for the United States as a whole. The presence of such levels indicates that the effect of exposure to carbon monoxide may on occasion, depending upon the length of exposure, be sufficient to be harmful to the health of an exposed person. This would be particularly significant for people who are already suffering from chronic bronchopulmonary disease and coronary heart disease.

3. Other components of tobacco smoke, such as particulate matter and the oxides of nitrogen, have been shown in various concentrations to adversely affect animal pulmonary and cardiac structure and function. The extent of the contributions of these substances to illness in humans exposed to the concentrations present in an atmosphere contaminated with tobacco smoke is not presently known.

(n) The experiments on animals were not controlled and there is no reported concentration of carbon monoxide. It is not possible to relate the observations on animals to humans unless the concentration of cigarette smoke is known.

1005051204

- (7) ABELSON P H : A damaging source of air pollution. Science 158: 1527, 1967.

One of the toxic products of the automobile is carbon monoxide. Exposure for 1 hour to a concentration of this gas of 120 parts per million causes inactivation of about 5 percent of the body's hemoglobin and commonly leads to dizziness, headache, and lassitude. Concentrations of carbon monoxide as high as 100 ppm often occur in garages, in tunnels, and behind automobiles. Such concentrations are tiny in comparison with those (42,000 ppm) found in cigarette smoke. The smoker survives because most of the time he breathes air not so heavily polluted. However, in a poorly ventilated, smoke-filled room, concentrations of carbon monoxide can easily reach several hundred parts per million, thus exposing smokers and nonsmokers present to a toxic hazard.

In this article, the comparison of concentration of carbon monoxide is as follows: 100 ppm for garages and tunnels and 42,000 ppm in cigarette smoke. The latter represents 4.2 carbon monoxide in pure cigarette smoke. This concentration is not inhaled continuously, but is diluted by air in the lungs at the time of inhalation of cigarette smoke. After exhalation, atmospheric air enters to replace the cigarette smoke. Ringold et al. (1962) analyzed the expired air to determine the integrated concentration of carbon monoxide therein: that of heavy smokers had a concentration of 16.4 ppm, light smokers 7.7 ppm, and non-smokers 0.08 ppm (see page 19).

1005051205

(8) ANON : Carbon monoxide. Am Industr Hyg Ass J 26: 431-4, 1965.

The blood of cigarette smokers will contain from 2% to 10% carboxyhemoglobin and nonexposed adults will show a normal average background of 1% carboxyhemoglobin.<sup>1</sup>

The blood volumes for smokers are overestimated and those for nonsmokers underestimated. The overall mean value reported in 30 investigations is 3.76% for smokers (see page 12). The overall mean value recorded in 26 investigations is 1.45% for nonsmokers.

1005051206

- (9) ANON: Carbon monoxide poisoning - a timely warning. New England J Med  
278: 849-50, 1968.

The cigarette is another producer of carbon monoxide. Heavy cigarette smokers may have as much as 10 per cent carbon monoxide hemoglobin in their blood. Such levels may not be sufficient to cause impairment at sea level but will be enough to produce changes at altitudes of 8000 to 10,000 feet. A recent article has demonstrated behavioral impairment associated with small doses of carbon monoxide<sup>3</sup> - levels of exposure in the range accepted as tolerable in industry. The results indicate that impairment of cerebral function can occur at extremely low levels (50 to 250 ppm) during exposures of half an hour to two and a half hours. Further support for central-nervous-system effects comes from the observation that low levels of carbon monoxide hemoglobin can significantly raise the threshold of light sensitivity of the eye.<sup>4</sup>

That heavy cigarette smoking may cause as much as 10% carboxyhemoglobin in the blood is rare. Barach *et al* (1941) measured the peak levels in the blood of 18 subjects who smoked 20 cigarettes daily and noted a mean level of 5.7%, with a range of 2.2% to 12.3%. Fabre *et al* (1951), in a group of 5 subjects smoking 24 cigarettes, reported a mean level of 4.85%, which represented an increase of 2.7% over the level prior to smoking (see page 21).

1005051207



- (10) ANON : Warning: Cigarettes are dangerous to your health. American Cancer Society. Med Bull Montgomery County Med Soc 24: 45-7, 1968.

Some of these carcinogens may be as simple carbon monoxide — CO — one of the standard by-products of automobile exhaust. Colorless and odorless, this compound of carbon and oxygen is fatal when as little as one-tenth of one percent is breathed in continuously; in minute quantities and repeated doses, it can also be a slow cause of death as a producer of cancer.

Carbon monoxide is not a carcinogen. Carbon monoxide has been tested in mice with tumors induced by chemicals. Carbon monoxide did not influence the rate of tumor growth.

1005051208

(11) ANON: World action on smoking. Brit Med J 4: 65, 1971

Carbon monoxide may be a toxic ingredient of tobacco smoke that deserves more attention than it has received. P. Astrup<sup>1</sup> has recently reported that exposure of rabbits to low concentrations of carbon monoxide can lead to production of atheroma. The carboxyhaemoglobin content of the blood of cigarette smokers may exceed 10%, and Astrup believes that this may be more important than nicotine in relation to coronary disease. Further research on this aspect is certainly needed.

There are two statements that should be modified. Astrup's experiments were performed on rabbits fed with cholesterol and their blood levels showed 15% carboxyhemoglobin (see page 69). That the blood levels of smokers may exceed 10% is a rarity (see pages 20-21).

1005051209

(12) ANON : Cigarette smoking and carbon monoxide. Med Letter Drug Ther  
13: 91-2, 1971.

In addition to nicotine, tars, and other chemical compounds, carbon monoxide has been incriminated as a pathogenic factor in cigarette smoke. Recent studies have suggested that heavy cigarette smoking (more than 20 cigarettes a day) may result in an intake of carbon monoxide that could impair the performance of the smoker in driving a car or piloting an airplane.

The average concentration of carbon monoxide in cigarette smoke is about 20,000 parts per million or about 400 ppm in the inhaled mixture of smoke and air (J. R. Goldsmith and S. A. Landaw, *Science*, 162:1352, 1968). The additive effects of carbon-monoxide-polluted air must also be taken into account. In Los Angeles, where high atmospheric carbon monoxide levels have caused concern, the concentration in the air during a four-year study ranged from 7.3 to 20.2 ppm (A. C. Hexter and J. R. Goldsmith, *Science*, 172:265, 1971).

The reference to the paper by Goldsmith and Landaw (1968) regarding 400 ppm of an inhaled mixture of smoke and air appears in this article. Since Goldsmith and Landaw cite no reference, it is not possible to challenge their source. Jongbloed (1939) analyzed the alveolar air and noted a peak level of 31.5 ppm carbon monoxide. Ringold et al (1962) analyzed the expired air after a 20-second breath-holding period and noted a level of 16.4 ppm for heavy smokers (see page 19).

1005051210

- (13) ARONOW W S, KAPLAN M A and JACOB D : Tobacco: A precipitating factor in angina pectoris. Ann Int Med 69: 529-36, 1938.
- (14) ARONOW W S, DENDINGER J and ROKAW S N : Heart rate and carbon monoxide level after smoking high-, low-, and non-nicotine cigarettes. A study in male patients with angina pectoris. Ann Int Med 74: 697-702, 1971.
- (15) ARONOW W S and ROKAW S N : Nonnicotine cigarettes. Effects in angina pectoris. Circulation 44: 782-8, 1971.
- (16) ARONOW W S, HARRIS C N, ISBELL M W, ROKAW S N and IMPARATO B : Effect of freeway travel on angina pectoris. Ann Int Med 77: 669-76, 1972.

These four publications from Aronow's group have been widely quoted as supporting the theory that carbon monoxide causes coronary heart disease. It should be noted that the investigation concerns carbon monoxide contained in cigarette smoke and in vehicular exhaust. There is no comparative study using carbon monoxide in air to ascertain that the results are due to carbon monoxide contained in vehicular exhaust or cigarette smoke.

Not all four reports include blood analysis for carboxyhemoglobin. In the first, none is recorded, in the second, blood was analysed, in the third, alveolar-expired air was used, and in the fourth, both blood and air had analysis.

1005051211

SIEGEL P V and MOHLER S R : Medical factors in U. S. general aviation accidents.  
Aerospace Med 40: 180-4, 1969.

A 184

STEVENSON P J : The search for the cause of an accident. Proc Roy Soc Med 61: 1076-9, 1968. A 185

VOROSMARTI J Jr, BRADLEY M E, LINAWEAVER P G, KLECKNER J C and ARMSTRONG F W :  
Helium-oxygen saturation diving: I. Hematologic, lactic acid dehydrogenase and carbon  
monoxide-carboxyhemoglobin studies. Aerospace Med 41: 1347-53, 1970.

A 186

1005051252

- (19) AYRES S M: Roles of carbon monoxide and nicotine in circulatory effects of cigarette smoke. JAMA 219: 520, 1972.

### Roles of Carbon Monoxide and Nicotine in Circulatory Effects of Cigarette Smoke

**Q** I would like information about the specific factors that cause cigarette smoking to have an adverse effect on the cardiovascular system. Is it only nicotine that is involved, or can carbon monoxide and its effect on carboxyhemoglobin levels be implicated?

ARTHUR H. SCHLES, MD  
New Rochelle, NY

**A** Until recently, the pharmacology of tobacco smoke was thought to be essentially that of nicotine, and most early articles on the toxicity of cigarette smoke emphasized nicotine to the exclusion of other considerations. It seems likely, however, that the 3% to 4% of carbon monoxide found in cigarette smoke may play an important role intensifying the recognized cardiovascular toxicity of nicotine. Carboxyhemoglobin levels as low as 3% to 4% may increase the oxygen debt of exercise<sup>1</sup> and we have shown that levels of carboxyhemoglobin between five and ten percent may produce abnormal myocardial metabolism in patients with coronary artery disease. Smokers generally have 3% to 7% of their hemoglobin saturated with carbon monoxide. The whole subject has been recently reviewed in a New York Academy of Science monograph on carbon monoxide<sup>2</sup> and in the progress reports from the Surgeon General's office on cigarette smoking and health.

The toxicity of tobacco smoke appears to derive from both its nicotine and carbon monoxide content. Nicotine increases cardiac work by increasing heart rate and blood pressure. Carbon monoxide interferes with the ability of the heart to extract oxygen from the perfusing blood. The combination of increased oxygen requirements and decreased oxygen availability may well lead to myocardial ischemia, particularly in patients with coronary artery disease.

SIMON M. AYRES, MD  
St. Vincent's Hospital  
and Medical Center  
New York

1. Chevalier RB, Krumholz RA, Ross JC: Effects of carbon monoxide inhalation on the cardiopulmonary responses of non-smokers to exercise. *J Lab Clin Med* 62:167, 1953.

2. Colburn RF (ed): *Biological Effects of Carbon Monoxide*. New York, New York Academy of Science, 1970.

1005051213

Commentary (19) Ayres Cont.

The author cites Chevalier *et al*, a reference which appeared in 1963 as an abstract and does not contain results of blood levels of carboxyhemoglobin. In a subsequent paper (*JAMA* 198: 1061-64, 1966) Chevalier *et al* reported the results of inhaling 0.15% carbon monoxide, which caused a carboxyhemoglobin level of 3.95%. This value is lower than results of others, who obtained levels of 10% to 15% of carboxyhemoglobin. Chevalier *et al* used an indirect technique based on analysis of alveolar air, so that it is possible that by this technique they underestimated the true value if blood analysis was used. The technique employed was as follows:

For determination of COHb levels in the blood, the relationship of Haldane and Smith was used.<sup>6</sup> Forster and co-workers,<sup>7</sup> adapted the Haldane relationship for determination of COHb by using a modification of Sjöstrand<sup>7</sup> in which alveolar gas is equilibrated with pulmonary blood and the equilibrated capillary carbon monoxide tension is measured directly in a sample of expired alveolar gas. This method, as utilized by Forster and co-workers,<sup>7</sup> gave results in the same range as those found by analysis of whole venous blood for COHb. Our technique for equilibration of capillary carbon monoxide tension was similar to that described above with certain modifications.<sup>8,9</sup> The Haldane relation states the following (atm signifies atmosphere):

$$\text{COHb/O.Hb} = \frac{210 \text{ CO pressure in atm}}{100 \text{ O}_2 \text{ pressure in atm}}$$

$$\% \text{ COHb} = \frac{\text{O}_2 \text{ pressure in atm}}{210 \text{ CO pressure in atm}} + 1$$

When there is little reduced hemoglobin present, 210 is the M fraction established by Haldane. This technique for determination of COHb showed a satisfactory relationship when compared with the COHb level as determined spectrophotometrically on the venous blood of nine control subjects.

1005051214

- ROBIN E, RAVENS K G and BING R J : Die Wirkung von Alkohol, Nikotin und Zigaretten-  
rauchen auf das Herz. (The effect of alcohol, nicotine and cigarette smoking on the  
heart.) Deutsch Med J 20: 19-29, 1969. 327
- SCHIEVELBEIN H and EBERHARDT R : Cardiovascular actions of nicotine and smoking. J Na  
Canc Ins 48: 1785-94, 1972. 328
- SZÖLLÖSI E, MEDVE F and JENEY E : Angaben zur Wirkung des niedrigen Kohlenmonoxyd-  
Gehaltes in der Luft auf den Menschen. (Data on the effect of a low carbon monoxide con-  
tent in the air on man.) Z Arbeitsmed 20: 263-8, 1970. 329
- TIBBLIN G : Hjärtinfarkt och rökning. (Harmful clinical effects of smoking. Myocardial in-  
farkt and smoking.) Soc Med Tid 2: 65-7, 1971. 330
- WIKTOR Z : Sprawozdanie z posiedzen naukowych wroclawskiego oddzialu towarzystwa intern-  
istow polskich w r, 1952. (Report on the scientific session of the Wroclaw branch of the  
Polish Society of Internal Medicine in 1952. Pol Arch Med Wewn 24: 596-7, 1954. 331
- ZEH E : (Heart function disorders after carbon monoxide or E605 poisoning. Med Welt 1:  
339-40, 1960. 332

1005051129



### Commentary (20) Banyai, Cont.

motor vehicles discharge 66 million tons of carbon monoxide annually. No wonder that in all major cities at busy intersections during hours of peak traffic the concentration of this harmful gas is much higher than the maximum allowable concentration, victimizing drivers, pedestrians and traffic policemen. Carbon monoxide in cigarette smoke is an incomplete combustion product even though the temperature at the burning zone of the cigarette is 884°C (1,635.2°F) while air is being drawn through the cigarette. Carboxyhemoglobin level of the blood is 4-6 percent in moderate smokers and up to 12 percent in heavy smokers. Its potential hazard can be estimated by adding these figures to those pertaining to motor vehicle drivers.

Andrew L. Banyai, M.D.

a. If hyperventilation increases carbon monoxide uptake, it will also hasten its removal from the blood. One technique for promoting the elimination of carbon monoxide in acute poisoning is to increase depth of respiration by having the patient inhale 5% carbon dioxide in oxygen.

b. In urban populations, the carboxyhemoglobin level in the blood is greater than 0.62% to 1.24%. In the 25 investigations reported in the literature the overall mean for 1,662 subjects was 1.45%. Residents of London, Los Angeles and Milan show the following mean levels respectively: 3.5%, 2.3% and 2.8%. These values represent a significant contribution by carbon monoxide to pollution in the atmosphere (see pages 32 and 33).

c. The effects listed in 15 lines are derived from toxic concentrations of carboxyhemoglobin ranging from 50% to 100%. The phrase "toxic concentration" appear only in the first sentence but applies to the next 3 sentences.

d. The experiments in rabbits fed with cholesterol are not supported by those in dogs reported by De Bias *et al* (1972). Chronic exposure to carbon monoxide does not exaggerate myocardial ischemia (see page 62).

e. There is no published report that psychomotor and cognitive areas in the brain can be influenced by levels of carboxyhemoglobin between 2% and 5%. More accurate statistics would be between 5% and 20% (see pages 88-89).

f. The effect of carbon monoxide on alveolar macrophages of experimental animals is encountered with levels of 0.5% to 2% in inspired air. Smokers have a level of carbon monoxide of 7 to 12 ppm in expired air, which is 1000 less than the concentration used in experiments on animals.

g. The levels of 4% to 6% carboxyhemoglobin in moderate smokers and 12% in heavy smokers is not supported by values stated in the literature. Balbo *et al* (1966) reported a mean level of 2.8% for 7 smokers, each consuming 30 cigarettes daily. Rouch *et al* (1971) reported a mean of 4.25% for 15 smokers using more than 10 cigarettes daily (see page 15).

1005051216

- (21) BARTLETT D Jr: Pathophysiology of exposure to low concentrations of carbon monoxide. Arch Environ Health 16: 719-27, 1968.

Regular cigarette smokers have repeatedly been shown to have COHb concentrations in the 5% to 10% range.<sup>27</sup> Smokers of pipes and cigars have COHb levels that are somewhat lower than those of cigarette smokers, but higher than those of nonsmokers. These findings have led to the widespread error of supposing that smokers may be more susceptible to environmental CO than nonsmokers. Carbon monoxide from cigarette smoke and CO in the ambient air are not additive in their biologic effect. Carbon monoxide is absorbed only when the Pco in the ambient air exceeds that in the pulmonary capillary blood. Thus, persons with COHb levels of 5% from smoking do not absorb further CO from the environment unless the ambient CO concentration is 30 ppm or more; on the contrary, they excrete CO at a rate roughly proportional to the Pco gradient between their blood and the ambient air. This suggests that smokers may be among the least susceptible of persons exposed to low atmospheric concentrations of CO, since their COHb concentrations are not increased by the exposure. This conclusion is modified, however, by the fact that smokers' CO excretion between cigarettes is slower in a CO-polluted environment than in pure air. Thus, their long-term average COHb concentrations are slightly higher in the presence of environmental CO than in its absence.

The death rate from coronary heart disease is considerably higher for smokers than for nonsmokers.<sup>27</sup> The rate for exsmokers is no higher than for persons who have never smoked. This pattern implies that the smoking effect is completely reversible when an individual stops smoking. Thus, smoking must cause myocardial hypoxia by some acute, reversible process, probably unrelated to the formation of hard, irreversible, atherosclerotic lesions. Carbon monoxide fits this epidemiologic pattern quite well, but nicotine or other components of cigarette smoke may be responsible, and the question remains unsolved.

The two paragraphs quoted from this article emphasize two points: (1) Carbon monoxide from cigarette smoke and that in the ambient air are not additive in their biologic effect; and (2) the effect of smoking on coronary heart disease is reversible. These two points were missed in the main text of this review and their conception properly belongs to Bartlett.

1005051217

(22) CAMM A J : The effects of smoking. Guy Hosp Gaz 81: 185-203, 1967.

Some three hundred different constituents of tobacco smoke have been identified, many of them in infinitesimally small quantities. The two substances present in the greatest amounts are carbon monoxide and nicotine. Although carbon monoxide is found in high proportions in the mainstream smoke of a cigarette, it is seldom found in high proportions combined with haemoglobin in the blood. The percentage of carboxy-haemoglobin rarely rises above five per cent unless cigarettes are "chain-smoked" in which case it may rise to ten per cent. This is not sufficient to be of clinical significance.

This review article includes a section on "contents of tobacco smoke". The paragraph quoted summarizes the present status of carboxyhemoglobin levels in cigarette smokers, concluding with the statement: "This is not sufficient to be of clinical significance".

1005051218

- (23) CONROY J P : Smoking and the anesthetic risk. Anest Anal 48: 388-400, 1969.

It takes 24 hours in a carbon monoxide free atmosphere to reduce a carboxyhemoglobin of 18 to 5 percent.<sup>31</sup> Is it extravagant to demand 72 smoking-free hours before anesthesia?

The 24 hours required to reduce carboxyhemoglobin in the blood from 18% does not apply to the time required to reduce from 5% the blood level of carboxyhemoglobin in cigarette smokers. In the literature, 4 to 12 hours after smoking, the mean blood level for 2,054 habitual smokers is 3.76%. This value, compared with the level for nonsmokers, represents an increase of 2.19% attributed to smoking. Because of endogenous and exogenous sources of carbon monoxide other than cigarette smoking, it is not possible to reduce the carboxyhemoglobin level below 1.5% (see pages 12-13). An answer to the question raised by Conroy should be as follows: Waiting 72 hours is unreasonable; 4 to 12 hours would be sufficient.

1005051219

- (24) CURPHEY T J : Carboxyhemoglobin in relation to smoking. Nat Cancer Inst Monogr 28: 231-5, 1968.

This is an excellent discussion of the significance of the carboxyhemoglobin level in relation to smoking. The entire article is reproduced.

### Carboxyhemoglobin in Relation to Smoking

THEODORE J. CURPHEY, M.D.,<sup>1</sup> *Chief Medical Examiner—Coroner, Los Angeles County, Los Angeles, California 90012*

THE main thrust of the Conference and the tenor of discussion in the general sessions and in this workshop have been to review and analyze the various agents in tobacco smoke with regard to their potential threat to the health and well-being of the cigarette smoker. The evidence already presented has dealt largely with those effects of certain components of tobacco smoke as they relate to such problems as myocardial infarction, blood coagulation, and carcinogenesis. What can be done to reduce such hazards as "tar" and nicotine, thus leading to the production of a less harmful cigarette, has been discussed.

This afternoon's workshop seems to me to be a variation on the general theme, being in the nature of a movement written in a minor key. It has dealt with certain components in tobacco smoke, e.g., nicotine, whose deleterious properties have not been experimentally and clinically established, but which are nevertheless under various degrees of suspicion. Therefore, these components must be examined in the process of writing the score for the orchestration of Dr. Wynder's symphony, entitled *Toward a Less Harmful Cigarette*.

Carbon monoxide (CO) is one of these components of tobacco smoke that has long been suspected of being harmful and, hence, has received much study over the years.

The problem of CO as a harmful constituent of tobacco smoke raises two questions:

1. Does the amount of CO in the blood differ between the smoker and nonsmoker?
2. If more CO is present in the blood of the smoker, does it produce either functional or structural pathological changes? Are such changes demonstrable by symptomatic, clinical, or laboratory evidence, and can they therefore be assumed to be detrimental to the health or well-being of the smoker as is true in the case of other components of tobacco smoke?


There is abundant evidence in the literature to answer unequivocally the question of the difference between the CO blood level concentration in the smoker and nonsmoker. The article by Larson *et al.* (1) is replete with references covering studies over the past 50 years of CO blood levels in smokers and nonsmokers under various conditions, as well as the effect of various quantity levels of smoking on the CO blood level.

Numerous studies on the normal blood level of CO in the nonsmoker show ranges from 0.5-2.5%. In our study, we used 1% as the normal level.

1005051220

Commentary (24) Curphey, Cont.

The data to be presented on the amount of CO in the blood of smokers are a good example of serendipity. Originally our study was aimed at determining whether there was any correlation between the postmortem CO blood levels of individuals handled by the Los Angeles County Medical Examiner's Office and the CO level of the ambient air at the time of death. We were not considering the cause or mode of death, but were looking for a way to use the CO blood level as a daily indicator of air pollution in the Los Angeles Basin (text-fig. 1).



TEXT-FIGURE 1.—Distribution median of CO in blood of cigarette-only smokers and nonsmokers.

After analyzing our data, we observed a significant association, which, however, was not noted for every location of the monitoring station. Goldsmith *et al.* (2) who had studied the blood CO levels of longshoremen in San Francisco in relation to their smoking habits suggested that the collected data be used to study the smoking habits of this postmortem population.

From November 1–June 30, 1961, 2,267 cases were surveyed, and the data were correlated with 1) the CO concentration of the ambient air at certain monitoring stations in Los Angeles and 2) the smoking habits of the study group (2). To determine the smoking habits of the group, a questionnaire was mailed to the next of kin, when known, or to a known informant. This reduced the group to a total of 1,578 persons, from whom we received usable smoking histories for 1,073 persons.

The 1,073 persons were divided into two groups: 1) nonsmokers (including ex-smokers and persons who never smoked), and 2) smokers. These two groups were further subdivided into (a) those under age 65 and (b) those over age 65. The blood CO levels in the entire group ranged from 0–11.6%. Over 89% of the nonsmokers, regardless of age, fell in the 1% or less CO level. A blood CO level of 5%, regardless of smoking habit, was considered abnormally high.


Forty-six persons had values of 5% and all of these were smokers, except 3 who were ex-smokers. Only 7 of the 46 persons were age 65 or over; in other words, 85% of the persons were in the younger age group. Furthermore, with the use of 1% CO as the normal blood level for nonsmokers, 62% of the nonsmokers had less than this level, whereas only 22% of the smokers had values this low (text-fig. 2). Also, the smokers tended to have a much greater frequency at the extreme values of more than 4%. Moreover, smokers over 65 years old had almost twice as high a percentage value under 1% as the smokers under 65 years (text-fig. 2). The interpretation of this finding offers room for speculation, with one possibility being that older smokers might smoke less than their younger counterparts.

1005051221


Commentary (24) Curphey, Cont.

Another interesting fact gleaned from a study of the observed median CO values is that the values of male nonsmokers were greater than those of female nonsmokers by a factor of nearly 2. On the other hand, for smokers, the distribution by sex did not show consistent differences.

That there is a direct correlation between the height of the CO blood level and the number of cigarettes smoked is a well-established fact, as demonstrated by Goldsmith (2) in his study of a group of San Francisco longshoremen (text-fig. 3). This is seen in the graph of percentage cumulative frequency of expired CO measured in ppm as related to the smoking habits of his study group.



TEXT-FIGURE 2.—Percentage cumulative frequency and percentage of blood CO in smokers and nonsmokers.



TEXT-FIGURE 3.—Distribution of expired CO in longshoremen, by smoking pattern: ILWU study, 1961. \*Percent carboxyhemoglobin concentration based on regression:  $\text{COHb}\% = 0.21 + 0.19 \times (\text{CO ppm})$ .

1005051222

# Commentary (24) Curphey, Cont.

The graphs of Goldsmith's cases of live persons and of our cases show very good correlation in the CO expressed in ppm of expired air with that obtained from a study of postmortem blood expressed in percentage terms of carboxyhemoglobin concentration.

Goldsmith did not correlate the CO blood levels with the general health of his subjects; and obviously in our series, we were denied that opportunity, since the deaths we studied included those from natural causes due to disease and also homicidal, suicidal, and accidental deaths. In point of fact neither of these studies answers the question "Is smoking dangerous to health?"

Fortunately, there is good evidence available which bridges this gap, namely, the study made by Sievers *et al.* (4) of the effect of exposure to known concentrations of CO on a group of 156 police traffic officers, between 32 and 51 years old, who were assigned to duty in the Holland Tunnel for a period of 13 years. These officers were exposed to an average of 70 ppm of CO, which is equivalent to 10% (COHb saturation), with brief exposures up to 260-300 ppm at times and with the heaviest level for a 24-hour period of 66 ppm (14% COHb). Infrequently, the CO level exceeded 260 ppm (32% COHb) and rarely rose as high as 300 ppm (40% COHb) for a few minutes at a time.

This study on police traffic officers is particularly valuable for the purpose of this workshop, for it demonstrated that these men showed no evidence of injury to their health, as determined by serial physical examinations, blood and urine studies, EKG tracings, blood pressure readings, and neurological examinations. In this latter connection, an excellent test for judging the integrity of the nervous system was the pistol marksmanship record of these officers. The Port Authority pistol team was composed of 7 officers, 6 of whom had tunnel duty, and the team consistently finished in first or second place in formal competition with pistol teams from other police organizations for 7 consecutive years.

Even more pertinent to our charge at this time is the study of the smoking habits of the officers in relation to their blood CO levels. Variation in the entire group ranged from 0.5-13.1% saturation, the highest values being obtained in those who smoked and were stationed on the upgrade section of the tunnel and who were exposed to atmospheric CO readings slightly above 100 ppm (16% COHb saturation) for a 2-hour period in contrast to the average daily value of 70 ppm (10% COHb saturation).

What appears to be the most significant observation in this study of traffic officers in the Holland Tunnel is that the blood CO levels of non-smokers in the tunnel on the average exceeded those of smokers in an environment free from any occupational exposure to CO. Since these men remained healthy after being consistently exposed for 13 years to CO levels appreciably higher than those found in tobacco smoke, the conclusion then is inescapable that smokers with CO levels that lie well within these same ranges are similarly unaffected by CO.

## REFERENCES

- (1) LARSON, P. S., HAAQ, H. B., and SILVERSTEIN, H.: Experimental clinical studies. In Tobacco, Baltimore, Williams & Williams Co, 1931, pp 106-110.
- (2) GOLDSMITH, J., SCHURTER, F., and NOVIK, N.: Appraisal of carbon monoxide exposure from analysis of expired air. In Proceedings of XIV International Congress on Occupational Health, Madrid, September 1963.
- (3) CURPHEY, T. J., HOOD, L. D. L., and PIERCE, N. M.: Carboxyhemoglobin in relation to air pollution and smoking: Postmortem studies. Arch Environ Health (Chicago) 10: 170-187, 1965.
- (4) SEEVERS, R. F., EDWARDS, T. L., MURRAY, A. L., and SCHULZ, H. H.: Effect of exposure to known concentrations of carbon monoxide: Study of traffic officers stationed at the Holland Tunnel for 13 years. JAMA 149: 7-8, 1942.

1005051223



(25) DINMAN B D : Carbon monoxide and cigarette smoking. JAMA 212: 1785, 1970.

Although the community is concerned with carbon monoxide body burdens arising from ambient concentrations of this gas, it ignores almost totally the most significant source of carbon monoxide intake—the cigarette. This is paradoxical, since the air of our cities rarely contains more than 30 parts of carbon monoxide per million parts of air (ie, 30 ppm); cigarette smoke streams have been reported to contain from 400 to 40,000 ppm carbon monoxide!

The body itself manufactures carbon monoxide in the course of the breakdown of hemoglobin to the extent of about 1 teaspoonful per day. This small amount of carbon monoxide converts about 0.8% of hemoglobin to inactive carboxyhemoglobin. By contrast, the light smoker converts about 3% of his hemoglobin while the heavy smoker inactivates approximately 8% of this blood pigment. Pipe and cigar smokers rarely achieve such loadings. The body has "learned" to adapt to the small amount of self-produced carbon monoxide over the course of evolution. However, the body burden arising from cigarette smoking probably extends beyond the limit of ready accommodation.

What is the significance of such cigarette-caused carbon monoxide body burdens? It is quite clear that visual acuity at low levels of light intensity is impaired with carbon monoxide loadings in the middle of that range of carboxyhemoglobin levels seen among cigarette smokers. Less clear at this time is the effect of carbon monoxide per se upon cardiac function. However, among patients with heart disease whose ability to accommodate is compromised, at levels of 7% to 9% carboxyhemoglobin, there is deterioration in several cardiac-function indices. On the basis of animal experimental data, it appears that long-term carbon monoxide exposures with about 12% carboxyhemoglobin loading are associated with increased deposition of cholesterol in blood vessels. On the basis

1005051224

Commentary (25) Dinman, Cont.

of epidemiological data, there are some suggestions that this might also apply to humans. In addition, research in mountainous areas suggests that the carbon monoxide loading stemming from cigarette smoking contributes significantly to the development of chronic mountain sickness.

(a) The level of 40,000 ppm carbon monoxide represents pure cigarette smoke which a smoker does not inhale continuously. The concentration of carbon monoxide in the expired air of a heavy smoker is 16.4 ppm and in that of a light smoker 7.7 ppm (see page 19).

(b) The carboxyhemoglobin levels of 3% as estimated for a light smoker and 8% for a heavy smoker are not supported by the available results of investigations. Balbo et al (1966) reported a mean level of 2.8% for 7 smokers consuming 30 cigarettes daily. Rouch et al (1971) recorded a mean level of 4.25% for 15 smokers using more than 10 cigarettes daily (see page 15).

(c) The investigation quoted refers probably to Aronow et al (1971), who noted a reduction in exercise tolerance when patients with angina pectoris smoked low-nicotine cigarettes. The carboxyhemoglobin blood level is elevated to 7.79% but a cause-and-effect relationship has not been demonstrated (see page 62).

(d) The cited work on cholesterol deposition involves rabbits fed with cholesterol in the diet. In dogs, chronically exposed to 100 ppm carbon monoxide, De Bias et al (1972) failed to exaggerate the signs of myocardial ischemia. There is no experimental support for the statement that a blood level of 10% carboxyhemoglobin is harmful to the ischemic heart (see page 62).

1005051225

Commentary (35) Rose, Cont.

When cardiac function is normal there is a significant margin of safety even though the CO intoxication is of long duration. A variety of electrocardiographic aberrations have been observed following CO poisoning, but typically there is a low-voltage pattern.<sup>33</sup> Recovery is apparently rapid following the restoration of oxygen and there is a reversion to a "normal" electrocardiogram; however, enzyme studies may show alterations indicating ischemic heart damage.<sup>32</sup> Epidemiologic studies of cigarette smokers indicate that the death rate from coronary heart disease is considerably higher for smokers than for nonsmokers,<sup>34</sup> and in patients with heart disease there is deterioration of cardiac-function indices at blood levels of seven to nine per cent COHb.<sup>35</sup>

(c) The cited reference for the last sentence is by Dinman (1970), an editorial that is commented upon elsewhere in this report (see page 160).

1005051237

(27) GIEL B G : Air pollution and your lungs. Public Health News 46: 246-53, 1965.

#### FACTS ABOUT SPECIFIC AIR POLLUTANTS

##### *Carbon Monoxide*

Carbon monoxide is well known to all of us. Yet many smokers are unaware that approximately seven to eight percent of their hemoglobin may be bound as carboxyhemoglobin. If in the mean time, such an individual should develop vascular insufficiency to vital organs, and then be forced to breath ambient air containing 30 ppm of CO for four to six hours or gotten into an atmosphere where he would be exposed to 120 ppm CO for one hour, he would bind an additional five percent of his hemoglobin and could suffer tragic results.

The figure of 7% to 8% carboxyhemoglobin among smokers is an over-estimation. A review of the literature shows an overall mean of 3.76% for 2,054 smokers 4 to 12 hours after smoking, and a peak level of 5.26% after smoking (see pages 12 to 14 and 20 to 21).

1005051227

- (28) GOLDSMITH J R : Carbon monoxide and coronary heart disease. Ann Int Med 71: 199-201, 1969.

A list of future investigations relating to carbon monoxide is included in this Editorial.

The completed jigsaw puzzle may give a clear picture showing that the role of smoking in atherosclerotic diseases is mediated in part by carbon monoxide through such mechanisms as those in the preceding paragraph. Missing pieces are needed: [1] the effects of carbon monoxide exposure with and without vasoactive agents on production of anginal symptoms with exercise; [2] prognostic importance of carboxy-hemoglobin levels over both short and long periods of time in cardiovascular diseases; [3] mechanisms and significance of altered hemoglobin binding of oxygen with age, smoking, and other factors; and [4] the possibility of prevention of cardiovascular effects of cigarette smoking when such smoking does not lead to carbon monoxide absorption.

1005051228

- (29) GOLDSMITH J R : Carbon monoxide and coronary heart disease: Compelling evidence in angina pectoris. Ann Int Med 77: 808-10, 1972.

This Editorial is devoted largely to the investigations of Aronow. The 2 paragraphs reproduced herewith contain suggestions for future investigations which have not been discussed elsewhere in this review.

We still lack decisive epidemiologic evidence that there is a risk of more rapid development of coronary heart disease in cigarette smokers with high carbon-monoxide uptake, compared with those of the same age and smoking habits but with low carbon-monoxide uptake. We do, however, have convincing evidence that the death rates from coronary heart disease are higher in cigarette smokers than in non-smokers (6). The 1972 Surgeon General's smoking and health report says, "Experimental and epidemiological investigations implicate the elevation of carboxyhemoglobin levels in smokers as a contributor to the development of CHD and arteriosclerotic peripheral vascular disease." If one estimates the number of excess deaths caused by arteriosclerotic heart disease in smokers in comparison with deaths of people of the same age and sex who are nonsmokers, the potential for prevention is vast. Of the approximately 580,000 deaths in the U.S. caused by arteriosclerotic heart disease, cigarette smoking makes the greatest proportionate contribution to the deaths of those under 60 years of age, doubling the mortality ratios in several studies. If we can substantially reduce this toll of death and of associated disability by reducing carbon monoxide exposure, it would be a massive public health achievement. We are challenged to develop an alteration in cigarette-smoking behavior that does not permit an increase in carboxyhemoglobin to occur.

It is conceivable that with a cigarette that has a catalytic filter or in which the combustion processes are altered there would be less uptake of carbon monoxide by the smoker. Possibly an alteration in the type of tobacco used would have such effects. Difference in tobacco type is credited with the differential effect of smoking cigars or pipes. Smoking these produces relatively little increase in carboxyhemoglobin compared with that from cigarettes. Although much attention is given to cigarettes that are low in tar and nicotine, practically no attention has so far been given to the public health importance of cigarette smoking that produces a low output of carbon monoxide. Such attention is urgently indicated by the evidence that even small increases in carboxyhemoglobin, as we used to think of them, can decrease the work capacity of persons with angina pectoris.

1005051229

- (30) GOLDSTEIN R E and EPSTEIN S E : Medical management of patients with angina pectoris. Prog Cardiovas Dis 14: 360-98; 1972.

#### *Cigarette Smoking*

Patients with angina who smoke a single cigarette before exercise experience significant decreases in the duration of exercise required to precipitate ischemic pain.<sup>31</sup> The decreased exercise capacity after cigarette smoking is associated with a greater heart rate and blood pressure both at rest and after equal amounts of exercise. Since the pressure-rate product was the same at angina before and after smoking, Aronow and Kaplan concluded that smoking increased MVO<sub>2</sub> during exercise, thus precipitating ischemic pain sooner without appreciably altering myocardial oxygen delivery. These authors attributed the alteration in circulatory response to exercise to ganglionic stimulation by nicotine, although subsequent studies<sup>32</sup> using low-nicotine cigarettes yielded the same results as those using ordinary cigarettes. Cigarette smoking might also impair myocardial oxygen delivery in some individuals by converting hemoglobin to carboxyhemoglobin, a change which impairs or destroys the ability of hemoglobin to convey oxygen to tissues. Up to 15% conversion to carboxyhemoglobin can result from heavy smoking.<sup>33</sup>

The last sentence quotes Ayres *et al*, which is actually an abstract. The statement that heavy smoking can produce a blood level of 15% carboxyhemoglobin is not based on any experimental observations.

#### *Carboxyhemoglobin (COHB) and the Access to Oxygen: An Example of Human Counter-Evolution*

STEPHEN M. AYRES, MD, FACC; STANLEY GIANNELLI, Jr., MD, FACC; HILTRUD MUELLER, MD, New York, New York

Cigarette smoking and exposure to community air pollution produce COHB saturations between 3 and 15% and decrease both oxygen capacity and the unloading tension of circulating hemoglobin. An individual with 15% COHB has regressed to a hemoglobin which is functionally intermediate between that of an elephant and a newborn goat.

Acute studies performed in 26 subjects demonstrated that elevation of COHB levels to an average of 7.98% increased cardiac output from 5.01 to 5.56 liters/min, increased minute ventilation from 6.36 to 8.64 liters/min, and decreased arterial and mixed venous oxygen tension from 81 and 39 to 76 and 31 mm Hg, respectively. Myocardial studies performed by coronary sinus catheterization demonstrated that similar elevations of COHB increased coronary blood flow, decreased coronary artery-coronary sinus oxygen content and decreased coronary sinus oxygen tension. The changes were most marked in patients with coronary artery disease or chronic emphysema. Lactate extraction decreasing in 10 of 15 patients. The possibility of adaptation was studied by achieving similar COHB concentrations with both a low and high concentration of carbon monoxide. Hemodynamic changes appeared more marked with administration of the high concentration even though COHB was the same.

These studies suggest that COHB concentrations between 5 and 10% may produce abnormal myocardial function in certain individuals. The well-known deleterious effect of cigarette smoking on the heart may be explained by the interaction of COHB and nicotine, the latter increasing cardiac work and the former decreasing oxygen availability.

1005051230

- (31) LINDQUIST V A Y : Carbon monoxide: Its relationship to air pollution and cigarette smoking. Public Health London '86: 20-6, 1970.

☞ Cigars and pipe tobacco produce more CO than cigarettes, but cigarette smokers tend to have higher levels of COHb because they usually inhale. The gas phase of cigarette smoke contains 1-3% CO (Osborne *et al.*, 1965; Robinson & Niessen, 1961) and the concentrations tend to be highest as the cigarette is smoked down toward the butt-end. Continuous exposure to such ambient concentrations would normally render a man unconscious in a few minutes. However, inhalation of cigarette smoke is both transient and intermittent and the gas is diluted with atmospheric air so that a single cigarette would not be expected to produce an immediate rise in COHb of more than 3%. During the course of a day, those who smoke heavily and inhale usually have mean COHb concentrations in excess of 4% and may even exceed 10% (Ayres *et al.*, 1965; Goldsmith & Landaw, 1968).

- (a) This is a more accurate summary of carboxyhemoglobin levels in cigarette smokers.

#### Myocardial Oxygenation

☞ Those organs with high oxygen consumption leave little "reserve" in the blood supplying them and therefore rely more on increased perfusion to meet any extra demand for oxygen. The myocardium is a typical example. A combination of high oxygen demand, poor perfusion, lowered oxygen capacity and impaired oxygen uncoupling from the blood will obviously prejudice tissue respiration. This was classically demonstrated by Ayres *et al.* (1969, 1970). In a group of noncoronary disease patients undergoing cardiac catheterization raising the COHb level to 9.0% produced a significant increase in coronary perfusion. Despite this, however, the oxygen tension of coronary sinus blood, and presumably of the myocardium itself, dropped slightly. After raising the COHb to a similar degree in a group with established coronary disease, the increase in perfusion was less marked and there was definite reversal of lactate and pyruvate extraction in addition to a drop in oxygen extraction. This indicated significant myocardial hypoxia.

- (b) The experiments of Ayres *et al.* (1969, 1970), consisting of inhalation of carbon monoxide causing blood levels of 9.0% are probably in error. This is discussed elsewhere in the present commentary (see page 148).

1005051231



(32) NAHUM L H : Smoking and thrombosis. Conn Med 29: 853-4, 1965.

- The next problem is to determine by what mechanism does smoking accelerate thrombosis. Employing the method of multiple working hypotheses, a number of possibilities immediately present themselves. One is that smoking produces an increase in carbon monoxide hemoglobin up to 15-20 per cent which somehow promotes thrombosis. Another is that nicotine absorption itself sets into operation the thrombotic process. A third possibility is the well known effect of nicotine in liberating increased amounts of epinephrine which may be the real culprit in accelerating thrombosis. A fourth possibility is that smoke irritation in the bronchial mucosa liberates into the blood stream a thrombus inducing agent. Perhaps there are other possibilities as yet unidentified.

That cigarette smoking produces an increase in carboxyhemoglobin of up to 15% to 20% is a rarity. In a review of the literature, it was found that out of 30 investigations only Meigs (1948) reported a mean level of 16.2% for a group of 6 habitual smokers. The overall mean level for 2,054 subjects reported in 29 investigations was 3.76% (see page 12).

1005051232

- (33) NAHUM L H: Toxic products in cigarette smoke: pleasure or poison.  
Conn Med 32: 154-5, 1963.

Okay, well let's see what it is that the cigarette smoker inhales. Most people when they consider air pollution think of the automobile, the smoke-stack or the trash burners. It's time then to point to a most damaging source of air pollution, the cigarette. One of the toxic products of the automobile exhaust is carbon monoxide (CO). Exposure for one hour to a concentration of this gas of 120 parts per million causes inactivation of about 5 per cent of the body hemoglobin by forming CO hemoglobin. Concentrations of CO as high as 100 ppm, often occur in garages, in tunnels, and behind automobiles. Such concentrations are tiny in comparison with those (12,000 ppm) found in cigarette smoke.

The smoker survives because most of the time he breathes air not so heavily polluted. Nevertheless the smoker can carry 15 to 20 per cent CO hemoglobin for hours and seriously reduce the oxygen supply to already compromised areas in the brain, heart and elsewhere whose arteries are narrowed by atherosclerotic disease. Furthermore, in a poorly ventilated smoke-filled room concentrations of CO can easily reach several hundred parts per million thus exposing smokers and non-smokers present to a toxic hazard. The headache and fatigue that those exposed experience after a time in such an atmosphere is no accident and not psychosomatic.

The concentration of carbon monoxide that a smoker is exposed to continuously is not 42,000 ppm. This is the concentration of pure cigarette smoke which reaches the lung diluted with atmospheric air. The concentration of carbon monoxide in expired air would be a more reasonable estimation. Ringold et al (1962) have reported the following observations with regard to the latter: heavy smokers had a concentration of 16.4 ppm, light smokers 7.7 ppm and non-smokers 0.8 ppm (see page 19).

1005051233

(34) NAHUM L H: The effects of carbon monoxide on human health. Conn Med  
33: 90-2, 1969.

CO. occurs in high concentration in cigarette smoke greater than 2 per cent, this means 20,000 ppm, although an estimate of the average concentration in smoke is much less—400 ppm. In a population of longshoremen smoking produced 6 per cent of COHb. When it comes to occupational exposure 12-14 per cent of employed persons had occupations in which there is a likelihood of exposure. Various forms of indoor combustion may emit CO, and a number of deaths each year are due to poisoning from this source. Gas fired baseboard heaters were incriminated by Michigan State Department of Public Health. Open fires and charcoal braziers produce substantial amounts of CO.

(a) Blood levels of habitual smokers have been reported by 11 groups of investigators. The overall peak after smoking was 5.26% (see page 20). The blood levels 4 to 12 hours after smoking have been reported by 30 groups of investigators with an overall mean of 3.76%. The smoker throughout 24 hours sustains a level between 3.76% and 5.26% (see page 12).

Grut<sup>2</sup> alleged that 46 per cent of 721 drivers had chronic CO. poisoning characterized by fatigue, headache, irritability, dizziness, disturbed sleep and other symptoms. Some subjects had abnormal neurological symptoms. From the epidemiological point of view it is desirable to obtain data which would show whether there are CO. associated increases in such relatively frequent events as motor vehicle accidents or in fatality rates with myocardial infarction to confirm the data from the Los Angeles Hospitals where an association of CO. and case fatality rates in 3,080 patients with myocardial infarction was observed. The central nervous system effects are definitely due to anoxia. The mechanism of myocardial effects probably are similarly produced. Lindenberg<sup>3</sup> did obtain significant electrocardiographic changes on exposing dogs for six weeks to 50 ppm. CO. They also showed dilatation of the right ventricle, scarring of heart muscle and fatty degeneration. A very important question for epidemiologists to study is whether exposure to low concentrations of CO. have a role in the development of human heart disease. Inferring from results of acute toxicologic and experimental studies, we can begin to appreciate the abundant data linking cigarette smoking to coronary heart disease. As far as cigarette smoking is concerned we must keep in mind that high levels of COHb. imply also increased respiratory absorption of other ingredients of tobacco smoke.<sup>4</sup>

1005051234

Commentary (34) Nahum, Cont.

It is safe to say that exposure to CO<sub>2</sub> is widespread, that the smoker who inhales 6 per cent CO<sub>2</sub> is developing a blood concentration which is a serious threat to the health in persons with underlying cardiovascular disease. It is also true that community air pollution may produce COHb in non-smokers similar to that observed in smokers. Even low and commonly occurring CO<sub>2</sub> exposures may impair performance of complex psychomotor tasks. Finally that CO<sub>2</sub> has a role in motor vehicle accidents, is supported by data of high COHb in many drivers involved in accidents.

L.H.N.

## References

1. Goldsmith, J. R. and Landow, S. A.: Carbon monoxide and human health. Science, 162: 1352, December 20, 1968.
2. Grut, A.: Chronic carbon monoxide poisoning. (Munksgaard) Copenhagen, 1969, p. 44.
3. Lindenberg, R., Levy, D., Pierosi, T. and Christensen, M.: paper presented at a meeting of the American Industrial Hygiene Association, Washington, D.C., 1962.
4. Nahum, L. H.: Toxic products in cigarette smoke: pleasure or poison. Conn. Med., 32: 154, March, 1968.

(b) Each statement in this paragraph can be challenged by work of other investigators. For instance, at a meeting Lindenberg et al presented the electrocardiographic changes on exposing dogs for 6 weeks to 50 ppm carbon monoxide. De Bias et al (1972) published the results of their investigations indicating that exposure of dogs to 100 ppm carbon monoxide for 14 weeks does not influence the heart that has been previously infarcted (see page 62).

1005051235

- (35) ROSE E F and ROSE M: Carbon monoxide: A challenge to the physician.  
Clin Med 78: 12-21, 1971.

The amount of CO in cigarette smoke varies between 1 and 2.5 per cent by volume,<sup>4</sup> and can create discomfort not only for the addict but also for others in the environment. For the average smoker, the concentration of CO reaching the alveoli is about 0.01 per cent (400 ppm). The importance of cigarette-produced CO is amply demonstrated by samplings from fleet submarines submerged for extended periods. In 30 hours, concentrations may reach 0.01 per cent (100 ppm),<sup>5</sup> with cigarette smoke accounting for 75 to 80 per cent of the CO production.<sup>6</sup> This exceeds the allowable maximum concentrations recommended by the American Conference of Governmental Industrial Hygienists, which states that CO concentrations in the atmosphere should be kept below 50 ppm.<sup>7</sup>

(a) The concentration of carbon monoxide reaching the alveoli is not 400 ppm for the average smoker. Jongbloed (1939) noted the highest level of 31.5 ppm after a subject finished smoking a 4th cigarette (see page 19).

Because individuals who smoke from 20 to 30 cigarettes daily have a COHb level ranging from three to 10 per cent,<sup>16</sup> there is a widespread mistaken idea that smokers are more susceptible to environmental CO than are nonsmokers. CO from cigarettes and CO in the ambient air are not additive in their biologic effects. CO is absorbed into the bloodstream only when the pressure of CO in the ambient air exceeds that in the pulmonary capillary blood. Thus, persons with COHb levels of five per cent as a result of smoking do not absorb further CO unless the environmental CO concentrations exceed 0.003 to 0.004 per cent.

1005051236

(b) The cited reference 16 is Barach et al (1941). Although the range is 3% to 10%, the mean value is 5.7% for the carboxyhemoglobin level. Other groups of investigators reported mean values of 4.3 and 4.85% (see page 21).

Commentary (35) Rose, Cont.

c' When cardiac function is normal there is a significant margin of safety even though the CO intoxication is of long duration. A variety of electrocardiographic aberrations have been observed following CO poisoning, but typically there is a low-voltage pattern.<sup>33</sup> Recovery is apparently rapid following the restoration of oxygen and there is a reversion to a "normal" electrocardiogram; however, enzyme studies may show alterations indicating ischemic heart damage.<sup>32</sup> Epidemiologic studies of cigarette smokers indicate that the death rate from coronary heart disease is considerably higher for smokers than for nonsmokers,<sup>34</sup> and in patients with heart disease there is deterioration of cardiac-function indices at blood levels of seven to nine per cent COHb.<sup>35</sup>

(c) The cited reference for the last sentence is by Dinman (1970), an editorial that is commented upon elsewhere in this report (see page 160).

1005051237

- (36) SELTZER C C: The effect of cigarette smoking on coronary heart disease. Arch Environ Health 20: 418-23, 1970.

This article reviews the effect of cigarette smoking on coronary heart disease. Seltzer discusses the shortcomings of interpretations appearing in the Surgeon General's report. The question as to the role of carbon monoxide in cigarette smoke is discussed. It should be noted that the work of Ayres quoted in the article has been commented upon elsewhere (see page 148).

6. Does the carbon monoxide constituent of cigarette smoke result in or contribute to increased myocardial infarction or sudden death either in normal individuals or in persons with already impaired coronary circulation due to CHD?

Studies have shown that the carbon monoxide constituent of cigarette smoke does effect increases (2% to 10%) in the levels of carboxyhemoglobin (COHb) saturation when heavy cigarette smokers and nonsmokers were compared, with the consequent displacement of oxyhemoglobin. In addition, carbon monoxide effects a shift to the left of the oxygen-hemoglobin dissociation curve, which may result in a decreased release of oxygen at the tissue level.<sup>21</sup>

On the whole, experimental and clinical investigations bearing on this question are few. The most salient work in this area has been performed by Ayres and associates.<sup>21</sup> In 20 human subjects before and after carbon monoxide inhalation, these investigators found no significant change in oxygen tension. In another experiment, after exposure to carbon monoxide, coronary blood flow increased significantly in seven non-CHD patients but not in four patients with arteriographically proven CHD. In the patients with CHD, myocardial lactate and pyruvate extraction decreased or shifted to actual production, suggesting anaerobic metabolism.

If carbon monoxide does in fact appreciably decrease oxygen extraction at the myocardial level, the matter of oxygen consumption may hinge on the extent of increase in coronary blood flow in normal persons, while in persons with diseased coronary arteries, the increase in blood flow is slight or absent. Hence, it may be a question of the ultimate balance of these opposing forces. In normal persons, there is the presumption that the increased coronary blood flow more than matches the presumed decrease in oxygen extraction. Whether or not this fails to occur in patients with obvious CHD, to such an extent as to "trigger" a coronary event is as yet unknown and much work remains to be done in this area.

1005051238

- (9) ANON: Carbon monoxide poisoning - a timely warning. New England J Med  
278: 849-50, 1968.

The cigarette is another producer of carbon monoxide. Heavy cigarette smokers may have as much as 10 per cent carbon monoxide hemoglobin in their blood. Such levels may not be sufficient to cause impairment at sea level but will be enough to produce changes at altitudes of 8000 to 10,000 feet. A recent article has demonstrated behavioral impairment associated with small doses of carbon monoxide<sup>3</sup> - levels of exposure in the range accepted as tolerable in industry. The results indicate that impairment of cerebral function can occur at extremely low levels (50 to 250 ppm) during exposures of half an hour to two and a half hours. Further support for central-nervous-system effects comes from the observation that low levels of carbon monoxide hemoglobin can significantly raise the threshold of light sensitivity of the eye.<sup>4</sup>

That heavy cigarette smoking may cause as much as 10% carboxyhemoglobin in the blood is rare. Barach et al (1941) measured the peak levels in the blood of 18 subjects who smoked 20 cigarettes daily and noted a mean level of 5.7%, with a range of 2.2% to 12.3%. Fabre et al (1951), in a group of 5 subjects smoking 24 cigarettes, reported a mean level of 4.85%, which represented an increase of 2.7% over the level prior to smoking (see page 21).

1005051207



## Additional Bibliography List, No. 1

## ANALYSIS OF CARBOXYHEMOGLOBIN IN THE BLOOD

- AINSWORTH C A, SCHLOEGEL E L, DOMANSKI T J and GOLDBAUM L R : A gas chromatographic procedure for the determination of carboxyhemoglobin in postmortem samples. J Forensic Sci 12: 529-37, 1967. A 1 Reprint
- ANDERHUB H P, HOFER P and SCHERRER M : Normalwerte der Hb-CO-Sättigung des Blutes. (Normal values of Hb-CO saturation of blood.) Schweiz Med Wochenschr 100: 739-45, 1970. A 2
- ANDERSON S R and ANTONINI E : The binding of carbon monoxide by human hemoglobin. Proof of validity of the spectrophotometric method and direct determination of the equilibrium. J Biol Chem 243: 2918-20, 1968. A 3
- AYRES S M, CRISCITIELLO A and GIANNELLI S Jr : Determination of blood carbon monoxide content by gas chromatography. J Appl Physiol 21: 1368-70, 1966. A 4
- BAUER H J : Beitrag zur Bestimmung von Kohlenoxidhamoglobin. (Contribution to the determination of carbon monoxide hemoglobin). Pharm Zentralhalle Deutsch 104: 25-6, 1965. A 5
- BEECKMANS J M : Determination of the carboxyhaemoglobin saturation of blood by spectrophotometric analysis. Brit J Industr Med 24: 71-2, 1967. A 6
- BJURE J and NILSSON N J : Spectrophotometric determination of oxygen saturation of hemoglobin in the presence of carboxyhemoglobin. Scand J Clin Lab Invest 17: 491-500, 1965. A 7
- BLACKMORE D J : The determination of carbon monoxide in blood and tissue. Analyst 95: 439-58, 1970. A 8
- BREYSSE P A, BOVEE H H and GABAY L F : Comparison of field methods for estimating carbon monoxide hemoglobin percentages. Amer Industr Hyg Ass J 27: 256-9, 1966. A 9
- BUCHWALD H : A rapid and sensitive method for estimating carbon monoxide in blood and its application in problem areas. Amer Industr Hyg Ass J 30: 564-9, 1969. A 10
- CIUHANDU G, RUSU V, DIACONOVICI M and KISS L : Indirekte Bestimmung des Kohlenmonoxyds im Blut. (Indirect determination of carbon monoxide in blood). Z Klin Chem 4: 247-9, 1966. A 11
- COLLISON H A, RODKEY F L and O'NEAL J D : Determination of carbon monoxide in blood by gas chromatography. Clin Chem 14: 162-71, 1968. A 12
- COMMINS B T and LAWTHORP P J : A sensitive method for the determination of carboxyhaemoglobin in a finger prick sample of blood. Brit J Industr Med 22: 139-43, 1965. A 13
- DATSENKO I I : (Determination of the carboxyhemoglobin in the blood with photoelectric colorimeters). Lab Delo 2: 108-10, 1965. A 14
- DOMINGUEZ A M, HALSTEAD J R and DOMANSKI T J : The effect of postmortem changes on carboxyhemoglobin results. J Forensic Sci 9: 330-41, 1964. A 15
- DORSCH J and KOSTER E : Eine Methode zur schnellen Bestimmung von Kohlenmonoxyd im Blut mittels Atlas-Oxymeter. (A rapid test for the determination of carbon monoxide in the blood by means of the Atlas oxymeter). Med Welt 19: 1068-70, 1965. A 16
- EFFENBERGER E : Untersuchungen über den CO-Hb-Gehalt des Blutes in Abhängigkeit von der CO-Konzentration der Luft bei ruhenden Versuchspersonen. (Studies on the COHb content of blood in relation to the CO concentration of the air in subjects at rest). Arch Hyg Bakt 151: 455-74, 1967. A 17

1005051240

## Additional Bibliography List No. 3

## ELIMINATION AND METABOLISM OF CARBON MONOXIDE

- |   | Reprint |
|---|---------|
| ARIN L M and WARNECK P: Reaction of ozone with carbon monoxide. <u>J Phys Chem</u> 76: 1514-6, 1972.  | A 151   |
| COBURN R F, FORSTER R E and KANE P B: Considerations of the physiological variables that determine the blood carboxyhemoglobin concentration in man. <u>J Clin Invest</u> 44: 1899- 910, 1965.  | A 152   |
| COBURN R F, SWERDLOW M, LUGMANMAKI K J, FORSTER R E and POWELL K: Uptake of carbon monoxide from the urinary bladder of the dog. <u>Am J Physiol</u> 215: 1010-23, 1968.  | A 153   |
| COBURN R F: The carbon monoxide body stores. <u>Ann NY Acad Sci</u> 174: 11-22, 1970.   | A 154   |
| FENN W O: The burning of CO in tissues. <u>Ann NY Acad Sci</u> 174: 64-71, 1970.  | A 155   |
| FORSTER R E: Carbon monoxide and the partial pressure of oxygen in tissue. <u>Ann NY Acad Sci</u> 174: 233-41, 1970.  | A 156   |
| LUOMANMAKI K: Studies on the metabolism of carbon monoxide. <u>Ann Med Exp Fenn</u> 44 Suppl 2: 1-55, 1966.   | A 157   |
| LUOMANMAKI K and COBURN R: Effects of metabolism and distribution of carbon monoxide on blood and body stores. <u>Am J Physiol</u> 217: 354-63, 1969.   | A 158   |
| MULHAUSEN R O, ASTRUP P and MELLEMGAAARD K: Oxygen affinity and acid-base status of human blood during exposure to hypoxia and carbon monoxide. <u>Scand J Clin Lab Invest</u> 22 Suppl 103: 9-15, 1968.  | A 159   |
| PATTONO R, MARCHIARO G, CAPELLARO F and ORIONE G: Dinamica della eliminazione del CO in varie condizioni di rianimazione. (Dynamics of elimination of CO in various conditions of resuscitation). <u>Rass Med Industr</u> 33: 456-7, 1964.  | A 160   |
| PETERSON J E and STEWART R D: Absorption and elimination of carbon monoxide by inactive young men. <u>Arch Environ Health</u> 21: 165-71, 1970.   | A 161   |
| POWER G G: Solubility of O <sub>2</sub> and CO in blood and pulmonary and placental tissue. <u>J Appl Physiol</u> 24: 468-74, 1968.   | A 162   |
| TRUHAUT R, BOUDENE C and CLAUDE J R: Recherches sur les effets de l'exposition prolongée du lapin et du rat à de très faibles concentrations d'oxyde de carbone. I. Etude du rythme de fixation et d'élimination du toxique. Discussion de la notion de remanence de l'oxyde de carbone dans le sang (1). (Effects of prolonged exposure of rabbits and rats to very weak carbon monoxide concentrations. I. Study of the rhythm of fixation and elimination of the poison. Discussion of the concept of residual carbon monoxide in the blood). <u>Arch Mal Prof Paris</u> 28: 341-56, 1967. | A 163   |
| WRANNE L: Studies on erythro-kinetics in infancy. VI. A method for the quantitative estimation of pulmonary excretion of carbon monoxide in infancy. <u>Acta Paediat Scand</u> 56: 374-80, 1967.  | A 164   |

1005051250

- LEHMANN J : On the conservation and determination of 5-hydroxytryptamine in blood. Scand J Clin Lab Invest 21: 368-74, 1968. A 35
- LINDERHOLM H : A micromethod for the determination of carbon monoxide in blood. Acta Physiol Scand 64: 372-6, 1965. A 36
- LINDERHOLM H, SJOSTRAND T and SODERSTROM B : A method for determination of low carbon monoxide concentration in blood. Acta Physiol Scand 66: 1-8, 1966. A 37
- MARKIEWICZ J : Badania in vitro nad COHb pochodzenia endogennego w sekcyjnie pobranych probkach krwi. (Studies in vitro on oxidocarbonic hemoglobin of endogenic origin in blood samples taken at autopsy). J Przegl Lek 22: 615-6, 1966. A 38
- McCREDIE R M and JOSE A D : Analysis of blood carbon monoxide and oxygen by gas chromatography. J Appl Physiol 22: 863-4, 1967. A 39
- NOBEL S and RICKER A : Dumbbell-diffusion screening for carbon monoxide, volatile alcohols, and chlorinated hydrocarbons in blood. Clin Chem 13: 276-80, 1967. A 40
- PALMA-CARLOS A G, PALMA-CARLOS M L and SOARES A D : L'etude de la concentration du CO sanguin en quelques hemopathies. (The study of serum CO concentration in some hemopathies). Sangre (Barc) 9: 289-92, 1964. A 41
- RODKEY F L and COLLISON H A : An artifact in the analysis of oxygenated blood for its low carbon monoxide content. Clin Chem 16: 896-9, 1970. A 42
- RODKEY L F, COLLISON H A and O'NEAL J D : Influence of oxygen and carbon monoxide concentration on blood carboxyhemoglobin saturation. Aerosp Med 42: 1274-8, 1971. A 43
- SMALL K A, RADFORD E P, FRAZIER J M, RODKEY F L and COLLISON H A : A rapid method for simultaneous measurement of carboxy- and methemoglobin in blood. J Appl Physiol 31: 154-60, 1971. A 44
- SUNDSTROM G : Enkel COHb-metodik vid hemolytiska tillstand. (Simple COHb method in the hemolytic state) Nord Med 84: 1499, 1970. A 45
- SUNDSTROM G : Blood carboxyhemoglobin: Results with conventional standards compared with those with a submicroliter reference of gaseous CO. Clin Chem 18: 188-92, 1972. A 46
- TAKANO M, GOTO Y, TAKEYA K and SAKAHIGASHI M : (Problems and re-evaluation of quantitative determination of carbon monoxide in the blood). Jap J Clin Pathol 19: Suppl: 505, 1971. A 47
- TAYLOR J D and MILLER J D M : A source of error in the cyanmethemoglobin method of determination of hemoglobin concentration in blood containing carbon monoxide. Amer J Clin Path 43: 265-71, 1965. A 48
- WALD N J and FENTON J : Artefact due to traces of detergent on laboratory glassware affecting carboxyhaemoglobin measurement. Clin Chim Acta 39: 266-8, 1972. A 49
- WEATHERBURN M W and LOGAN J E : Hemoglobinometry. Clin Chem 15: 261-3, 1969. A 50
- WEDERKINCH W F : Semi-micro method for the determination of carboxy-hemoglobin by microdiffusion, reduction of palladium ions and titration of iodide. Scand J Clin Lab Invest 16: 365-71, 1964. A 51
- WIECZOREK H : Bestimmung kleiner Kohlenmonoxidgehalte in biologischem Material. (Determination of small amounts of carbon monoxide in biological material). Mikrochim Acta 4: 802-5, 1968. A 52

1005051242

WOJAHN H: Die CO-Hb-Konzentration in Blutunterlaufungen bei todtlichen Kohlenmonoxyd-vergiftungen. (The CO-Hb concentration in blood suffusions in fatal carbon monoxide poisoning). Deutsch Z Ges Gerichtl Med 59: 99-101, 1967.

A 53

ZORN H: Nachweis von Kohlenmonoxid zur Diagnostik der CO-Vergiftung. (Demonstration of carbon monoxide in the diagnosis of CO poisoning). Dtsch Med Wochensch 94: 1692-5, 1969.

A 54

1005051243

## REACTION OF CARBON MONOXIDE WITH HEMOGLOBIN

- AINSWORTH S and BINGHAM W S W : Reactions of partially oxidised hemoglobin solutions. II. A matrix rank analysis of the initial rates of binding carbon monoxide. Biochim Biophys Acta 160: 10-7, 1968. Reprint  
A 55
- ALBEN J O and CAUGHEY W S : An infrared study of bound carbon monoxide in the human red blood cell, isolated hemoglobin, and heme carbonyls. Biochem 7: 175-83, 1968. A 56
- ALFSEN A, CHIANCONE E, ANTONINI E, WAKS M and WYMAN J : Studies on the reaction of haptoglobin with hemoglobin and hemoglobin chains. III. Observations on the kinetics of the reaction of the haptoglobin-hemoglobin complexes with carbon monoxide. Biochim Biophys Acta 207: 395-403, 1970. A 57
- ALPERT B and BANERJEE R : Photodissociation of carbon monoxide from some new hemoglobin derivatives. A possible case of energy transfer. Biochem Biophys Res Commun 42: 608-14, 1971. A 58
- ANDERSEN M E and GIBSON Q H : A kinetic analysis of the binding of oxygen and carbon monoxide to Lamprey hemoglobin. J Biol Chem 246: 4790-99, 1971. A 59
- ANDERSON N M, REED T A and CHANCE B : A lower limit for the rate of a conformation change in hemoglobin. Ann NY Acad Sci 174: 189-92, 1970. A 60
- ANTONINI E, SCHUSTER T M, BRUNORI M and WYMAN J : The kinetics of the Bohr effect in the reaction of human hemoglobin with carbon monoxide. J Biol Chem 240: 2262-4, 1965. A 61
- ANTONINI E, BUCCI E, FRONTICELLI C, WYMAN J and ROSSI-FANELLI A : The properties and interactions of the isolated  $\alpha$  and  $\beta$  chains of human haemoglobin. III. Observations on the equilibria and kinetics of the reactions with gases. J Mol Biol 12: 375-84, 1965. A 62
- ANTONINI E, BRUNORI M, WYMAN J and NOBLE R W : Preparation and kinetic properties of intermediates in the reaction of hemoglobin with Ligands. J Biol Chem 241: 3236-8, 1966. A 63
- ANTONINI E, CHIANCONE E and BRUNORI M : Studies on the relations between molecular and functional properties of hemoglobin. VI. Observations on the kinetics of hemoglobin reactions in concentrated salt solutions. J Biol Chem 242: 4360-6, 1967. A 64
- ANTONINI E, ANDERSON N M and BRUNORI M : Properties of the product of partial photodissociation of carbon monoxide hemoglobin. J Biol Chem 247: 319-21, 1972. A 65
- APPLEBY C A : Electron transport systems of rhizobium japonicum. I Haemoprotein P-450, other co-reactive pigments, cytochromes and oxidases in bacteroids from  $N_2$ -fixing root nodules. Biochim Biophys Acta 172: 71-87, 1969. A 66
- ASTRUP P : Intracellular 2, 3-diphosphoglycerate and carbon monoxide exposure. Ann NY Acad Sci 174: 252-4, 1970. A 67
- BANERJEE R, DOUZOU P and LOMBARD A : Kinetics of association of carbon monoxide with haemoglobin at low temperature. Nature 217: 23-5, 1968. A 68
- BENESCH R, GIBSON H Q and BENESCH R E : Preliminary communications. Rates of reaction of hemoglobin H with Ligands. J Biol Chem 239: 1668-9, 1964. A 69
- BENESCH R E, MAEDA N and BENESCH R : 2, 3-Diphosphoglycerate and the relative affinity of adult and fetal hemoglobin for oxygen and carbon monoxide. Biochim Biophys Acta 257: 178-182, 1972. A 70
- BETKE K and SHEPARD M K : Verteilung von Kohlenmonoxid auf die Erythrozytenpopulation bei Partialsättigung. Folia Haematologica 89: 135-8, 1968. A 71

1005051244

- BETKE K: Cytological differentiation of haemoglobin. Bibl Haemat 29: 1085-93, 1968. A 72
- BIDE R W and COLLIER H B: A note on the methemoglobin<sup>reductase</sup> activity of rabbit erythrocytes. Canad J Biochem 42: 669-73, 1964. A 73
- BRUNORI M: The carbon monoxide Bohr effect in hemoglobin from *Thunnus thynnus*. Arch Biochem 114: 195-199, 1966. A 74
- BRUNORI M, ANTONINI E, WYMAN J, TENTORI L, VIVALDI G and CARTA S: The hemoglobin of amphibia. VII. Equilibria and kinetics of the reaction of frog hemoglobin with oxygen and carbon monoxide. Comp Biochem Physiol 24: 519-26, 1968. A 75
- BRUNORI M, BONAVENTURA J, BONAVENTURA C, ANTONINI E and WYMAN J: Carbon monoxide binding by hemoglobin and myoglobin under photodissociating conditions. Proc Nat Acad Sci 69: 868-71, 1972. A 76
- CAUGHEY W S, ALBEN J O, McCOY S, BOYER S H, CHARACHE S and HATHAWAY P: Differences in the infrared stretching frequency of carbon monoxide bound to abnormal hemoglobins. Biochem 8: 59-62, 1969. A 77
- CAUGHEY W S: Part III. Biochemical and physiological effects of carbon monoxide. Carbon monoxide bonding in hemoproteins. Ann NY Acad Sci 174: 148-53, 1970. A 78
- CHEAH K S: The membrane-bound carbon monoxide-reactive hemoproteins in the extreme halophiles. Biochim Biophys Acta 197: 84-6, 1970. A 79
- CHRISTIANSEN E and MAGID E: Effects of phosphate, HEPES,  $N_2O$  and CO on the kinetics of human erythrocyte carbonic anhydrases B and C. Biochim Biophys Acta 220: 630-2, 1970. A 80
- CROSETTI L, RUBINO G F and PETTINATI L: Osservazioni in tema di detossicazione dell'emoglobina nell'uomo esposto a rischio prolungato da ossido di carbonio. (Observations on detoxication of hemoglobin in men exposed to prolonged risk of carbon monoxide). Minerva Med 57: 268-9, 1966. A 81
- DINMAN B D, EATON J W and BREWER G J: Effects of carbon monoxide on DPG concentrations in the erythrocyte. Ann NY Acad Sci 174: 246-51, 1970. A 82
- ENGEL R R, RODKEY F L, O'NEAL J D and COLLISON H A: Relative affinity of human fetal hemoglobin for carbon monoxide and oxygen. Blood 33: 37-45, 1969. A 83
- FORMAN H J and FEIGELSON P: Kinetic evidence indicating the absence during catalysis of an unbound ferroporphyrin form of tryptophan oxygenase. Biochem 10: 760-3, 1971. A 84
- FUKUI K and KAKIUCHI Y: The kinetics of the reaction of  $CO + O_2Hb = O_2COHb$  in human-hemoglobin in solution at a low pCO range. Jap J Physiol 20: 332-347, 1970. A 85
- GEDDES R and STEINHARDT J: Factors affecting the release by dilute acid of hidden prototropic groups in carbonylhemoglobin. J Biol Chem 243: 6056-63, 1968. A 86
- GEORGE P and SCHEJTER A: The reactivity of ferrocytochrome c with iron-binding Ligands. J Biol Chem 239: 1504-8, 1964. A 87
- GIBSON Q H, PALMER G and WHARTON D C: The binding of carbon monoxide by cytochrome c oxidase and the ratio of the cytochromes a and  $a_3$ . J Biol Chem 240: 915-20, 1965. A 88
- GIBSON Q H, HELLER P and YAKULIS V: The rate of reaction of carbon monoxide with hemoglobins M. J Biol Chem 241: 1650-1, 1966. A 89
- GIBSON Q H and KAMEN M D: Kinetic analysis of the reaction of cytochrome cc' with carbon monoxide. J Biol Chem 241: 1969-76, 1966. A 90

1005051245

- GIBSON Q H and PARKHURST L J : Kinetic evidence for a tetrameric functional unit in hemoglobin. J Biol Chem 243: 5521-4, 1968. A 91
- GLASS H I, EDWARDS R H T, De GARRETA A C and CLARK J C :  $^{14}\text{C}$  red cell labeling for blood volume and total hemoglobin in athletes: effect of training. J Appl Physiol 26: 131-3, 1969. A 92
- GRAY R D and GIBSON Q H : Kinetic investigation of haemoglobin Bohr effect by flash photolysis. Nature London 226: 77-8, 1970. A 93
- GRAY R D and GIBSON Q H : The binding of carbon monoxide to  $\alpha$  and  $\beta$  chains in tetrameric mammalian hemoglobin. J Biol Chem 246: 5176-8, 1971. A 94
- GRAY R D and GIBSON Q H : The effect of inositol hexaphosphate on the kinetics of CO and  $\text{O}_2$  binding by human hemoglobin. J Biol Chem 246: 7168-74, 1971. A 95
- GRIFFIN J B and HOLLOCHER T C : Evidence for the binding of oxygen and carbon monoxide by succinic dehydrogenase. Biochem Biophys Res Commun 26: 405-10, 1967. A 96
- GUIDOTTI G and KONIGSBERG W : The characterization of modified human hemoglobin. I. Reaction with iodoacetamide and N-ethylmaleimide. J Biol Chem 239: 1474-84, 1964. A 97
- HANISCH G, ENGEL J, BRUNORI M and FASOLD H : Ligand induced conformational changes in various normal and modified hemoglobins as indicated by changes in optical rotatory dispersion. Europ J Biochem 9: 335-42, 1969. A 98
- HAYASHI N, MOTOKAWA Y and KIKUCHI G : Studies on relationships between structure and function of hemoglobin  $\text{M}_{\text{Iwate}}$ . J Biol Chem 241: 79-84, 1966. A 99
- HILDEBRANDT A G, FRANKLIN M R, ROOTS E and ESTABROOK R W : The inhibitory effect of metyrapone on cytochrome P-450-catalyzed mixed-function oxidation reactions as compared to the effect of carbon monoxide. Chem Biol Interact 3: 276-8, 1971. A 100
- HOLLAND R A B : Cell and solution velocity constants for the reaction  $\text{CO} + \text{Hb} = \text{COHb}$  at different temperatures in mammals with different red cell sizes. J Gen Physiol 49: 199-220, 1965. A 101
- HOLLAND R A B : Kinetics of combination of  $\text{O}_2$  and CO with human hemoglobin F in cells and in solution. Res Physiol 3: 307-17, 1967. A 102
- HOLLAND R A B : Rate of  $\text{O}_2$  dissociation from  $\text{O}_2\text{Hb}$  and relative combination rate of CO and  $\text{O}_2$  in mammals at  $37^\circ\text{C}$ . Res Physiol 7: 30-42, 1969. A 103
- HOLLAND R A B : Rate at which CO replaces  $\text{O}_2$  from  $\text{O}_2\text{Hb}$  in red cells of different species. Resp Physiol 7: 43-63, 1969. A 104
- HOLLAND R A B : Reaction rates of carbon monoxide and hemoglobin. Ann NY Acad Sci 174: 154-71, 1970. A 105
- HORIE S : On the absorption spectrum of cytochrome  $\text{a}_3$ . J Biochem 56: 57-71, 1964. A 106
- HORIE S : An alternative method for the calculation of reduced and oxidized absorption spectra of the respiratory enzyme. J Biochem 57: 650-6, 1965. A 107
- HUBER R, EPP O and FORMANEK H : Structures of deoxy- and carbonmonoxy-erythrocyruorin. J Mol Biol 52: 349-54, 1970. A 108
- JEFCOATE C R E and GAYLOR J L : A model for the interaction of cytochrome P-450 with carbon monoxide. J Am Chem Soc 91: 4610-1, 1969. A 109

1005051246

- KEYES M, MIZUKAMI H and LUMRY R: Equilibrium measurement in the reactions of Heme-proteins with gaseous Ligands. Anal Biochem 18: 126-42, 1967. A 110
- KOSMIDER S, ZURKOWSKI Z and WEGIEL A: Wplyw Hg, KCN i CO in vitro na zasadowa fosfataze granulocytow. (In vitro effects of Hg, KCN and CO on granulocyte alkaline phosphatase). Pol Arch Med Wewn 35: 477-81, 1965. A 111
- LANDAW S A, CALLAHAN E W and SCHMID R: Catabolism of heme in vivo: comparison in the simultaneous production of bilirubin and carbon monoxide. J Clin Invest 49: 914-25, 1970. A 112
- LAWSON W H Jr: Effect of anemia, species, and temperature on CO kinetics with red blood cells. J Appl Physiol 31: 447-57, 1971. A 113
- LEVIN W, ALVARES A P and KUNTZMAN R: Distribution of radioactive hemoprotein and CO-binding pigment in rough and smooth endoplasmic reticulum of rat liver. Arch Biochem Biophys 139: 230-5, 1970. A 114
- MacQUARRIE R and GIBSON Q H: Use of a fluorescent analogue of 2, 3-Diphosphoglycerate as a probe of human hemoglobin conformation during carbon monoxide binding. J Biol Chem 246: 5832-5, 1971. A 115
- MANSLEY G E, STANBURY J T and LEMBERG R: Cytochrome oxidase and its derivatives. VI. The CO combining capacity of cytochrome c oxidase. Biochem Biophys Acta 113: 33-40, 1966. A 116
- Mc CONNELL H M, DEAL W and OGATA R T: Spin-labeled hemoglobin derivatives in solution, polycrystalline suspensions, and single crystals. Biochem 8: 2580-5, 1969. A 117
- MEDA E: Cinetica della reazione dell'emoglobina umana con l'ossido di carbonio. Rass Med Industr 33: 292-5, 1964. A 118
- MEL'NICHENKO R K: (Combined effect of carbon monoxide and hydrogen sulfide). Vrach Delo 7: 87-90, 1968. A 119
- MOON R B and RICHARDS J H: Nuclear magnetic resonance studies of CO binding to various heme globins. J Am Chem Soc 94: 5093-5, 1972. A 120
- MORRISON M and HORIE S: Cytochrome c oxidase components. VII. An evaluation of the carbon monoxide-combining capacity in order to determine the stoichiometry of cytochromes a and a<sub>3</sub>. J Biol Chem 240: 1359-64, 1965. A 121
- and WYMAN J
- MURRAY J D: Facilitated diffusion. The case of carbon monoxide. J Biol Chem 246: 5903-6, 1971. A 122
- NAGANO T: Der einfluss von Hitze auf die Affinitat Zwischen Hamoglobin und Kohlenmonoxyd. (Influence of heat on the affinity between hemoglobin and carbon monoxide). Wakayama Med Rep 11: 161-8, 1967. A 123
- NAGEL R L and GIBSON Q H: Kinetics of the reaction of carbon monoxide with the hemoglobin-haptoglobin complex. J Mol Biol 22: 249-55, 1966. A 124
- NAGEL R L, GIBSON Q H and JENKINS T: Ligand binding in hemoglobin J Capetown. J Mol Biol 58: 643-50, 1971. A 125
- NAGEL R L, GIBSON Q H and HAMILTON H B: Ligand kinetics in hemoglobin Hiroshima. J Clin Invest 50: 1772-5, 1971. A 126
- NOBLE R W, BRUNORI M, WYMAN J and ANTONINI E: Studies on the quantum yields of the photodissociation of carbon monoxide from hemoglobin and myoglobin. Biochem 6: 1216-22, 1967. A 127



- NOBLE R W, PARKHURST L J and GIBSON Q H: The effect of pH on the reactions of oxygen and carbon monoxide with the hemoglobin of the carp, *cyprinus carpio*. J Biol Chem 245: 6628-33, 1970. A 128
- OMURA T, SATO R, COOPER D Y, ROSETHAL O and ESTABROOK R W: Function of cytochrome P-450 of microsomes. Fed Proc 24: 1181-9, 1965. A 129
- OSKI F A, GOTTLIEB A J, MILLER W W and DELIVORIA-PAPADOPOULOS M: The effects of deoxygenation of adult and fetal hemoglobin on the synthesis of red cell 2, 3-diphosphoglycerate and its in vivo consequences. J Clin Invest 49: 400-7, 1970. A 130
- PARKHURST L J and GIBSON Q H: The reaction of carbon monoxide with horse hemoglobin in solution, in erythrocytes, and in crystals. J Biol Chem 242: 5762-70, 1967. A 131
- PARKHURST L J, GERACI G and GIBSON Q H: Kinetics of the reaction of hybrid-heme hemoglobins with carbon monoxide. J Biol Chem 245: 4131-5, 1970. A 132
- PASECHNIK V I, SHTAMM E V, VLADISLAVLEV E I and ZAMAYATNIN A A: (Change in the volume of hemoglobin a solution induced by CO ligation). Biofizika 16: 939-41, 1971. A 133
- PAULET G and CHEVRIER R: Influence des ions cobalt sur la combinaison de l'hémoglobine avec l'oxyde de carbone. (Influence of cobalt ions on the combination of hemoglobin with carbon monoxide). C R Soc Biol Paris 160: 1726-7, 1966. A 134
- PETTER C, BOURBON J, MALTIER J P and JOST A: Pathologie du développement. - Production d'hémorragies des extrémités chez le fœtus de Rat soumis à une hypoxie in utero. (Production of hemorrhages of the extremities in the rat fetus subjected to hypoxia in utero). C R Acad Sci Paris 272: 2488-90, 1971. A 135
- PHELPS C and ANTONINI E: The combination of carbon monoxide-haem with apoperoxidase. Biochem J 114: 719-24, 1969. A 136
- PIRNAY F, FASSOTTE F, DEROANNE R and PETIT J M: Capacité de transfert du CO en fonction de la consommation d'O<sub>2</sub> chez l'adulte normal. (The capacity for CO transfer as a function of O<sub>2</sub> consumption by normal adults.) Arch Int Physiol 76: 151-2, 1968. A 137
- POLITZER P: A charge-transfer interpretation of the interactions of hemoglobin with oxygen and carbon monoxide. Biochim Biophys Acta 153: 799-803, 1968. A 138
- RAUSA G, DIANA L and PERIN G: Azione in vitro dell'ossido di carbonio e del piombo sull'attività dell'enzima acido delta - amino - levulinico deidratasi di eritrociti umani. (Action in vitro of carbon monoxide and lead on the delta-aminolevulinic acid dehydratase activity of human erythrocytes). Riv Ital Ig 28: 319-27, 1968. A 139
- REED T: Alterations in the heme-carbon monoxide bond strength. Ann NY Acad Sci 174: 172-6, 1970. A 140
- ROUGHTON F J W: The equilibrium of carbon monoxide with human hemoglobin in whole blood. Ann NY Acad Sci 174: 177-88, 1970. A 141
- RUMEN N M and CHANCE B: Heme-Heme interaction of crystalline lamprey hemoglobin. Kinetics and equilibrium of the reaction of carbon monoxide. Biochim Biophys Acta 207: 404-12, 1970. A 142
- SCHMELZER U, STEINER R, MAYER A, NEDETZKA T and FASOLD H: On the kinetics of the reaction of hemoglobin with CO. Flash - Photolysis experiments. Europ J Biochem 25: 491-7, 1972. A 143
- SIRS J A: Uptake of O<sub>2</sub> and CO by partially saturated haemoglobin in sheep erythrocytes. Biochim Biophys Acta 90: 90-9, 1964. A 144
- TZAGOLOFF A and WHARTON D C: Studies on the electron transfer system. LXII. The reaction of cytochrome oxidase with carbon monoxide. J Biol Chem 240: 2628-33, 1965. A 145

1005051248

- WHARTON D C : Valence state of copper after interaction of the cytochrome oxidase-carbon monoxide complex with ferricyanide. Biochim Biophys Acta 92: 607-9, 1964. A 146
- WINTERHALTER K H, AMICONI G and ANTONINI E : Functional properties of a hemoglobin carrying Heme only on  $\alpha$  chains. Biochem 7: 2228-32, 1968. A 147
- WITTENBERG B A, ANTONINI E, BRUNORI M, NOBLE R, WITTENBERG J B and WYMAN J : Studies on the equilibria and kinetics of the reaction of peroxidases with Ligands. III. The dissociation of carbon monoxide from carbon monoxide ferro-horseradish peroxidase. Biochem 6: 1970-4, 1967. A 148
- WRANNE L : Studies on erythro-kinetics in infancy. XIV. The relation between anaemia and haemoglobin catabolism in Rh-haemolytic disease of the newborn. Acta Paediat Scand 58: 49-53, 1969. A 149
- YAMAZAKI H, OHISHI S and YAMAZAKI I : CO as a Ligand of photosensitive P-630 transformed from horseradish peroxidase in the presence of indole-3 acetic acid. Arch Biochem Biophys 136: 41-6, 1970. A 150

1005051249

## Additional Bibliography List No. 3

## ELIMINATION AND METABOLISM OF CARBON MONOXIDE

- |  | Reprint |
|--|---------|
| ARIN L M and WARNECK P : Reaction of ozone with carbon monoxide. <u>J Phys Chem</u> 76: 1514-6, 1972.  | A 151   |
| COBURN R F, FORSTER R E and KANE P B : Considerations of the physiological variables that determine the blood carboxyhemoglobin concentration in man. <u>J Clin Invest</u> 44: 1899- 910, 1965.  | A 152   |
| COBURN R F, SWERDLOW M, LUGMANMAKI K J, FORSTER R E and POWELL K : Uptake of carbon monoxide from the urinary bladder of the dog. <u>Am J Physiol</u> 215: 1010-23, 1968.  | A 153   |
| COBURN R F : The carbon monoxide body stores. <u>Ann NY Acad Sci</u> 174: 11-22, 1970.   | A 154   |
| FENN W O : The burning of CO in tissues. <u>Ann NY Acad Sci</u> 174: 64-71, 1970.  | A 155   |
| FORSTER R E : Carbon monoxide and the partial pressure of oxygen in tissue. <u>Ann NY Acad Sci</u> 174: 233-41, 1970.  | A 156   |
| LUOMANMAKI K : Studies on the metabolism of carbon monoxide. <u>Ann Med Exp Fenn</u> 44 Suppl 2: 1-55, 1966.   | A 157   |
| LUOMANMAKI K and COBURN R : Effects of metabolism and distribution of carbon monoxide on blood and body stores. <u>Am J Physiol</u> 217: 354-63, 1969.   | A 158   |
| MULHAUSEN R O, ASTRUP P and MELLEMGAAARD K : Oxygen affinity and acid-base status of human blood during exposure to hypoxia and carbon monoxide. <u>Scand J Clin Lab Invest</u> 22 Suppl 103: 9-15, 1968.  | A 159   |
| PATTONO R, MARCHIARO G, CAPELLARO F and ORIONE G : Dinamica della eliminazione del CO in varie condizioni di rianimazione. (Dynamics of elimination of CO in various conditions of resuscitation). <u>Rass Med Industr</u> 33: 456-7, 1964.  | A 160   |
| PETERSON J E and STEWART R D : Absorption and elimination of carbon monoxide by inactive young men. <u>Arch Environ Health</u> 21: 165-71, 1970.   | A 161   |
| POWER G G : Solubility of O <sub>2</sub> and CO in blood and pulmonary and placental tissue. <u>J Appl Physiol</u> 24: 468-74, 1968.   | A 162   |
| TRUHAUT R, BOUDÈNE C and CLAUDE J R : Recherches sur les effets de l'exposition prolongée du lapin et du rat à de très faibles concentrations d'oxyde de carbone. I. Etude du rythme de fixation et d'élimination du toxique. Discussion de la notion de remanence de l'oxyde de carbone dans le sang (1). (Effects of prolonged exposure of rabbits and rats to very weak carbon monoxide concentrations. I. Study of the rhythm of fixation and elimination of the poison. Discussion of the concept of residual carbon monoxide in the blood). <u>Arch Mal Prof Paris</u> 28: 341-56, 1967. | A 163   |
| WRANNE L : Studies on erythro-kinetics in infancy. VI. A method for the quantitative estimation of pulmonary excretion of carbon monoxide in infancy. <u>Acta Paediat Scand</u> 56: 374-80, 1967.  | A 164   |

1005051250

## CARBON MONOXIDE IN POORLY VENTILATED AREAS

- BLACKMORE D J : Interpretation of carboxyhaemoglobin found at post mortem in victims of aircraft accidents. Aerospace Med 41: 757-9, 1970. Reprint A 165
- BLOOM J D : Some considerations in establishing divers' breathing gas purity standards for carbon monoxide. Aerospace Med 43: 633-6, 1972. A 166
- CONKLE J P, MABSON W E, ADAMS J D, ZEFT H J and WELCH B E : Detailed study of contaminant production in a space cabin simulator at 760 Mm of mercury. Aerospace Med 38: 491-9, 1967. A 167
- DILLE J R and MOHLER S R : Drug and toxic hazards in general aviation. Aerospace Med 40: 191-5, 1969. A 168
- EBERSOLE J H : Submarine medicine on Nautilus and Seawolf. Arch Industr Health(Chicago) 18: 200-7, 1958. A 169
- ELLIS W R and SEATONBERRY B W : A radioisotope method for locating points of entry of carbon monoxide into the cabin of an aircraft. Int J Appl Radiat 17: 81-4, 1966. A 170
- HODY G L and BAILEY R W : Weapons exhaust contaminants inside helicopters. Aerospace Med 39: 641-4, 1968. A 171
- JOHNSTON D G and BURGER W D : Injury and disease of scuba and skin divers. Postgrad Med 49: 134-9, 1971. A 172
- JUDD H J : Levels of carbon monoxide recorded on aircraft flight decks. Aerospace Med 42: 344-8, 1971. A 173
- KENT A D : Hazards from products of combustion and oxygen depletion in occupied spaces. Occup Health Rev 21: 1-18, 1970. A 174
- KERTESZ D, ANTONINI E, BRUNORI M, WYMAN J and ZITO R : Studies on the equilibria and kinetics of the reactions of peroxidases with Ligands. I. The reaction of ferropoxidases with carbon monoxide. Biochemistry 4: 2672-6, 1965. A 175
- LIGHTFOOT N F : Chronic carbon monoxide exposure. Proc Roy Soc Med 65: 798-9, 1972. A 176
- LOMONACO T : Breve compendio di medicina aeronautica e spaziale. Continuazione. (Brief compendium of aeronautic and space medicine. Continuation. Minerva Med 62: 3399-34-2, 1971. A 177
- MAGDALENO F M : Tabaco, alcohol y cafe en aviacion. (Tobacco, alcohol and coffee in aviation). Rev Aero Astro 28: 581-5, 1968. A 178
- McFARLAND R A : Human factors in relation to the development of pressurized cabins. Aerospace Med 42: 1303-18, 1971. A 179
- MERLI S, MARCHIORI A, DURANTE F, and MARRACINO F : Ulteriore contributo in tema di identificazione di vittime di disastro aereo. (Further contribution on the identification of air disaster victims). Zacchia 5: 411-37, 1969. A 180
- OKALYI Z : Occupational mortality and morbidity among divers in the Torres Straits. Med J Aust 1: 1239-42, 1969. A 181
- RODKEY F L, COLLISON H A and O'NEAL J D : Influence of oxygen and carbon monoxide concentrations on blood carboxyhemoglobin saturation. Aerospace Med 42: 1274-8, 1971. A 182
- SCHAEFER K E : Environmental physiology of submarines and spacecrafts. Arch Environ Health Chicago 9: 320-31, 1964. A 183

1005051251

SIEGEL P V and MOHLER S R : Medical factors in U. S. general aviation accidents.  
Aerospace Med 40: 180-4, 1969.

A 184

STEVENSON P J: The search for the cause of an accident. Proc Roy Soc Med 61: 1076-9, 1968. A 185

VOROSMARTI J Jr, BRADLEY M E, LINAWEAVER P G, KLECKNER J C and ARMSTRONG F W :  
Helium-oxygen saturation diving: I. Hematologic, lactic acid dehydrogenase and carbon  
monoxide-carboxyhemoglobin studies. Aerospace Med 41: 1347-53, 1970. A 186

1005051252

## ATMOSPHERIC SOURCES OF CARBON MONOXIDE

- ANON: Carbon monoxide changes with the season. New Scientist 56: 625, 1972. Reprint A 187
- DELWICHE C C: Carbon monoxide production and utilization by higher plants. Ann NY Acad Sci 174: 116-21, 1970. A 188
- DOUGLAS E: Carbon monoxide solubilities in sea water. J Phys Chem 71:1931-3, 1967. A 189
- JAFFE L S: Ambient carbon monoxide and its fate in the atmosphere. J Air Pollut Contr Ass 18: 534-40, 1968. A 190
- JAFFE L S: Carbon monoxide in the environment. Sources, characteristics, and fate of atmospheric carbon monoxide. Ann NY Acad Sci 174: 76-88, 1970. A 191
- JUNGE C, SEILER W, BROCK R, GREES K D and RADLER F: Uber die CO-Produktion von mikroorganismen. (Carbon monoxide production of microorganisms). Naturwissenschaften 58: 362-3, 1971. A 192
- MAUGH T H II: Carbon monoxide: Natural sources dwarf man's output. Science 177: 338-9, 1972. A 193
- McMILLAN C and COPE J M: Response to carbon monoxide by geographic variants in acacia farnesiana. Am J Bot 56: 600-2, 1969. A 194
- PICKWELL G V: The physiology of carbon monoxide production by deep-sea coelenterates: causes and consequences. Ann NY Acad Sci 174: 102-5, 1970. A 195
- ROBINSON E and ROBBINS R C: Atmospheric background concentrations of carbon monoxide. Ann NY Acad Sci 174: 89-95, 1970. A 196
- STUPFEL M and GODIN J: Emission d'oxyde de carbone et consommation d'oxygène. (Emission of carbon monoxide and oxygen consumption). C. R. Soc Biol Paris 162: 1684-8, 1969. A 197
- SWINNERTON J W, LINNENBOM V J and LAMONTAGNE R A: The ocean: a natural source of carbon monoxide. Science 167: 984-6, 1970. A 198
- SWINNERTON J W, LINNENBOM V J and LAMONTAGNE R A: Distribution of carbon monoxide between the atmosphere and the ocean. Ann NY Acad Sci 174: 96-101, 1970. A 199
- SWINNERTON J W, LAMONTAGNE R A and LINNENBOM V J: Carbon monoxide in rainwater. Science 172: 943-5, 1971. A 200
- WEINSTOCK B: Carbon monoxide: residence time in the atmosphere. Science 166: 224-5, 1969. A 201
- WEINSTOCK B and NIKI H: Carbon monoxide balance in nature. Science 176: 290-2, 1972. A 202
- WESTBERG K and COHEN N: Carbon monoxide: its role in photochemical smog formation. Science 171: 1013-5, 1971. A 203
- WOLFGANG R: Carbon monoxide as a basis for primitive life on other planets. Nature 225: 876, 1970. A 204

1005051253

## ENDOGENOUS ORIGIN OF CARBON MONOXIDE

- COBURN R F, WILLIAMS W J and KAHN S B : Endogenous carbon monoxide production in patients with hemolytic anemia. J Clin Invest 45: 460-8, 1966. Reprint A 205
- COBURN R F : Endogenous carbon monoxide production and body CO stores. Acta Med Scand Suppl 472: 269-82, 1967. A 206
- COBURN R F, WILLIAMS W J, WHITE P and KAHN S B : The production of carbon monoxide from hemoglobin in vivo. J Clin Invest 46: 346-56, 1967. A 207
- COBURN R F : Endogenous carbon monoxide production. New Eng J Med 282: 207-9, 1970. A 208
- COBURN R F : Enhancement by phenobarbital and diphenylhydantoin of carbon monoxide production in normal man. New Eng J Med 283: 512-15, 1970. A 209
- COBURN R F, WALLACE H W and ABBOUD R : Redistribution of body carbon monoxide after hemorrhage. Am J Physiol 220, 868-73, 1971. A 210
- COLTMAN C A Jr, DUDLEY G M III and LEVERETT S D Jr : Measurement of endogenous carbon monoxide production to determine the effect of high  $+G_x$  acceleration on the destruction rate of red cells. Aerospace Med 40: 627-31, 1969. A 211
- COLTMAN C A Jr and DUDLEY G M III : The relationship between endogenous carbon monoxide production and total heme mass in normal and abnormal subjects. Am J Med Sci 258: 374-85, 1969. A 212
- DESOILLE H, CASTILLON du PERRON M, CREMER G and LEBBE J : Absence de corrélation entre la pression partielle d'oxygène et l'oxyde de carbone dans le sang. A propos de l'oxycarbonisme endogène décrit par M. Loeper. Arch Mal Prof Paris 24: 603-7, 1963. A 213
- DESOILLE H, CREMER G and GIRARD C : Au sujet d'un prétendu oxycarbonisme d'origine endogène chez les diabétiques. (On the subject of a supposed blood carbon monoxide of endogenous origin in diabetics). Arch Mal Prof Paris 26: 625-30, 1965. A 214
- DRABKIN D L : The hemophagous organ of the placenta and in vitro studies of endogenous carbon monoxide production. Ann NY Acad Sci 174: 49-63, 1970. A 215
- FALLSTROM S P and BJURE J : Endogenous formation of carbon monoxide in newborn infants. II. Rh haemolytic disease of the newborn. Acta Paediat Scand 56: 365-73, 1967. A 216
- FALLSTROM S P and BJURE J : Endogenous formation of carbon monoxide in newborn infants. III. ABO Incompatibility. Acta Paediat Scand 57: 137-44, 1968. A 217
- FALLSTROM S P : Endogenous formation of carbon monoxide in newborn infants. IV. On the relation between the blood carboxyhaemoglobin concentration and the pulmonary elimination of carbon monoxide. Acta Paediat Scand 57: 321-9, 1968. A 218
- FALLSTROM S P : Endogenous formation of carbon monoxide in newborn infants. V. On the relation between the carboxyhaemoglobin concentration and the haemoglobin catabolism calculated from simultaneous determinations of carbon monoxide elimination and total haemoglobin. Acta Paediat Scand 57: 487-94, 1968. A 219
- FALLSTROM S P : On the endogenous formation of carbon monoxide in full-term newborn infants. Acta Paediat Scand Suppl 189: 1-27, 1969. A 220
- LANDAW S A and WINCHELL H S : Endogenous production of carbon-14 labeled carbon monoxide: an in vivo technique for the study of heme catabolism. J Nucl Med 7: 696-707, 1966. A 221
- LANDAW S A : Endogenous production of carbon monoxide: the human body as a cause of air pollution. Ann Int Med 70: 1275-6, 1969. A 222

1005051254

Additional Bibliography List No. 10

- SANNA-RANDACCIO F and NISSARDI G P : L'influenza della ventilazione sulla misura del 'trasfert' del CO in stato stabile. (Influence of ventilation on the measurement of 'transfer' of CO in stable state.) Boll Soc Ital Biol Sper 45: 762-5, 1969. A 442
- SARTORELLI E : Alterazioni della diffusione alveolo-capillare del CO nella silicosi. (Changes in alveolo-capillary diffusion of CO in silicosis.) Acta Geront (Milano) 17: 182-6, 1967. A 443
- SATAKE T, HIDA S, USUI A, TATSUBANA K, YAMAZAKI J and MATSUOKA T : (Studies on the measurement of the carbon monoxide diffusing capacity by the single-breath method, with special reference to the influence of the uneven distribution of ventilation-perfusion of the lungs.) Jap Circ J 32: 35-40, 1968. A 444
- SHAW D B, CINKOTAI F and THOMSON M L : Syncope induced by application of negative pressure to the lower body and its effect on lung CO diffusing capacity. Aerospace Med 37: 154-7, 1966. A 445
- SIKAND R S and PIPPER J : Pulmonary diffusing capacity for CO in dogs by the single breath method. Resp Physiol 1: 172-92, 1966. A 446
- SIMONE M, REGGIANI A and BET E : La capacità di diffusione per il monossido di carbonio nella valutazione preoperatoria. (Carbon monoxide diffusion capacity in preoperative evaluation.) Acta Anaesth (Padova) 16: Suppl 4: 103-14, 1965. A 447
- STEINER S H, FRAYSER R and ROSS J C : Alterations in pulmonary diffusing capacity and pulmonary capillary blood volume with negative pressure breathing. J Clin Invest 44: 1623-30, 1965. A 448
- TARTULIER M, TOURNIAIRE A, DEYRIEUX F and BLUM J : Étude de transfert alvéolo-capillaire de l'oxyde de carbone dans la sténose mitrale. (Study of alveolo-capillary transfer of carbon monoxide in mitral stenosis.) Arch Mal Coeur 60: 843-64, 1967. A 449
- TLUSTÝ L, HLOUŠKOVÁ Z, KROFTA K and DAUM S : Difúzní kapacita plicní pro CO, vyšetřena metodou 'steady state' a metodou 'single breath'. (Pulmonary diffusing capacity for CO examined by 'steady state' and 'single breath' method.) Cas Lek Cesk 111: 149-53, 1972. A 450
- TRINQUET G, CLAUZEL A M, CARRÉ R and MEYER A : Les valeurs du transfert de l'oxyde de carbone chez l'homme normal en état ventilatoire stable au repos. (Values for carbon monoxide transfer in normal humans in stable ventilatory state at rest.) J Franc Med Chir Thorac 21: 365-82, 1967. A 451
- VANDENBERGH E, BILLIET L, WOESTIJNE K P and GYSELEN A : Relation between single-breath diffusing capacity and arterial blood gases in chronic obstructive lung disease. Scand J Resp Dis 49: 92-101, 1968. A 452
- VANROUX R and GREGOIRE M : La capacité de diffusion au CO chez le bronchiteux chronique. (The CO diffusion capacity in chronic bronchitis.) Acta Tuberc Belg 55: 488-97, 1964. A 453
- WAGNER P D, MAZZONE R W and WEST J B : Diffusing capacity and anatomic dead space for carbon monoxide ( $C^{18}O$ ). J Appl Physiol 31: 847-52, 1971. A 454
- WAGNER W W Jr, LATHAM L P, BRINKMAN P D and FILLEY G F : Pulmonary gas transport time: larynx to alveolus. Science 163: 1210-1, 1969. A 455
- WEBER J, MORET P and CHAUVET M : L'utilité du test de diffusion au monoxyde de carbone dans l'exploration fonctionnelle du poumon. (Utility of the diffusion test with carbon monoxide in functional exploration of the lung.) Helv Med Acta 34: Suppl: 101-2, 1967. A 456
- WEISKOPF R B and SEVERINGHAUS J W : Diffusing capacity of the lung for CO in man during acute acclimation to 14,246 ft. J Appl Physiol 32: 285-9, 1972. A 457
- WOLKONSKY P M : Pulmonary effects of air pollution. Arch Environ Health 19: 586-92, 1969. A 458
- WOOLF C R : An assessment of the fractional carbon monoxide uptake and its relationship to pulmonary diffusing capacity. Dis Chest 46: 181-9, 1964. A 459

1005051269



## CARBON MONOXIDES AS AIR POLLUTANT.

- |  | Reprint |
|--|---------|
| ABELSON P H : Man-made environmental hazards. I. How man shapes his environment. <u>Am J Public Health</u> 58: 2043-9, 1968.   | A 240   |
| AMERICAN THORACIC SOCIETY : Air pollution and health. <u>Am Rev Res Dis</u> 93: 302-12, 1966.  | A 241   |
| ANDERSON D O : The effects of air contamination on health: a review. <u>Canad Med Ass J</u> 97: 585-93, 1967.  | A 242   |
| ANON : Carbon monoxide - a stealthy killer. <u>Anesth Analg Cleveland</u> 45: 664-8, 1966.   | A 243   |
| AYRES S M : The effects of air pollution on health. <u>Delaware Med J</u> 41: 9-14, 1969.  | A 244   |
| and<br>AYRES S M, EVANS R G / BUCHLER M E : Air pollution: A major public health problem. <u>CRC Crit Rev Clin Lab Sci</u> 3: 1-40, 1972.  | A 245   |
| CARNOW B W : Air pollution and physician responsibility. <u>Arch Int Med</u> 127: 91-5, 1971.  | A 246   |
| CLAYTON G D : Discussion. Carbon monoxide and lead: an environmental appraisal. <u>J Air Pollut Contr Ass</u> 19: 688-9, 1969.   | A 247   |
| EISENBUD M and EHRLICH L R : Carbon monoxide concentration trends in urban atmospheres. <u>Science</u> 176: 193-4, 1972.   | A 248   |
| ELFIMOVA E V and KHACHATURYAN M K : Features specific to the reflex action of sub-threshold concentrations of sulphurous anhydride in combination with phenol and carbon monoxide in the atmosphere. <u>Gig Sanit</u> 33: 3-6, 1968. | A 249   |
| ENVIRONMENTAL PROTECTION AGENCY: National primary and secondary ambient air quality standards. <u>Fed Reg</u> 36: 8186-8201, 1971.   | A 250   |
| EPSTEIN S S : Toxicologic and epidemiologic bases for air quality criteria. <u>J Air Pollut Contr Ass</u> 19: 629-31, 1969.  | A 251   |
| FELDSTEIN M : Toxicity and analysis of air pollutants. <u>J Forensic Sci</u> 14: 337-51, 1969.   | A 252   |
| GIEVER P M and RUCH W E : Statistical analyses of air pollution sampling frequency in an epidemiologic study. <u>Am Indus Hyg Ass J</u> 32/4: 260-6, 1971.   | A 253   |
| GOLDSMITH J R : Environmental epidemiology and the metamorphosis of the human habitat. <u>Am J Pub Health Na Health</u> 57:1532-49, 1967.  | A 254   |
| GRUNDY R D : Toxicologic and epidemiologic bases for air quality criteria. Carbon monoxide and lead. Session summary. <u>J Air Pollut Contr Ass</u> 19: 729-32, 1969.  | A 255   |
| HEXTER A C and GOLDSMITH J R : Carbon monoxide: association of community air pollution with mortality. <u>Science</u> 172: 265-7, 1971.  | A 256   |
| LANGMANN R : Schadklima. (The injurious climate). <u>Hippokrates</u> 35: 588-96, 1964.   | A 257   |
| MORROW P E : Adaptations of the respiratory tract to air pollutants. <u>Arch Environ Health</u> 14: 127-36, 1967.  | A 258   |
| SCORER R S : Review paper. New attitudes to air pollution - the technical basis of control. <u>Atmos Environ</u> 5: 903-34, 1971.  | A 259   |
| SOKOLOVSKY M S, PINCHUK E M and PRAVOVEROV K N : Hygienic assessment of air and microclimate in gas-supplied kitchens. <u>Gig Sanit</u> 32: 16-20, 1967.   | A 260   |
| STOKINGER H E : The spectre of today's environmental pollution - USA brand: New perspectives from an old scout. <u>Am Indus Hyg Ass J</u> 30: 195-217, 1969.   | A 261   |

1005051256

- TONOMURA M, YAMATE N and TSUJI K : Observations of air pollution with the aid of automatic continuous analysers. II. Relationship among the concentrations of various kinds of pollutants in atmosphere. Bull Nat Inst Hyg Sci Tokyo 85: 35-40, 1967. A 262
- TOYAMA T : Air pollution and its health aspect in Tokyo areas. Asian Med J 11: 5-15, 1968. A 263
- WHO EXPERT COMMITTEE: Atmospheric pollutants. Wld Hlth Org Techn Rep Ser 271: 3 - 16, 1963. A 264
- WHO EXPERT COMMITTEE: Environmental change and resulting impacts on health. Wld Hlth Org Techn Rep Ser 292: 3-24, 1964. A 265
- WHO SCIENTIFIC GROUP: Research into environmental pollution. Wld Hlth Org Techn Rep Ser No. 406: 34-44, 1968. A 266
- WHO EXPERT COMMITTEE: Urban air pollution with particular reference to motor vehicles. Wld Hlth Org Techn Rep Ser 410: 3-53, 1969. A 267
- WHO CHRONICLE: Public health aspects of climate in cities. WHO Chron 25: 161-7, 1971. A 268
- WOLKONSKY P M : Pulmonary effects of air pollution. Arch Environ Health 19: 586-92, 1969. A 269

1005051257

VEHICULAR EXHAUST AS SOURCE OF CARBON MONOXIDE

- ANDERSON D E : Problems created for ice arenas by engine exhaust. Am Ind Hyg Assoc J 32: 790-801, 1971. A 270
- ANON : Air pollution: CO not OK in Europe. Nature (Lond) 227: 3-4, 1970. A 271
- AOYAMA M : Effect of automotive exhaust on the living body. Jap J Clin Pathol 18: 690-693, 1970. A 272
- FISHER R S, MASEMORE W C and SOPHER I M
- BAKER S P : Fatal unintentional carbon monoxide poisoning in motor vehicles. Am J Public Health 62: 1463-1467, 1972. A 273
- BARTEK P, GAUME J G and ROSTAMJI H J : Dynamics analysis for time of useful function (TUF) predictions in toxic combusive environments. Aerosp Med 41: 1392-1395, 1970. A 274
- BARTH D S : Federal motor vehicle emission goals for CO, HC and NOx based on desired air quality levels. J Air Pollut Control Assoc 20: 519-524, 1970. A 275
- BLUMER W : Verkehrsabgase als ursache von kopfschmerzen. Praxis 59: 201-209, 1970. A 276
- BOCCADORO S and INGIULLA M : Determinazione dell'ossido di carbonio come indice di inquinamento atmosferico da traffico motorizzato nei negozi e nelle abitazioni del centro di Firenze. Ann Sanita Pubblica 29: 1507-1522, 1968. A 277
- BOVÉ J L and SIEBENBERG S : Airborne lead and carbon monoxide at 45th Street, New York City. Science 167: 986-987, 1970. A 278
- BRANDT A : Über die herabsetzung des CO-gehaltes der auspuffgase von verbrennungsmotoren. Dtsch Gesundheitsw 20: 1202-1206, 1965. A 279
- BRICE R M and ROESLER J F : The exposure to carbon monoxide of occupants of vehicles moving in heavy traffic. J Air Pollut Control Assoc 16: 597-600, 1966. A 280
- BYSTROM G : En okonventionell lösning av parkerings- och kommunikationsproblem inom stadsdelen Orrholmen, Karlstad. Nord Hyg Tidskr 51: 20-24, 1970. A 281
- CAMPBELL J M : A Breath of New York. Science 159: 693-694, 1968. A 282
- CHICHIKALO D I, BENEVELSKY I A and MINSKY I A : On 2 cases of poisoning by carbon monoxide fumes inside automobiles. Cig Sanit 31: 76-77, 1966. A 283
- DUNLAP R : Carbon monoxide: the silent killer. Today's Health 26-27, 67-68, and 71-72, 1961. A 284
- GOLDSMITH J R and ROGERS L H : Health hazards of automobile exhaust. Public Health Report 74: 551-558, 1959. A 285
- GOLDSMITH J R and DEANE M : Outdoor workers in the United States and Europe. Milbank Mem Fund Q 43: 107-116, 1965. A 286
- HAAGEN-SMIT A J : Carbon monoxide levels in city driving. Arch Environ Health 12: 548-551, 1966. A 287
- INGIULLA M, GRASSO C and MARIOTTINI E : Ulteriori determinazioni dell'ossido di carbonio quale indice di inquinamento atmosferico. (Further determinations of carbon monoxide as an indication of atmospheric pollution.) Is Med 61: 582-591, 1968. A 288

1005051258

- JOHNSON K L, DWORETZKY L H and HELLER A N : Carbon monoxide and air pollution from automobile emission in New York City. Science 160: 67-68, 1968. A 289
- LANDAU E, SMITH R and LYNN D A : Carbon monoxide and lead: an environmental appraisal. J Air Pollut Control Assoc 19: 684-687, 1969. A 290
- LANGMANN R and KETTNER H : Die problematik einer chronischen intoxication durch kohlenoxyd und untersuchungen seiner immissionen in grobstadten. (Problems of chronic carbon monoxide intoxication and studies on the emission in cities). Oeff Gesundheitswes 30: 7-11, 1968. A 291
- LE BRETON R and GARAT J : Intoxications oxycarbonées aiguës provoquées par les clés ou coupe-tirage. (Acute carbon monoxide poisonings caused by valves or dampers). Ann Med Leg (Paris) 44: 366-368, 1964. A 292
- LUDERITZ P : Experimentelle untersuchungen über die wirkungen von kohlenmonoxid aus kraftfahrzeugabgasen auf den gesundheitszustand von verkehrspolizisten. (Experimental studies in the effect of carbon monoxide from automobile exhausts on the health of traffic policemen). Z Gesamte Hyg 17: 645-646, 1971. A 293
- MIRKIN A J : The role of the physician in accident prevention. Arch Environ Health 13: 519-524, 1966. A 294
- MURPHY S D, LENG J K, ULRICH C E and DAVIS H V : Effects on animals of exposure to auto exhaust. Arch Environ Health 7: 60-70, 1963. A 295
- NISHII S : (Effects of air pollution due to automobile exhaust on the human body.) Naika 21: 813-819, 1968. A 296
- PERRELLI G and ROSETTANI E : Criteri di valutazione del rischio ambientale da co. (Criteria of evaluation of the environmental risk from carbon monoxide). Folia Med (Napoli) 47: 1062-1067, 1964. A 297
- PERRELLI G, ROSETTANI E and BRAGUZZI E : Valutazione del rischio negli esposti all'in-alazione di piccole quantità di ossido di carbonio. (Evaluation of the risk in exposure to inhalation of small quantities of carbon monoxide). Minerva Med 56: 3477-3479, 1965. A 298
- PETROVA A, DALAKMANSKI Y and BAKALOV D : Study of contamination of the atmosphere by injurious road transport and industrial products. J Hyg Epidemiol Immunol (Praha) 10: 383-391, 1966. A 299
- RAMSEY J M : Concentrations of carbon monoxide at traffic intersections in Dayton, Ohio. Arch Environ Health 13: 44-46, 1966. A 300
- RAUSA G, PERIN G and DIANA L : Inquinamento da ossido di carbonio, piombo e ossidi di azoto in una autorimessa a più piani del Veneto. (Air pollution with carbon monoxide, lead and nitrogen oxides in an automobile parking garage several stories high in Venetia). Riv Ital Ig 27: 402-422, 1967. A 300
- RAYFIELD J D : Field testing program for carbon monoxide. J Med Assoc State Ala 36: 1494-1502, 1967. A 301
- REED L E and TROTT P E : Continuous measurement of carbon monoxide in streets 1967-1969. Atmos Environ 5: 27-39, 1971. A 302
- RISPLER L and ROSS C R : Ventilation for engine exhaust gases. Occup Health Rev 17: 19-22, 1965. A 303
- RONDIA D, GUYAUX J and HEUSGHIEM C : Concentration de l'oxyde de carbone dans les rues, à Liège. (Concentration of carbon monoxide in the streets of Liège). Arch Belg Med Soc 24: 73-87, 1966. A 304

1005051259

- SFORZOLINI G S and SAVINO A : Azione dell'ozono su alcuni indici chimici di inquinamento degli ambienti confinati. (Action of ozone on several chemical indices of air pollution in closed areas). Riv Ital Ig 29-30: 174-188, 1970, Italian. A 305
- SHOJI H, YAMAMOTO T, NISHIDA K, ISHIKAWA Y, TAKADA S and INOUE K : (Studies on air pollution owing to the automobile exhaust gases. The concentrations of C1-C6 hydrocarbons and some inorganic gases in the exhaust gases). Jap J Hyg 22: 341-353, 1967. A 306
- SILVER H M : Carbon monoxide in health and politics. Med Ann DC 40: 169-171, 1971. A 307
- SPINAZZOLA A, MARRACCINI L, DEVOTO G and ZEDDA S : Studio sul comportamento di alcuni prodotti tossici volatili quale indice di inquinamento atmosferico nella città di Cagliari. (Study of the behavior of some volatile toxic products as indices of air pollution in the city of Cagliari. V. Carbon monoxide). Folia Med (Napoli) 49: 659-668, 1966. A 308
- TROMPEO G, TURLETTI G and GIARRUSSO O T : Concentrazioni di CO nelle autorimesse interrato. (Concentrations of CO in underground garages). Rass Med Industr 33: 392-393, 1964. A 309
- UBISCH H and WESTERLUND K : The concentration of carbon monoxide in a city building and in underground garages. Nord Hyg Tidskr 52: 48-51, 1971. A 314
- UMEZAWA T : (Air pollution and blood gases). Naika 21: 836-840, 1968, (Jap.). A 310
- VEIL C : L'attitude épidémiologique. (The epidemiologic attitude). Evol Psychiatr (Paris) 33: 253-262, 1968. A 311
- VOORHOEVE R J H, REMEIK A J P, FREELAND P E and MATTHIAS B T : Rare-earth oxides of manganese and cobalt rival platinum for the treatment of carbon monoxide in auto exhaust. Science 177: 353-354, 1972. A 312
- WEAVER N K : Toxicologic implications of motor gasoline and auto emissions. Ind Med Surg 40: 31-34, 1971. A 313
- WIETHAUP H : Zum anteil der autoabgase an der luftverunreinigung. (On the role of automobile exhausts in air pollution). Med Klin 63: 1088-1089, 1968. (Ger) A 315
- WOODRUFF P S : Environmental pollution-nuisance, menace or disaster? Med J Aust 1: 1162-1167, 1970. A 316
- YAMATE N and MATSUMURA T : (Summary of carbon monoxide levels during the past six years (1964-1969) at three air monitoring stations in Tokyo). Bull Natl Inst Hyg Sci (Tokyo) 89: 107-111, 1971. A 317

1005051260

## Additional Bibliography List No. 9

## ANALYSIS OF CARBON MONOXIDE IN THE AIR

- |  | Reprint |
|--|---------|
| AJEMIAN R S and WHITMAN N E : Monitoring carbon monoxide in ambient air. <u>J Air Pollut Contr Ass</u> 20: 310-1, 1970.  | A 318   |
| ANON : Tentative method of analysis for methane and carbon monoxide content of the atmosphere (gas chromatographic method by reduction to methane). <u>Health Lab Sci</u> 9: 58-61, 1972.  | A 319   |
| AUBEAU R, LEROY J and CHAMPEIX L : Influence du degré d'hydratation de l'adsorbant sur l'analyse chromatographique des gaz permanents. (Influence of degree of hydration of the adsorbent on the chromatographic analysis of permanent gases.) <u>J Chromatogr</u> 19: 249-62, 1965. | A 320   |
| BARRETT J C, BENNETT R and BUCKMASTER J : Automatic carbon monoxide monitor. <u>Am Industr Hyg Ass J</u> 27: 402-6, 1968.  | A 321   |
| BLANC C, HUYNH C T and ESPAGNO L : Enrichissement des isotopes du carbone et du néon par chromatographie en phase gazeuse. 1ère partie. (Enrichment of carbon and neon isotopes by gas phase chromatography.) <u>J Chromatogr</u> 28: 177-93, 1967.                                  | A 322   |
| BOTTEAU H L and MOUSSION C : Un appareil simple pour le dosage de quelques toxiques gazeux ou volatils. (A simple apparatus for the determination of various gaseous or volatile poisons.) <u>Ann Biol Clin(Paris)</u> 25: 215-27, 1967.   | A 323   |
| CELEGIN M, HANSSON R and SUNDSTROM G : A sub-microliter sampling device for quantitative collection of gases. Application to 'gas chromatographic' analysis. <u>Scand Clin Lab Invest</u> 27: 367-70, 1971.  | A 324   |
| CIUHANDU G, DIACONOVICI M, KISS L and RUSU V : Bestimmung des ausgeatmeten Kohlenmonoxyds. (Determination of exhaled carbon monoxide.) <u>Hoppe Seyler Z Physiol Chem</u> 339: 104-9, 1964.  | A 325   |
| CIUHANDU G and RUSU V : Photometrische Mikrobestimmung von Kohlenmonoxid in Luft und Blut. (Photometric microdetermination of carbon monoxide in air and blood.) <u>Z Klin Chem Biochem</u> 6: 204-8, 1968.  | A 326   |
| CLARK J C and BUCKINGHAM P D : The preparation and storage of carbon-11 labelled gases for clinical use. <u>Int J Appl Radiat Isot</u> 22: 639-46, 1971.   | A 327   |
| DAVIES G M, JONES J G and WARNER C G : A continuously recording atmospheric carbon monoxide monitoring system with fully automatic alarms in a blast furnace area. <u>Brit J Industr Med</u> 22: 270-8, 1965.  | A 328   |
| DUBOIS L, ZDROJEWSKI A and MONKMAN J L : The analysis of carbon monoxide in urban air at the ppm level, and the normal carbon monoxide value. <u>J Air Pollut Contr Ass</u> 16: 135-9, 1966.   | A 329   |
| DUBOIS L and MONKMAN J L : Continuous determination of carbon monoxide by frontal analysis. <u>Anal Chem</u> 44: 74-6, 1972.   | A 330   |
| DUCROS H : Epuration de l'oxyde de carbone par l'hopcalite dans une enceinte close. (Purification of carbon monoxide by hopcalite in a closed environment.) <u>Rev Corps Sante Armées</u> 9: 645-64, 1968.   | A 331   |
| FELDSTEIN M : The colorimetric determination of blood and breath carbon monoxide. <u>J Forensic Sci</u> 10: 35-42, 1965.   | A 332   |

1005051261

## Additional Bibliography List No. 9

- FELDSTEIN M: Methods for the determination of carbon monoxide. Prog Chem Toxic 3: 99-119, 1967. A 333
- GORODINSKY S M, LEVINSKY S V and SCHERBAKOV V L: (Determination of standard values of noxious admixtures in the use of insulating respiratory apparatus.) Gig Sanit 32: 42-7, 1967. A 334
- IABLOCHKIN V D: (A modified method of photoelectrocolorimetric determination of carbon monoxide in the decomposition products of several polymers.) Gig Sanit 31: 59-62, 1966. A 335
- KIM T S and RYO U Y: A modified chemical analysis for carbon monoxide in respiratory gases. J Lab Clin Med 67: 873-8, 1966. A 336
- KUNZ C, DONDES S and HARTECK P: A spectroscopic study of the alpha-ray-induced luminescence in gases; the helium-carbon monoxide system. Radiat Res 41: 288-98, 1970. A 337
- LINCH A L and PFAFF H V: Carbon monoxide - evaluation of exposure potential by personnel monitor surveys. Am Industr Hyg Ass J 32: 745-52, 1971. A 338
- MCDOWELL R S: Metal carbonyl vapors: rapid quantitative analysis by infrared spectrophotometry. Am Industr Hyg Ass J 32: 621-4, 1971. A 339
- McFEE D R, LAVINE R E and SULLIVAN R J: Carbon monoxide, a prevalent hazard indicated by detector tabs. Am Industr Hyg Ass J 31: 749-53, 1970. A 340
- MOKHOV L A and CHUIKOVA N J: (On assessing the results of determining carbon monoxide in the air.) Lab Delo 4: 225-7, 1967. A 341
- MORGANSTERN A S, ASH R M and LYNCH J R: The evaluation of gas detector tube systems: I. Carbon monoxide. Am Industr Hyg Ass J 31: 630-2, 1970. A 342
- PIERCE J O and COLLINS R J: Calibration of an infrared analyzer for continuous measurement of carbon monoxide. Am Industr Hyg Ass J 32: 457-62, 1971. A 343
- RAMSEY J M: Potassium pallado sulfite detection of carbon monoxide in exhaled air as an estimate of carboxyhemoglobin. Am Industr Hyg Ass J 28: 531-4, 1967. A 344
- ROBBINS R C, BORG K M and ROBINSON E: Carbon monoxide in the atmosphere. J Air Pollut Contr Ass 18: 106-10, 1968. A 345
- RODKEY F L: Discussion: Carbon monoxide estimation in gases and blood by gas chromatography. Ann NY Acad Sci 174: 261-7, 1970. A 346
- SANZ RAMOS PEDRERO P and RODRIGO P: La toxicología del óxido de carbono con las técnicas para su determinación. Un nuevo método polarográfico para la evaluación en sangre y en el ambiente. (The toxicology of carbon monoxide with the new technics for its determination. A new polarographic method for its evaluation in the blood and in the air.) An Acad Farm(Madrid) 30: 228-59, 1964. A 347
- SEMAR M, TRESER G and LANGE K: Quantitative comparative immunohistology. Clin Chim Acta 15: 505-15, 1967. A 348
- SILVERMAN L and GARDNER G R: Potassium pallado sulfite method for carbon monoxide detection. Am Industr Hyg Ass J 26: 97-105, 1965. A 349
- SLATER K: Notes on experimental technique and apparatus. The detection and measurement of dangerous quantities of carbon monoxide gas. J Sci Instrum 44: 642-3, 1967. A 350
- SMITH R G, BRYAN R J, FELDSTEIN M, LEVADIE B, MILLER F A, STEPHENS E R and WHITE N G: Tentative method of continuous analysis for carbon monoxide content of the atmosphere (nondispersive infrared method). Health Lab Sci 7: Suppl:81-6, 1970. A 351

1005051262

## Additional Bibliograph List No. 9

- SMITH R G, BRYAN R J, FELDSTEIN M, LEVADIE B, MILLER F A, STEPHENS E R and WHITE N G: Tentative method for preparation of carbon monoxide standard mixtures. Health Lab Sci 7: Suppl: 72-4, 1970. A 352
- SMITH R G, BRYAN R J, FELDSTEIN M, LEVADIE B, MILLER F A, STEPHENS E R and WHITE N G: Tentative method of analysis for carbon monoxide content of the atmosphere (manual- colorimetric method). Health Lab Sci 7: Suppl: 75-7, 1970. A 353
- SMITH R G, BRYAN R J, FELDSTEIN M, LEVADIE B, MILLER F A, STEPHENS E R and WHITE N G: Tentative method of analysis for carbon monoxide content of the atmosphere (infrared absorption method). Health Lab Sci 7: Suppl: 78-80, 1970. A 354
- SMITH R G, BRYAN R J, FELDSTEIN M, LEVADIE B, MILLER F A, STEPHENS E R and WHITE N G: Tentative method of analysis for carbon monoxide content of the atmosphere (hopcalite method). Health Lab Sci 9: 84-7, 1972. A 355
- SMITH T C: Carbon monoxide, neon, and acetylene analysis in the presence of anesthetic gases. J Appl Physiol 21: 745-6, 1966. A 356
- STAMM E: Eine photometrische Methode zur Bestimmung kleiner CO-Mengen in der Luft. (A photometric method for the determination of small CO quantities in the air.) Z Ges Hyg 13: 160-3, 1967. A 357
- VANNESTE W H: A radiochemical method for determination of carbon monoxide in chemical combination. Anal Biochem 15: 65-76, 1966. A 358
- WEASE D F : An automated, closed system for gas analysis. Assay of carbon monoxide in blood. US Air Force Sch Aerospace Med 1-12, 1967. A 359
- XINTARAS C, JOHNSON B L, ULRICH C E, TERRILL R E and SOBECKI M F: Application of the evoked response technique in air pollution toxicology. Toxicol Appl Pharmacol 8: 77-87, 1966. A 360

1005051263



## CARBON MONOXIDE USE FOR MEASURING PULMONARY DIFFUSION CAPACITY

- AYRES S M, BUEHLER M E and ARMSTRONG R G : Diffusing capacity of the lung in pulmonary emphysema. J Appl Physiol 19: 981-9, 1964. A 361
- BATES D V, CHRISTIE R V and VARVIS C J : The measurement of the pulmonary capillary blood volume and membrane diffusion component on exercise. J Physiol London 154: 13P-14P, 1960. A 362
- BEDELL G N and OSTIGUY G L : Transfer factor for carbon monoxide in patients with airways obstruction. Clin Sci 32: 239-48, 1967. A 363
- BEEREL F R and VANCE J W : Daily  $P_{CO_2}$  and pH fluctuations in pulmonary emphysema with carbon dioxide retention. Am Rev Resp Dis 92: 894-9, 1965. A 364
- BEUMER H M : Determination of the transfer factor and half time of carbon monoxide during apnea in the human lungs in normal and pathological cases. Med Thorac 21: 204-22, 1964. A 365
- BEUMER H M : De Bepaling van de diffusiecapaciteit van de longen met Behulp van Koolmonoxyde. (The determination of the diffusion capacity of the lungs with the aid of carbon monoxide.) Nederl Milit Geneesk T 18: 89-102, 1965. A 366
- BJURE J : Pulmonary diffusing capacity for carbon monoxide in relation to cardiac output in man. Scand J Clin Lab Invest 17: Suppl 81: 1-113, 1965. A 367
- BOLLINELLI R, ROUCH Y, PUJOL M, CARRIERE R and CARLES P : Interet du test au CO en etat stable, dans le depistage de la bronchite chronique, en medecine du travail. (Importance of the steady-state CO diffusion test in the detection of chronic bronchitis in industrial medicine.) Med Lav 62: 101-10, 1971. A 368
- BOOZ J : Fréquence ventilatoire et "capacité de diffusion" de l'oxyde de carbone mesurée en "steady state". (Ventilatory and diffusion capacity of carbon monoxide measured at steady state.) Arch Int Physiol 77: 939-43, 1969. A 369
- BOUHUYS A, GEORG J, JÖNSSON R, LUNDIN G and LINDELL S E : The influence of histamine inhalation on the pulmonary diffusing capacity in man. J Physiol 152: 176-81, 1960. A 370
- BURGESS J H, GILLESPIE J, GRAF P D and NADEL J A : Effect of pulmonary vascular pressures on single-breath CO diffusing capacity in dogs. J Appl Physiol 24: 692-6, 1968. A 371
- CANEPA F, CAVALLO F and MUZIO M : Aspetti tecnici di funzione respiratoria: descrizione di un apparecchio per la determinazione successiva della meccanica respiratoria della diffusione polmonare del CO. (Technical aspects of respiratory function: description of an apparatus for successive determination of respiratory mechanics of pulmonary diffusion of CO.) Arch Maragliano Pat Clin 24: 359-63, 1968. A 372
- CASULA D, NISSARDI G P, SANNA-RANDACCIO F and FRAU P : Ricerche sugli scambi respiratori nella silicosi polmonare. III. Studio delle correlazioni esistenti fra  $PaO_2$  e "trasfert" del CO. (Research on respiratory exchanges in pulmonary silicosis. Study of the correlation between  $PaO_2$  and "transfer" of CO.) Boll Soc Ital Biol Sper 45: 1290-2, 1969. A 373
- CASULA D, NISSARDI G P, SANNA-RANDACCIO F and FRAU P : Ricerche sugli scambi respiratori nella silicosi polmonare. IV. Studio delle correlazioni esistenti fra  $SaO_2$  "transfert" del CO. (Research on respiratory exchanges in pulmonary silicosis. IV. Study of the correlations between  $SaO_2$  and "transfer" of CO.) Boll Soc Ital Biol Sper 45: 1292-3, 1969. A 374

1005051264

- CHINET A, MICHELI J L and HAAB P : Inhomogeneity effects on  $O_2$  and CO pulmonary diffusing capacity estimates by steady-state methods. *Theory. Res Physiol* 13: 1-22, 1971. A 375
- CINKOTAI F F and THOMSON M L : Diurnal variation in pulmonary diffusing capacity for carbon monoxide. *J Appl Physiol* 21: 539-42, 1966. A 376
- CIUHANDU G, DIACONOVICI M, KISS L and RUSU V : Die Beziehung zwischen Kohlenmonoxidausscheidung in der Ausatemluft und beruflicher Exposition. (Relationship between carbon monoxide exhalation in the breath and occupational exposure.) *Z Arbeitsmed* 18: 172-6, 1968. A 377
- COTES J E, DABBS J M, EVANS M R and HOLLAND P : Effect of CS aerosol upon lung gas transfer and alveolar volume in healthy men. *Q J Exp Physiol* 57: 199-206, 1972. A 378
- DALY W J and WALDHAUSEN J A : Physiologic studies of the pulmonary capillary bed after barium sulfate embolization. *J Clin Invest* 46: 1617-24, 1967. A 379
- DALY W J : Pulmonary diffusing capacity for carbon monoxide and topography of perfusion during changes in alveolar pressure in man. *Am Rev Resp Dis* 99: 548-53, 1969. A 380
- DALLE M, TOURNAIRE C, BRUDIEUX R and DELOST P : Évolution néo-natale des corticostéroïdes surrénaliens et plasmatiques chez le Cobaye. (Comparative study of respiratory dead space simultaneously measured with CO and  $CO_2$ .) *J Physiol (Paris)* 63: 34A, 1971. A 381
- DeGRAFFA A C Jr, TAYLOR H F, ORD J W, CHUANG T H and JOHNSON R L Jr : Exercise limitation following extensive pulmonary resection. *J Clin Invest* 44: 1514-22, 1965. A 382
- FLETCHER E C : A nomogram for the transfer factor for carbon monoxide in the lungs. *Thorax* 27: 382-5, 1972. A 383
- FORBES W H, SARGENT F and ROUGHTON F J W : The rate of carbon monoxide uptake by normal men. *Am J Physiol* 143: 594-608, 1945. A 384
- FORBES W H : Carbon monoxide uptake via the lungs. *Ann NY Acad Sci* 174: 72-5, 1970. A 385
- FORSTER R E, ROUGHTON F J W, CANDLER L, BRISCOE W A and KREUZER F : Apparent pulmonary diffusing capacity for CO at varying alveolar  $O_2$  tensions. *J Appl Physiol* 11: 277-89, 1957. A 386
- FREYSCHUSS U and HOLMGREN A : On the variation of  $D_{LCO}$  with increasing oxygen uptake during exercise in healthy ordinarily untrained young men and women. *Acta Physiol Scand* 65: 193-206, 1965. A 387
- GABRIEL S : On the prediction of pulmonary diffusing capacity with a steady state carbon monoxide method in chronic lung diseases. *Scand J Clin Lab Invest* 24: 377-81, 1969. A 388
- GATTO E, CANEPA F, CAVALLO F and MASSIMILIA A : La diffusione polmonare del monossido di carbonio in soggetti normali. (Pulmonary diffusion of carbon monoxide in the normal subject.) *Arch Maragliano Pat Clin* 23: 699-706, 1967. A 389
- GERHARDT T, GOTHERT M, MALORNY G and WILKE H : Aufnahme von Kohlenoxid in den Organismus des Menschen bei Atmung von CO-Luftgemischen unter erhöhtem Druck. (Carbon monoxide uptake in human beings breathing CO-air mixtures under increased pressure.) *N Schmied Arch Pharmacol* 266: 332-3, 1970. A 390
- GULERIA J S, PANDE J N, SEITHI P K and ROY S B : Pulmonary diffusing capacity at high altitude. *J Appl Physiol* 31: 536-43, 1971. A 391
- GULZOW M : Die Kohlenoxydgasvergiftung. (Poisoning by carbon monoxide gas.) *Z Aerztl Fortbild* 51: 838-41, 1957. A 392

1005051265

- GUYATT A R, NEWMAN F, CINKOTAL F F, PALMER J I and THOMSON M L: Pulmonary diffusing capacity in man during immersion in water. J Appl Physiol 20: 878-81, 1965. A 393
- HAAB P, ROBERT M and FIPER J: Comparaison de la capacité de diffusion pulmonaire du Chien narcotisé mesurée simultanément par l'oxygène et le monoxyde de carbone. (Comparison of the pulmonary diffusion capacity of anesthetized dogs measured simultaneously by oxygen and carbon monoxide.) J Physiol Paris 60 Suppl 2: 455-6, 1968. A 394
- HAMILTON L H and KERSTING D J: A study of gas analysis for measurement of pulmonary diffusing capacity for carbon monoxide by chromatographic techniques. Am Rev Resp Dis 102: 916-20, 1970. A 395
- HAMM J: Diffusionskapazität. (Diffusion capacity.) Beitr Klin Erforsch Tuberk 133: 292-304, 1966. A 396
- HATZFELD C, WIENER F and BRISCOE W A: Effects of uneven ventilation - diffusion ratios on pulmonary diffusing capacity in disease. J Appl Physiol 23: 1 - 10, 1967. A 397
- HILPERT P: Die Änderung der Diffusionskapazität der Lunge für CO durch die Hämoglobinkonzentration des Blutes. (Variation of the carbon monoxide diffusing capacity of the lung with the hemoglobin concentration of the blood.) Respiration 28: 518-25, 1971. A 398
- HOLMGREN A: On the reproducibility of steady state  $D_{LCO}$  measurements during exercise in man. Scand J Clin Lab Invest 17: 110-6, 1965. A 399
- HOLMGREN A: On the significance of pulmonary mean capillary CO back pressure corrections for repeated measurements of  $D_{LCO}$  during exercise in man. Scand J Clin Lab Invest 17: 117-22, 1965. A 400
- HOLMGREN A: On the significance of validity and precision in the determination of alveolar carbon dioxide tension for measurements of alveolar ventilation and  $D_{LCO}$ . Scand J Clin Lab Invest 17: 123-9, 1965. A 401
- HOLMGREN A: On the variation of  $D_{LCO}$  with increasing oxygen uptake during exercise in healthy trained young men and women. Acta Physiol Scand 65: 207-20, 1965. A 402
- HSIEH Y C, ROSS J C, SMALL G R and THOMPSON E C: Effect of limb vascular occlusion on pulmonary diffusing capacity ( $D_{LCO}$ ) during rest and leg exercise. Am J Med Sci 256: 9-17, 1968. A 403
- HYDE R W, MARIN M G, RYNES R I, KARREMAN G and FORSTER R E: Measurement of uneven distribution of pulmonary blood flow to CO diffusing capacity. J Appl Physiol 31: 605-12, 1971. A 404
- JAMES F and RUMBLE L Jr: Carbon monoxide diffusing studies in the clinical evaluation of chronic lung diseases. Dis Chest 52: 387-91, 1967. A 405
- JOHNSON R L Jr, TAYLOR H F and LAWSON W H Jr: Maximal diffusing capacity of the lung for carbon monoxide. J Clin Invest 44: 349-55, 1965. A 406
- JOHNSON R L Jr, TAYLOR H F and DeGRAFF A C Jr: Functional significance of a low pulmonary diffusing capacity for carbon monoxide. J Clin Invest 44: 789-800, 1965. A 407
- JOHNSON R L Jr and MILLER J M: Distribution of ventilation, blood flow, and gas transfer coefficients in the lung. J Appl Physiol 25: 1-15, 1968. A 408
- KANAZIRSKY P, TSANEV B and GEORGIEV G: (Diffusion capacity of lungs for carbon monoxide.) Surv Med(Sofia) 16: 614-9, 1965. A 409
- KAWAMOTO T: (A study of CO diffusing capacity with reference to pulmonary hemodynamics and ventilatory functions in patients with cardiac or chronic pulmonary disease.) Jap Circ J 30: 251-65, 1966. A 410

1005051266

- KOTTER D, HUCH A, STOTZ H and PIIPER J : Single breath CO diffusing capacity in anesthetized dogs with increased oxygen consumption. Resp Physiol 6: 202-8, 1969. A 411
- KRAL B, CERNOCHOVA Z and TUSL M : Diffusing capacity of the lungs for CO and its components in healthy men of different age at rest and in physical load. Sborn Ved Prac Lek Fak Karlov Univ 9: 625-9, 1966. A 412
- KREUKNIET J and VISSER B F : The pulmonary CO diffusing capacity according to Bates and according to Filley in patients with unequal ventilation. Pflueg Arch Ges Physiol 281: 207-11, 1964. A 413
- KREUZER F and CAMPAGNE P L : Resting pulmonary diffusing capacity for CO and O<sub>2</sub> at high altitude. J Arr. Physiol 20: 519-22, 1965. A 413a
- LACOSTE J and ROUCH Y : Mesures simultanées chez l'Homme de l'efficacité des échanges pulmonaires pour le CO<sub>2</sub> et le CO avec addition d'un espace mort. (Simultaneous measurements in humans of the efficiency of pulmonary exchanges for CO<sub>2</sub> and CO with addition of a dead space.) C R Soc Biol(Paris) 160: 1667-70, 1966. A 414
- LACOSTE J : Exploration fonctionnelle respiratoire. La ductance DuCO (monoxyde de carbone): évaluation globale, non sanglante, de l'échangeur pulmonaire. (Functional respiratory investigations. Ductance of CO (carbon monoxide): global estimation of pulmonary exchanger, omitting blood tests.) Presse Med 79: 1781-4, 1971. A 415
- LAWSON W H Jr : Rebreathing measurements of pulmonary diffusing capacity for CO during exercise. J Appl Physiol 29: 896-900, 1970. A 416
- LAWSON W H Jr : Effect of drugs, hypoxia, and ventilatory maneuvers on lung diffusion for CO in man. J Appl Physiol 32: 788-94, 1972. A 417
- LEWIS C M and BRINK A J : Carbon monoxide diffusion of lungs in assessment of pulmonary blood flow in patients with intracardiac shunts. Brit Heart J 28: 359-65, 1966. A 418
- LO COCO A : Sul comportamento della capacità di diffusione per il monossido di carbonio in due gruppi di soggetti, affetti da tubercolosi polmonare e da broncopatia cronica ostruttiva. (Behavior of diffusion capacity for carbon monoxide in 2 groups of subjects with pulmonary tuberculosis and chronic obstructive bronchopathy.) G Ital Mal Torace 24: 27-30, 1970. A 419
- LOPEZ-MAJANO V : Reproducibility of the carbon monoxide diffusion capacity method. Respiration 28: 114-9, 1971. A 420
- MAUDERLY J L : Steady-state carbon monoxide-diffusing capacity of unanesthetized Beagle dogs. Am J-Vet Res 33: 1485-91, 1972. A 421
- MÄHRLEIN W, KRAUSE M and MÜLLER H R : Apnoetechnik zur Bestimmung der pulmonalen Diffusionskapazität für Kohlenmonoxid. (Apnea technic for the determination of the pulmonary diffusion capacity of carbon monoxide.) Z Ges Med 22: 373-5, 1967. A 422
- MENKES H A, SERA K, ROGERS R M, HYDE R W, FORSTER R E II and DuBOIS A B : Pulsatile uptake of CO in the human lung. J Clin Invest 49: 335-45, 1970. A 423
- MITCHELL M M and RENZETTI A D Jr : Application of the single-breath method of total lung capacity measurement to the calculation of the carbon monoxide diffusing capacity. Am Rev Resp Dis 97: 581-4, 1968. A 424
- MITTMAN C : Nonuniform pulmonary diffusing capacity measured by sequential CO uptake and washout. J Appl Physiol 23: 131-8, 1967. A 425
- NAIRN J R, POWER G G, HYDE R W, FORSTER R E, LAMBERTSEN C J and DICKSON J : Diffusing capacity and pulmonary capillary blood flow at hyperbaric pressures. J Clin Invest 44: 1591-9, 1965. A 426

1005051267

- NISSARDI G P, SANNA-RANDACCIO F, TORRAZZA P L and CASCIU G: Ricerche sulla capacità di diffusione alveolo-capillare per il CO (DCO). I. Comportamento della D<sub>CO</sub> nei silicotici in condizioni di riposo. (Research on the alveolo-capillary diffusion capacity for CO (D CO). I. Behavior of D CO in silicotics at rest.) Boll Soc Ital Biol Sper 41: 866-70, 1965. A 427
- NISSARDI G P, SANNA-RANDACCIO F, TORRAZZA P L and GARIEL G: Ricerche sulla capacità di diffusione alveolo-capillare per il CO (D CO). II. Modificazioni della D CO nei silicotici nel corso del lavoro muscolare. (Research on the alveolo-capillary diffusion capacity for CO (D CO). II. Changes in D CO in silicotics during muscular work.) Boll Soc Ital Sper 41: 870-4, 1965. A 428
- NISSARDI G P, TORRAZZA P I and ANEDDA G: Ricerche sulla capacità di diffusione per il monossido di carbonio. Effetto del broncospasmo indotto mediante aerosol di acetilcolina. (Research on the diffusion capacity for carbon monoxide. Effect of bronchospasm induced with acetylcholine aerosol.) Boll Soc Ital Biol Sper 43: 130-4, 1967. A 429
- NISSARDI G P, SANNA-RANDACCIO F and SARNA R: Ricerche sulla capacità di diffusione nei silicotici rapporto tra D CO e alterazioni ventilatorie. (Research on diffusion capacity in silicotics. Relation between D CO and ventilatory changes.) Boll Soc Ital Biol Sper 44: 75-80, 1968. A 430
- PAEZ P N: Carbon monoxide levels in patients with chronic obstructive pulmonary disease. (COPD). Chest 58: 287, 1970. A 431
- PIPER J and SIKAND R S: Determination of D CO by the single breath method in inhomogeneous lungs: Theory. Resp Physiol 1: 75-87, 1966. A 432
- PIPER J, PFEIFER K and SCHEID P: Carbon monoxide diffusing capacity of the respiratory system in the domestic fowl. Resp Physiol 6: 309-17, 1969. A 433
- PIRNAY F, FASSOTTE A, GAZON J, DEROANNE R and PETIT J M: Diffusion pulmonaire au cours de l'exercice musculaire. Int Z Angew Physiol 28: 31-7, 1969. A 434
- PIRNAY F, PETIT J M and ROBERTS M: Evolution de la capacité de transfert du CO pendant le bronchospasme provoqué. (Development of the CO transfer capacity during induced bronchospasm.) Acta Tuberc Pneumol Belg 61: 95-9, 1970. A 435
- PODLESCH I and STEVANOVIC M: Die Altersabhängigkeit der Diffusionskapazität der Lunge in Ruhe und während Belastung. (Effect of age on the pulmonary diffusion capacity at rest and during exercise.) Med Thorac 23: 144-59, 1966. A 436
- POWER G G Jr, HYDE R W, SEVER R J, HOPPIN F G Jr and NAIRN J R: Pulmonary diffusing capacity and capillary blood flow during forward acceleration. J Appl Physiol 20: 1199-204, 1965. A 437
- POWER G G, AOKI V S, LAWSON W H Jr and GREGG J B: Diffusion characteristics of pulmonary blood-gas barrier at low temperatures. J Appl Physiol 31: 438-46, 1971. A 438
- RANDOWA D and SIERAWSKI S: Oznaczanie pojemności dyfuzyjnej płuc za pomocą tlenku węgla (metoda pojedynczego oddechu) w przewlekłych chorobach płuc. (Determination of the diffusion capacity of the lung with aid of carbon monoxide (single respiration method) in chronic lung diseases.) Pol Tyg Lek 19: 829-31, 1964. A 439
- RANDOWA D: Zaburzenia dyfuzji w przewlekłych chorobach płuc (metoda tlenkowęgłowa). (Disturbances in diffusion in chronic pulmonary diseases (carbon monoxide method).) Gruzlica 35: 465-72, 1967. A 440
- REMMERS J E and MITHOEFER J C: The carbon monoxide diffusing capacity in permanent residents at high altitudes. Resp Physiol 6: 233-44, 1969. A 441

1005051268

Additional Bibliography List No. 10

- SANNA-RANDACCIO F and NISSARDI G P : L'influenza della ventilazione sulla misura del 'trasferimento' del CO in stato stabile. (Influence of ventilation on the measurement of 'transfer' of CO in stable state.) Boll Soc Ital Biol Sper 45: 762-5, 1969. A 442
- SARTORELLI E : Alterazioni della diffusione alveolo-capillare del CO nella silicosi. (Changes in alveolo-capillary diffusion of CO in silicosis.) Acta Geront (Milano) 17: 182-6, 1967. A 443
- SATAKE T, IIDA S, USUI A, TATSUBANA K, YAMAZAKI J and MATSUOKA T : (Studies on the measurement of the carbon monoxide diffusing capacity by the single-breath method, with special reference to the influence of the uneven distribution of ventilation-perfusion of the lungs.) Jap Circ J 32: 35-40, 1968. A 444
- SHAW D B, CINKOTAI F and THOMSON M L : Syncope induced by application of negative pressure to the lower body and its effect on lung CO diffusing capacity. Aerospace Med 37: 154-7, 1966. A 445
- SIKAND R S and PIIPER J : Pulmonary diffusing capacity for CO in dogs by the single breath method. Resp Physiol 1: 172-92, 1966. A 446
- SIMONE M, REGGIANI A and BET E : La capacità di diffusione per il monossido di carbonio nella valutazione preoperatoria. (Carbon monoxide diffusion capacity in preoperative evaluation.) Acta Anaesth (Padova) 16: Suppl 4: 103-14, 1965. A 447
- STEINER S H, FRAYSER R and ROSS J C : Alterations in pulmonary diffusing capacity and pulmonary capillary blood volume with negative pressure breathing. J Clin Invest 44: 1623-30, 1965. A 448
- TARTULIER M, TOURNIAIRE A, DEYRIEUX F and BLUM J : Étude de transfert alvéolo-capillaire de l'oxyde de carbone dans la sténose mitrale. (Study of alveolo-capillary transfer of carbon monoxide in mitral stenosis.) Arch Mal Coeur 60: 843-64, 1967. A 449
- TLUSTÝ L, HLOUŠKOVÁ Z, KROFTA K and DAUM S : Difúzní kapacita plicni pro CO, vyšetřena metodou 'steady state' a metodou 'single breath'. (Pulmonary diffusing capacity for CO examined by 'steady state' and 'single breath' method.) Cas Lek Cesk 111: 149-53, 1972. A 450
- TRINQUET G, CLAUZEL A M, CARRÉ R and MEYER A : Les valeurs du transfert de l'oxyde de carbone chez l'homme normal en état ventilatoire stable au repos. (Values for carbon monoxide transfer in normal humans in stable ventilatory state at rest.) J Franc Med Chir Thorac 21: 365-82, 1967. A 451
- VANDENBERGH E, BILLIET L, WOESTIJNE K P and GYSELEN A : Relation between single-breath diffusing capacity and arterial blood gases in chronic obstructive lung disease. Scand J Resp Dis 49: 92-101, 1968. A 452
- VANROUX R and GREGOIRE M : La capacité de diffusion au CO chez le bronchiteux chronique. (The CO diffusion capacity in chronic bronchitis.) Acta Tuberc Belg 55: 488-97, 1964. A 453
- WAGNER P D, MAZZONE R W and WEST J B : Diffusing capacity and anatomic dead space for carbon monoxide ( $C^{18}O$ ). J Appl Physiol 31: 847-52, 1971. A 454
- WAGNER W W Jr, LATHAM L P, BRINKMAN P D and FILLEY G F : Pulmonary gas transport time: larynx to alveolus. Science 163: 1210-1, 1969. A 455
- WEBER J, MORET P and CHAUVET M : L'utilité du test de diffusion au monoxyde de carbone dans l'exploration fonctionnelle du poulmon. (Utility of the diffusion test with carbon monoxide in functional exploration of the lung.) Helv Med Acta 34: Suppl: 101-2, 1967. A 456
- WEISKOPF R B and SEVERINGHAUS J W : Diffusing capacity of the lung for CO in man during acute acclimation to 14,246 ft. J Appl Physiol 32: 285-9, 1972. A 457
- WOLKONSKY P M : Pulmonary effects of air pollution. Arch Environ Health 19: 586-92, 1969. A 458
- WOOLF C R : An assessment of the fractional carbon monoxide uptake and its relationship to pulmonary diffusing capacity. Dis Chest 46: 181-9, 1964. A 459

1005051269

## Additional Bibliography List No. 11

## LESIONS OF THE CENTRAL NERVOUS SYSTEM IN CARBON MONOXIDE POISONING

- AITKEN R C B, DALY R J, KREITMAN N, MATTHEW H and PROUDFOOT A T: Coal gas and the brain. Brit Med J 1: 706-7, 1968. A 460
- AMYOT R, GIARD N and ROBERT F: Anoxie cérébrale aigue. (Acute cerebral anoxia.) Un Med Canad 96: 680-8, 1967. A 461
- ANDO S: (Cerebral lesions caused by carbon monoxide poisoning.) Iryo 20: 638-48, 1966. A 462
- ANON: Carbon monoxide poisoning. W Virginia Med J 60: 344-5, 1964. A 463
- ANON: Neurological complications of carbon monoxide poisoning. Lancet 1: 77, 1968. A 464
- ANON: Coal gas and the brain. Brit Med J 1: 398, 1968. A 465
- ARAI H: Clinical and pathological study of akinetic mutism: 12 autopsy cases. Brain Nerve Tokyo 21: 615-32, 1969. A 466
- ARNOTT G, PETIT H and CHABRIER A: Coma oxycarbone, survenue apres important intervalle libre, d'un syndrome akineto-hypertonique avec trismus et mutisme, evolution favorable. (Carbon monoxide coma. Occurrence, after an important free interval, of an akineto-hypertonic syndrome with trismus and mutism. Favorable evolution.) Lille Med 9: 728-33, 1964. A 467
- ASAI K and TORU M: Changes of the brain focal symptoms in two cases of carbon monoxide intoxication (interval form). Psychiat Neurol Jap 71: 776-89, 1969. A 468
- BADAL J: Lazenske lecení diskogenni nemoci krční páteře. (Balneological treatment of diskogenic disease of the cervical spine.) Cesk Otolaryng 13: 354-62, 1964. A 469
- BEARON A: Carbon monoxide intoxication. Minnesota Med 48: 1537-43, 1965. A 470
- BERTONE E: Complicazioni neuropsichiche dovute ad avvelenamento subacuto di monossido di carbonio. (Neuropsychic complications due to subacute poisoning with carbon monoxide.) Minerva Med 56: 1657, 1965. A 471
- BONNET M H, GRATADOU M, BONNET H and LECIAK J P: État confuso-déirant post-intervallaire après coma oxycarbone. (Post-interval confusional and delirious state after carbon monoxide coma.) J Med Lyon 48: 1367-9, 1967. A 472
- BOUR H, GUY-GRAND B, TUTIN M and TAMINIAUX A M: Glycorégulation et coma oxycarbonné. (Glycoregulation and carbon monoxide coma.) J Ann Diabet Hotel Dieu 7: 365-76, 1967. A 473
- BOUR H, TUTIN M and PASQUIER P: The central nervous system and carbon monoxide poisoning. I. Clinical data with reference to 20 fatal cases. Progr Brain Res 24: 1-30, 1967. A 474
- GRAMOND W A: Organic psychosis. Brit Med J 4: 561-4, 1968. A 475
- D'AMORE V, GIORDANO P L and PENATIG: A proposito di intossicazione acuta da ossido di carbonio a rapida evoluzione demenziale. Considerazioni su due casi. (Apropos of acute poisoning by carbon monoxide with rapid evolution toward dementia. Consideration on 2 cases.) G Psychiat Neuropat 96: 1-16, 1968. A 476
- DIMIZIO M, FELICI F and BIETTI C: Su un caso di avvelenamento da ossido di carbonio: contributo clinico. (On a case of carbon monoxide poisoning: clinical case.) Riv Neurol 39: 425-34, 1969. A 477
- DOBOSZ J and LUCZYWEK E: W sprawie stosunku zaburzeń wyższych czynności nerwowych uwarunkowanych ogniskowymi uszkodzeniami mózgu do ogólnego obniżenia sprawności intelektualnej. (Relationship of higher nervous activity disorders due to focal brain lesions to the general decrease in intellectual capacity.) Neurol Neurochir Pol 5: 497-502, 1971. A 478

1005051270

- DROGICHINA E A and RYZHKOVA M N : (The clinical picture and diagnosis of diencephalic pathology in occupational poisonings.) Gig Tr Prof Zabol 11: 20-4, 1967. A 479
- FAURE J, VINCENT D, ESCHAPASSE P, LOISEAU P and CASTAING R : Confrontation des signes électrocliniques observés au cours d'intoxications par oxyde de carbone. (Comparison of electro-clinical signs observed in carbon monoxide poisoning.) Rev Neurol (Paris) 112: 287-92, 1965. A 480
- GARLAND H and PEARCE J : Neurological complications of carbon monoxide poisoning. Q J Med 36: 445-55, 1967. A 481
- GARREL S, PERRET J, PELLAT J and ARNOULD P : Syndrome neuro-psychiatrique d'altération frontale: complication post-intervallaire d'une intoxication oxycarbonée. (Neuro-psychiatric syndrome of frontal aspect: delayed complications of carbon monoxide poisoning.) Rev Neurol (Paris) 122: 445-7, 1970. A 482
- GARREL S, PERRET J, PELLAT J and ARNOULD P : Neuro-psychiatric syndrome following carbon monoxide poisoning. Electroenceph Clin Neurophysiol 29: 534, 1970. A 483
- GAULTIER M, FOURNIER E, GERVAIS P and BODIN F : Encéphalopathie pancréatique survenue au décours d'une intoxication oxycarbonée. Comparaison avec l'encéphalopathie post-intervallaire de l'intoxication oxycarbonée. (Pancreatic encephalopathy occurring after carbon monoxide poisoning. Comparison with the post-interval encephalopathy of carbon monoxide poisoning.) Presse Med 72: 3263-5, 1964. A 484
- GIRARDI G, CIS C and PIATTI A : La sindrome nucleo-reticolare cronica nelle intossicazioni professionali. (Chronic nucleo-reticular syndrome in occupational poisonings.) Arch Ital Otol 78: 756-70, 1967. A 485
- GORALSKI H and JANUSZKO L : Zespoły neurologiczne i psychiatryczne po zatruciu tlenkiem węgla. (Neurological and psychiatric syndromes after carbon monoxide poisoning.) Neurol Neurochir Pol 2:633-7, 1968. A 486
- GORDON E B : Carbon monoxide encephalopathy. Brit Med J 5444: 1343, 1965. A 487
- GROHME S, SCHNEIDER H and MASSHOFF W : Encephalopathien bei Vita reducta. (Encephalopathies in vita reducta.) Internist (Berlin) 10: 430-42, 1969. A 488
- GUNTHER K D : Lange verkannte CO-Vergiftungen mit schweren neurologischen Symptomen. (Chronically misdiagnosed CO poisoning with severe neurological symptoms.) Psychiat Neurol Med Psychol (Leipz) 23: 368-77, 1971. A 489
- HAMEL-PUSKARIC H, BERITIC T, JUSIC A and FRANJIC F : Neurološke komplikacije otrovanja ugljičnim monoksidom. (Neurologic complications of carbon monoxide poisoning.) Neuropsihijatrija 18: 147-55, 1970. A 490
- HANSEN D : Berufshedingte Riechstörung infolge chronischer Kohlenoxydeinwirkung. (Occupationally induced anosmia due to chronic carbon monoxide effect.) HNO 5: 140-2, 1970. A 491
- HARADA M and KOZUMA Z : (A case of carbon monoxide poisoning with so-called Sudeck's syndrome, and various neurologic symptoms.) Brain Nerve (Tokyo) 20: 1095-9, 1968. A 492
- HIRAI T : (The relationship between the disturbance of consciousness and behavior disorders.) Adv Neurol Sci (Tokyo) 14: 712-22, 1971. A 493
- IKUTA T : (Somatosensory evoked responses in patients with carbon monoxide poisoning as compared with those in Schizophrenics.) Fol Psychiatr Neurol Jap 23: 285-9, 1969. A 494

1005051271



## Additional Bibliography List No. 11

- INANAGA K : (Clinical problems of memory.) Adv Neurol Sci (Tokyo) 10: 648-60, 1966. A 497
- INANAGA K : Application of the averaged photopalpebral reflex in clinical neurology. Proc Aust Ass Neurol 5: 651-2, 1968. A 498
- ISHIKAWA H : (Effects of carbon monoxide on electrical activity of the brain: comparative studies on asphyxia, N<sub>2</sub>-inhalation and CO-inhalation.) J Kumamoto Med Soc 43: 870-84, 1969. A 499
- JESCHECK J : Laryngologische Untersuchungsergebnisse und ihre Beziehungen zur Neurologie. Msschr Ohrenheilk 101:378-94, 1967. A 500
- JORDI A : Schädigungen des Nervensystems durch gewerbliche Gifte, unter besonderer Berücksichtigung der Frühsymptome. (Damage to the nervous system by industrial poisons with special consideration of early symptoms.) Praxis 56: 610-8, 1967. A 501
- KATSUKI S : (Neuropsychic symptoms of patients in carbon monoxide poisoning caused by mining explosions.) Jap J Clin Med 23: 1928-36, 1965. A 502
- KEHL H and KEHL R : Morbus Basedow nach Kohlenmonoxydgasvergiftung. (Basedow's disease following carbon monoxide poisoning.) Msschr Unfallheilk 70: 349-55, 1967. A 503
- KHROLENKO D E : (Nervous system involvement in acute carbon monoxide poisoning.) Klin Med (Moskva) 47: 130-1, 1969. A 504
- KOLB K P : Lokale ischämische Kontrakturen der Hand nach Suizidversuchen. (Local ischemic contractures of the hand following suicide attempts.) Munch Med Wosenschr 110: 1873-4, 1968. A 505
- KRUG H : Die Hirnkonsistenz bei tödlicher Kohlenmonoxydvergiftung. (Brain consistency and fatal carbon monoxide poisoning.) Deutsch Z Gerichtl Med 56: 74-80, 1965. A 506
- KUROIWA Y, SHIDA K and KATO M : (Neurological aspect of acute carbon monoxide poisoning.) Adv Neurol Sci (Tokyo) 13: 4-10, 1969. A 507
- LANGAUER-LEWOWICKA H : (Early and remote neurological disorders due to acute carbon monoxide poisoning.) Med Pracy 17: 340-3, 1966. A 508
- LAPRESLE J and FARDEAU M : Les leuco-encéphalopathies de l'intoxication oxycarbonée. (The leukoencephalopathies caused by carbon monoxide poisoning. Study of sixteen anatomical-clinical observations.) Acta Neuropath (Berlin) 6: 327-48, 1966. A 509
- LAPRESLE J and FARDEAU M : The central nervous system and carbon monoxide poisoning. II. Anatomical study of brain lesions following intoxication with carbon monoxide (22 cases). Progr Brain Res 24: 31-74, 1967. A 510
- LAPRESLE J and FARDEAU M : Intoxication aigue par l'oxyde de carbone. Étude anatomique d'une encéphalopathie remarquable par la diffusion des lésions nécrotiques et l'importance des dépôts calciques. (Acute carbon monoxide poisoning. Anatomic study of an unusual encephalopathy caused by the diffusion of necrotic lesions and the importance of calcium deposits.) Bol Estud Med Biol 27: 9-17, 1971. A 511
- MACKINTOSH T F : Akinetic mutism following coal-gas poisoning with subsequent recovery. Postgrad Med J 41: 567-73, 1965. A 512
- MATTHEW H and PROUDFOOT A T : Coal gas and the brain. Brit Med J 1: 638, 1965. A 513
- MOORE M E and FINESTONE A J : The case of the disappearing headache. New Eng J Med 276: 1216, 1968. A 514
- MOROVIC-BUDAK A : Die Bedeutung des Befundes von punktförmigen Blutungen im Gebiet des Tegmen tympani. (Significance of punctiform hemorrhages in the area of the Tegmen

1005051272

## Additional Bibliography List No. 11

- MUROFUSHI K and MINAGAWA M: (An autopsy case of carbon monoxide poisoning with senile findings.) Adv Neurol Sci (Tokyo) 13:39-48, 1969. A 516
- OKUMA T, ISHINO H, SUNAMI Y and MOTOIKE M: An autopsy case of relapsing form carbon monoxide intoxication with special reference to the apallic syndrome and sleep cycle pattern. Fol Psychiat Neurol Jap 22: 43-51, 1968. A 517
- OSTROWSKA D: Morfologia odczynów tkankowych w encefalopatii tlenkoweglowej. (Morphology of tissue reactions in carbon monoxide encephalopathy.) Neurol Neurochir Pol 1: 561-8, 1967. A 518
- OSWALD E: Organische Demenz bei Intoxikationen. (Organic dementia in intoxications.) Wien Med Wochenschr 116-964-6, 1966. A 519
- PALADE D, MIHAI E, GOILAV N and SOVAREL G: Tulburarile neuropsihice la bolnavii cu oxicarbonism acut internati in sectia de boli interne a spitalului Piatra neamt intre anii 1964 si 1966. (Neuropsychological disorders in patients with acute carbon monoxide poisoning hospitalized in the department of internal medicine in the Piatra Neamt Hospital in the years 1964-1966.) Rev Medicochir Iasi 73: 1005-9, 1969. A 520
- PAULEIKHOFF B, MÜLLER-FAHLBUSCH MESTER H and MEIßNER U: Über spätfolgen, insbesondere Merkschwäche, nach Vergiftung mit Kohlenmonoxyd. (Late sequelae, especially weakness of memory, following carbon monoxide poisoning.) Fortschr Neurol Psychiatr 39: 349-77, 1971. A 521
- PEARCE J: Coal gas and the brain. Brit Med J 1: 767, 1968. A 522
- RINGEL S P and KLAUANS H L Jr: Carbon monoxide-induced Parkinsonism. J Neurol Sci 16: 245-51, 1972. A 523
- ROSENBLUTH P: Differential diagnosis of coma. Industr Med Surg 37: 108-12, 1968. A 524
- SANDERS H I and WARRINGTON E K: Memory for remote events in amnesic patients. Brain 94: 661-8, 1971. A 525
- SCHOTT B, TOMMASI M, BOURRAT C and MICHEL D: Neuropathie périphérique démyélinisante au cours d'une intoxication par l'oxyde de carbone. (Demyelinating peripheral neuropathy in the course of poisoning by carbon monoxide.) Rev Neurol (Paris) 116: 429-37, 1967. A 526
- SESSA T and SANNA G: La velocita' di conduzione nervosa nelle malattie professionali. (Nerve conduction rate in occupational diseases.) Fol Med (Napoli) 49: 809-15, 1966. A 527
- SHIDA K and KUROIWA Y: (Psychiatric signs, aphasia, apraxia and agnosia in acute carbon monoxide poisoning, with special reference to experiences in coal mine explosions.) Jap J Clin Med 27: 2333-7, 1969. A 528
- SHIMOJIMA W: (Classification of cerebral herniation.) Brain Ner (Tokyo) 22:577-86, 1970. A 529
- SHIRAKI H: (The neuropathology of carbon monoxide poisoning in humans - with special reference to the changes of globus pallidus.) Adv Neurol Sci (Tokyo) 13: 25-33, 1969. A 530
- SHIRUKI H and TATETSU M: (Toxic mental disorders.) Psychiat Neurol Jap 69: 994-1029, 1967. A 531
- SMITH J S, BRIERLEY H and BRANDON S: Akinetic mutism with recovery after repeated carbon monoxide poisoning. Psychol Med 1: 172-7, 1971. A 532
- SNYDER R D: Carbon monoxide intoxication with peripheral neuropathy. Neurology 20: 177-80, 1970. A 533
- TAKAMATSU I, TAKEICHI M and YUKITAKE A: (Light and electron microscopic observation on brains of experimentally induced CO poisoning cats.) Adv Neurol Sci (Tokyo) 13: 59-65, 1969.

1005051273

## Additional Bibliography List No. 11

- TOMASINI M : Un caso di encefalopatia demenziale postumo di intossicazione acuta da ossido di carbonio. (A case of demential encephalopathy after acute carbon monoxide poisoning.) Med Lav 58: 632-5, 1967. A 535
- TRILLET M, GIRARD P F and BOULETREAU P : Myelopathie oxycarbonée? (Carbon monoxide myelopathy?) Presse Med 78: 1843, 1970. A 536
- TRONZANO L and COSCIA G : Paralisi del radiale in un caso di intossicazione acuta da ossido di carbonio. (Paralysis of the radial nerve in a case of acute carbon monoxide poisoning.) Rass Med Industr 33: 401-3, 1964. A 537
- VITAL C and PICARD J : Neuropathologie de quelques intoxications aiguës. (Neuropathology of some acute poisonings.) J Med Bordeaux 144: 1029-38, 1967. A 538
- VUIA O : Leucoencéphalopathie souscorticale par intoxication au CO. (Subcortical leukoencephalopathy caused by CO poisoning.) Acta Neuropath(Berlin) 7: 305-14, 1967. A 539
- WENDER M : Studies of cerebral lipids in a relapsing case of carbon monoxide poisoning. Acta Neuropath(Berlin) 2: 371-7, 1963. A 540
- WITUSIK W : Erworbene chromatosthenopie und Chromatoanopie bei Kraftfahrern. (Acquired chromatosthenopia and chromatoanopia in automobile drivers.) Klin Monatsbl Augenheilkd 159: 689-92, 1971. A 541
- YASUKOCHI G and YASUOKA F : (Changes of personality as a sequela of acute carbon monoxide poisoning.) Psychiat Neurol Jap 69: 249-56, 1967. A 542
- YUKITAKE A : (Clinical investigation on acute carbon monoxide poisoning due to explosion in Miike coal mine 5 yrs. after the accident.) Psychiat Neurol Jap 72: 411-8, 1970. A 543

1005051274

## CARBON MONOXIDE ON HEPATIC CELLS

- ALVARES A P, SCHILLING G, LEVIN W and KUNTZMAN R: Studies on the induction of CO-binding pigments in liver microsomes by phenobarbital and 3-methyl cholanthrene. Biochem Biophys Res 29: 521-6, 1967. A 544 Reprint
- ALVARES A P, SCHILLING G, LEVIN W and KUNTZMAN R: Inability of substrates to alter the carbon monoxide and ethyl isocyanide difference spectra of microsomal hemoprotein. J Pharmacol Exp Ther 176: 1-10, 1971. A 545
- BELYAEV V A: (Concerning carbon monoxide action upon the glycogen-forming function of the liver.) Farmakol Toksik 30: 234-6, 1967. A 546
- BHATNAGAR S P: Release of cholinesterase from rat liver by nicotinamide and carbon tetrachloride. Biochem Pharmacol 19: 2009-16, 1970. A 547
- BOGDAN D P and JUCHAU M R: Characteristics of induced benzpyrene hydroxylase activity in the rat foeto-placental unit. Europ J Pharmacol 10: 119-26, 1970. A 548
- BRAUSER B, VERSMOLD H and BÜCHER T: Mechanisms of mixed function oxygenation. Redox kinetics and redox state of P-450 in whole liver. Hoppe-Seyler's Z Physiol Chem 349: 1589-90, 1968. A 549
- CHANCE B, ERECINSKA M and WAGNER M: Mitochondrial responses to carbon monoxide toxicity. Ann NY Acad Sci 174: 193-204, 1970. A 550
- CONNERY A H, LEVIN W, IKEDA M and KUNTZMAN R: Inhibitory effect of carbon monoxide on the hydroxylation of testosterone by rat liver microsomes. J Biol Chem 243: 3912-15, 1968. A 551
- COOPER D Y, SCHLEYER H and ROSENTHAL O: Some chemical properties of cytochrome P-450 and its carbon monoxide compound (P-450.CO). Ann NY Acad Sci 174: 205-17, 1970. A 552
- DATSENKO I I, DOTSENKO N S, MARTYNIUK V Z and PALCHEVSKY E I: (Characteristics of pathomorphological changes in the organism in carbon monoxide intoxication.) Vrach Delo 6: 77-80, 1965. A 553
- DATSENKO I I, DOTSENKO N S, MARTYNIUK V Z and PALCHEVSKY E K: (Histochemistry of polysaccharides and nucleoproteins in carbon monoxide poisoning.) Vrach Delo 1: 38-9, 1967. A 554
- ESTABROOK R W, FRANKLIN M R and HILDEBRANDT A G: Factors influencing the inhibitory effect of carbon monoxide on cytochrome P-450-catalyzed mixed function oxidation reactions. Ann NY Acad Sci 174: 218-32, 1970. A 555
- HERNANDEZ P H, MAZEL P and GILLETTE J R: Studies on the mechanism of action of mammalian hepatic azoreductase. II. The effects of phenobarbital and 3-methylcholanthrene on carbon monoxide sensitive and insensitive azoreductase activities. Biochem Pharmacol 16: 1877-88, 1967. A 556
- HLAVICA P, KIESE M, LANGE G and MOR G: Die Wirkung von Kohlenmonoxid auf die N-Hydroxylierung von Anilin durch Kaninchenlebermikrosomen. (Effect of carbon monoxide on the N-hydroxylation of aniline by rabbit liver microsomes.) Naunyn-Schmied Arch Pharm 263: 269-70, 1969. A 557
- HOCHSTRATE C and OBERDISSE E: Biochemische Veränderung im Serum der Ratte nach kombinierter Applikation lebervergrößernder und lebertoxischer Pharmaka. (Biochemical change in rat serum following combined application of inducing and hepatotoxic drugs. Naunyn-Schmied Arch Pharmacol 266: 357-8, 1970. A 558
- ICHIKAWA Y, HAGIHARA B and YAMANO T: Magnetic and spectrophotometric properties of the microsomal carbon monoxide binding pigment. Arch Biochem Biophys 120: 204-13, 1967. A 559

1005051275

- JEDRYCHOWSKI W, KUŚ J, PIOTROWSKI J and SAWICKI B : Zachowanie się aktywności oksydazy cytochromowej i peroksydazy w wątrobie szczura w doświadczalnym ostrym zatruciu tlenkiem węgla. (Cytochrome oxidase and peroxidase activities in the livers of rats in experimental carbon monoxide poisoning.) FoL Med Cracov 7: 429-35, 1965. A 560
- JEDRYCHOWSKI W, KUŚ J, PIOTROWSKI J and SAWICKI B : Zawartość glikogenu w wątrobie przy zatruciu tlenkiem węgla. (Glycogen content of the livers of rats poisoned with carbon monoxide.) FoL Med Cracov 7: 437-42, 1965. A 561
- KAMATAKI T and KITAGAWA H : (Studies of the binding of carbon monoxide with liver microsomal P-450 reduced by NADPH). J Pharm Soc Jap 91: 422-3, 1971. A 562
- KAMPFFMEYER H and KIESE M : The effect of carbon monoxide on the hydroxylation of aniline and N-Ethylaniline by microsomal enzymes. Naunyn Schmied Arch. Pharmacol 250: 1-8, 1965. A 563
- KRATZ F : Zur Frage der Entgiftung von Fremdstoffen durch die Leber. (On the problem of foreign substances detoxification through the liver.) Dtsch Med Wochensh 93: 2495-8, 1968. A 564
- KROBER F, LANGE G, MATHES S and MOR G : Änderung der Affinität isolierter Leber-mikrosomen für Anilin, N-Äthylanilin, Sauerstoff, Äthylisocyanid und Kohlenmonoxid durch die Behandlung junger Kaninchen mit Phenobarbital. (Change in the affinity of isolated liver microsomes for aniline, N-ethylaniline, oxygen, ethyl isocyanide and carbon monoxide by the treatment of young rabbits with phenobarbital.) Naunyn Schmied Arch Pharmacol 260: 161-3, 1968. A 565
- KROBER F, LANGE G, MATHES S and MOR G : Änderungen der spektralen Dissoziationskonstanten (Ks) von Anilin und N-Äthylanilin, der scheinbaren Michaelis-Konstanten (Km) für die Hydroxylierung dieser Substrate und der Affinität der beteiligten Enzyme für Sauerstoff und Kohlenmonoxid. (Qualitative changes in liver microsomes of phenobarbital-treated rabbits. Changes in spectral dissociation constants (Ks) of aniline and N-ethylaniline, in the apparent Michaelis constants (Km) for hydroxylation of these substrates, and in affinity of the enzymes involved for oxygen and carbon monoxide.) Naunyn Schmied Arch Pharmacol 267: 307-26, 1970. A 566
- KUNTZMAN R, LEVIN W, JACOBSON M and CONNEY A H : Studies on microsomal hydroxylation and the demonstration of a new carbon monoxide binding pigment in liver microsomes. Life Sci 7: 215-24, 1968. A 567
- LANGE P, KÄSTNER D and JUNG F : Die Beeinflussung der  $\text{CCl}_4$ -Hepatotoxizität und die Hemmung des mikrosomalen Stoffwechsels durch Diäthylthiokarbamat. (Influence of diethylthiocarbamate on the  $\text{CCl}_4$ -liver toxicity and the inhibition of the microsomal metabolism.) Acta Biol Med Ger 24: 29-33, 1970. A 568
- LATALSKI M and PAWLOWSKA A : Badania nad ultrastrukturą komórek wątrobowych po zatruciu gazem świetlnym. (Ultrastructure of the liver cells in lighting gas poisoning.) Patol Pol 20391-6, 1969. A 569
- LATALSKI M and PAWLOWSKA A : Fine structure of liver cell after lighting gas poisoning. Pol Med J 19: 891-5, 1970. A 570
- LEVIN W and KUNTZMAN R : Biphasic decrease of radioactive hemoprotein from liver microsomal CO-binding particles. J Biol Chem 244: 3671-6, 1969. A 571
- LEVIN W and KUNTZMAN R : Biphasic decrease of radioactive hemoprotein from liver microsomal carbon monoxide-binding particles. Effect of phenobarbital and chlordan. Mol Pharmacol 5: 499-506, 1969. A 572
- MAZALESKI S C, COLEMAN R L, DUNCAN R C and NAU C A : Subcellular trace metal alterations in rats exposed to 50 PPM of carbon monoxide. Am Industr Hyg Ass J 31: 183-8, 1970. A 573

1005051276

- MONTGOMERY M R and RUBIN R J : The effect of carbon monoxide inhalation on in vivo drug metabolism in the rat. J Pharmacol Exp Ther 179: 465-73, 1971. A 574
- NIZHEGORODOV V M and MARCHOZKY J D : (The effect of small concentrations of carbon monoxide and nitrogen oxides on the status of the supply and need of the animal organism for vitamin B-6) Gig Sanit 34: 96-7, 1969. A 575
- OMURA T and SATO R: The carbon monoxide-binding pigment of liver microsomes. I. Evidence for its hemoprotein nature. J Biol Chem 239: 2370-8, 1964. A 576
- RIKANS L E and VAN DYKE R A : Evidence for a different CO-binding pigment in solubilized rat hepatic microsomes. Biochem Pharmacol 20: 15-22, 1971. A 577
- RONDIA M D : Abaissement de l'activité de la benzopyrène-hydroxylase hépatique in vivo après inhalation d'oxyde de carbone. (Lowering of the activity of hepatic benzopyrene-hydroxylase in vivo after carbon monoxide inhalation.) C R Acad Sci Paris 271: 617-9, 1970. A 578
- WARBURG O, GEISLERAWand LORENZ S : Bemerkung über die Tryptophan-Oxygenase. (Note on tryptophan oxygenase.) Hoppe-Seyler Z Physiol Chem 348: 899-901, 1967. A 579
- WILSON L D, NELSON D H and HARDING B W : A mitochondrial electron carrier involved in steroid hydroxylations. Biochim Biophys Acta 99: 391-3, 1965. A 580
- WOHLRAB H and OGUNMOLA B G : Carbon monoxide binding studies of cytochrome  $a_3$  hemes in intact rat liver mitochondria. Biochemistry 10: 1103-6, 1971. A 581

1005051277

## I. INTRODUCTORY REMARKS

In recent years there has been increasing concern as to the harmful effects of carbon monoxide released as an air pollutant. The importance of cigarette smoking as a source of carbon monoxide has been recently stressed. This review is an attempt to clarify the role of carbon monoxide in cigarette smoking. The relationship will be analyzed in terms of carboxyhemoglobin blood levels and their influence on the respiratory, circulatory, nervous, renal, reproductive, endocrine and musculoskeletal systems.

At the outset it is pertinent to summarize the present state of knowledge relating to carbon monoxide in general and to carbon monoxide in cigarette smoke in particular. The information is summarized in the following publications:

a. The toxicity of carbon monoxide has been reviewed by Sayers and Davenport (1930), Killick (1940), Lilienthal (1950), Root (1962) and Theodore et al. (1971). These review articles have appeared at intervals of a decade and do not include the cigarette smoke as a source of carbon monoxide.

b. The importance of carbon monoxide as an air pollutant has been reviewed by Goldsmith (1964), Kaye (1965), Finck (1966), Giever (1967), Goldsmith and Landaw (1968), Beard (1969), Leclercq and Proteau (1970), Casarett (1971) and Jech (1972). These reviews appearing at yearly intervals emphasize the origin of carbon monoxide poisoning from sources other than cigarette smoking.

1005051069

- (26) DOYLE J T : Smoking and myocardial infarction. Circulation 39 & 40: Suppl 4: 136-43, 1969.

This review contains a paragraph on the role of carbon monoxide in pathogenesis of atherosclerosis. The author views the problem in the proper perspective.

The manner in which cigarette smoking accelerates atherosclerosis and its complications is, in short, unexplained. It is possible that in some way cigarette smoking damages the arterial intima. Carbon monoxide is the likeliest immediate candidate for such a role. Some presently mysterious interference with the normal mechanism of transport of lipids from the plasma through the vascular tunics to the lymphatics secondary to the inhalation of cigarette smoke is an alternative possibility: in all populations yet scrutinized, the prevalence and incidence of CHD rise with the serum cholesterol concentration.<sup>22</sup> It is, accordingly, a plausible hypothesis that inordinate cigarette smoking may be associated with an increased serum cholesterol concentration. Such a relationship does, indeed, exist, but is unimpressive. Although the serum cholesterol concentration in both men and women is consistently higher in cigarette smokers, the influence of increasing age is substantially greater (figs. 1 and 2).<sup>23</sup> The observation that heavy cigarette smokers have far more atheroma than nonsmokers is, possibly, complemented by Astrup's observation that fat-fed rabbits exposed to high tensions of carbon monoxide exhibit extreme hyperlipidemia and cholesterosis as compared to controls not exposed to carbon monoxide.<sup>24-26</sup> This interesting experimental model has, however, no recognized counterpart in human epidemiological studies. Obesity as a coronary risk factor is not related to cigarette smoking.<sup>2</sup> Lastly, the arterial blood pressure is not associated with cigarette habit.<sup>2</sup>

1005051226



## Additional Bibliography List No. 14

## ACCIDENTAL POISONING INVOLVING CARBON MONOXIDE AND OTHER CAUSATIVE FACTOR

- AITKEN R C B, BUGLASS D and KREITMAN N : The changing pattern of attempted suicide in Edinburgh, 1962 - 67. Brit J Prev Soc Med 23: 111 - 115, 1969. Reprint A 617
- AMENDT P and REDDEMANN H : Akzidentelle vergiftungen im kindesalter. (Accidental poisoning in childhood). Z Gesamte Inn Med 26: 188 - 90, 1970. A 618
- ANON : Burnt children. Brit Med J 1: 790, 1969. A 619
- BEAN, W B : President's address. The ecology of the soldier in World War II. Trans Am Clin Climatol Assoc 79: 1 - 12, 1968. A 620
- BRANDENBERGER H : Résolution de quelques problèmes de chimie toxicologique et légale par absorption atomique. (Solution of several problems of toxicologic and legal chemistry by atomic absorption). Ann Biol Clin (Paris) 25: 1053-62, 1967. A 621
- BURSTON G R : Self-poisoning in elderly patients. Geront Clin (Basel) 11: 279-89, 1969. A 622
- BURVILL P W : Methods of suicide in Western Australia. Med J Aust 2: 411-4, 1970. A 623
- DAVID A : Sebevražedné otravy toxickými průmyslovými látkami. (Suicidal poisonings using industrial poisons). Cas Lek Cesk 110: 118-22, 1971. A 624
- FARBEROW N L and SIMON M D : Suicides in Los Angeles and Vienna. An intercultural study of two cities. Public Health Rep 84: 389-403, 1969. A 625
- FATINI G and GALLENGA G : Considerazioni statistiche su 900 casi di morte improvvisa. (Statistical considerations on 900 cases of sudden death) Osped Ital Chir 18: 41-9, 1968. A 626
- FRANCOIS R C and BERTIN M : Etude statistique de la fréquence des certaines manifestations cliniques au cours de l'intoxication aigue par l'oxyde de carbone. Leur valeur pronostique immédiate. (Statistical study of the frequency of certain clinical manifestations during acute carbon monoxide poisoning. Their immediate prognostic value). Rass Med Industr 33: 380-91, 1964. A 627
- GAULTIER M, FREJAVILLE J P, BISMUTH C and PEBAY-PEYROULA F : Analyse des dossiers d'hypoxies survenues lors d'intoxications aiguës à l'hôpital Fernand-Widal. (Analysis of histories of cases of hypoxia during acute poisoning seen at the Hospital Fernand-Widal). Poumon Coeur 26: 931-6, 1970. A 628
- GOULDING R : The role of poisons information centres. Practitioner 194: 120-5, 1965. A 629
- GRAHAM J D P and HITCHENS R A N : Trends in hospitalized accidental poisoning. Brit J Prev Soc Med 22: 55-58, 1968. A 630
- GREGORY K L, MALINOSKI V F and SHARP C R : Cleveland clinic fire survivorship study, 1929-1965. Arch Environ Health 18: 508-15, 1969. A 631
- KAYE S : Bedside toxicology. Pediatr Clin North Am 17: 515-24, 1970. A 632
- KIM M W and PARK C S : Carbon monoxide poisoning in Korea. Proc Aust Ass Neurol 5: 385-90, 1968. A 633
- LACHNIT V : Vergiftungen in der krankenhauspraxis. (Poisoning in hospital practice). Wien Z Inn Med 45: 242-6, 1964. A 634

1005051280

- LITMAN R E : Psychological-psychiatric aspects in certifying modes of death. J Forensic Sci 13: 46-54, 1968. A 635
- MALIK M O A : Problems in the diagnosis of the causes of death in burned bodies. J Forensic Sci Soc 11: 21-8, 1971. A 636
- MATTHEW H : Acute poisoning: some myths and misconceptions. Br Med J 1: 519-22, 1971. A 637
- PRELLWITZ W, SCHUSTER H P, SCHYLLA G, BAUM P, SCHÖNBORN H, UNGERN-STERNBURG A, BRODERSEN H C and POEPLAU W : Zur differential diagnose von organbeteiligungen bei exogenen intoxicationen mit hilfe klinischer und klinisch-chemischer untersuchungen. (Differential diagnosis of organ involvement in exogenous poisoning by means of clinical and clinico-chemical studies). Klin Wochenschr 48: 51-3, 1970. A 638
- REDDEMANN H, AMENDT P and JÄHRIG K : Spätprognose akzidenteller vergiftungen im kindesalter. (Late prognosis of accidental poisoning in childhood). Dtsch Gesundheitsw 25: 2027-32, 1970. A 639
- ROPSCHITZ D H and OVENSTONE I M K : Two years' survey on self-aggressive acts, suicides and suicidal threats in the Halifax district between 1962 and 1964. Part I. Self-aggressive acts in the Halifax area. Int J Soc Psychiat 14: 165-87, 1968. A 640
- STURNER W Q : Some perspectives in "cot death". J Forensic Med 18: 96-107, 1971. A 641
- SZUCHOVSKY G, KENYERES I and HARSÁNYI L : Analyse von 2956 vergiftungs-selbstmordfällen. (Analysis of 2956 cases of suicidal poisoning). Deutsch Z Ges Gerichtl Med 66: 19-29, 1969. A 642
- VARESE L A and SORANZO L : Avvelenamenti acuti nell'infanzia. Avvelenamenti da ossido di carbonio. (Acute poisoning in children. VII. Carbon monoxide poisoning). Minerva Pediat 20: 871-83, 1968. A 643
- VIEWEG C, GRÜNEWALD G and ZIEGLER C : Die akute vergiftung. (Acute intoxication). Dtsch Gesundheitsw 25: 2480-5, 1970. A 644
- VOGEL, C : Vergiftungen im kindesalter. Eine klinische zehnjahresübersicht. (Poisonings in childhood. A 10-year clinical study). Z Aerztl Fortbild (Jena) 62: 486-91, 1968. A 645

1005051281

## Additional Bibliography List No. 15

## ACCIDENTAL POISONING INVOLVING CARBON MONOXIDE

- en dehors
- ALEXANDRE A : La pollution de l'atmosphère par l'oxyde de carbone/les lieux de travail. Dépistage. Manifestations cliniques. (Atmospheric pollution due to carbon monoxide outside places of employment. Detection. Clinical manifestations). Acta Tuberc Belg 58: 905-9, 1967. A 646
- ANDERSON T B : Natural gas, unnatural causes. Lancet 1: 466, 1970. A 647
- ANON : Natural gas: friend or foe? Lancet 2: 699-700, 1970. A 648
- ANON : Carbon monoxide poisoning. Brit Med J 1: 180, 1970. A 649
- ARONDEL E, GAUBERTI P and ROCHET J : Six cas d'anoxie accidentelle par appareils de chauffage mobiles a gaz butane. (Six cases of accidental anoxia caused by a mobile butane gas heating appliance). Ann Med Leg (Paris) 44: 443-5, 1964. A 650
- BERTIN M, FRANCOIS R C, PÉQUIGNOT H and SOULAIRAC A : Épidémiologie des asphyxies par le gaz. Remarques méthodologiques. (Epidemiology of asphyxia due to gas. Methodological remarks). Sem Hop Paris 46: 2657-71, 1970. A 651
- BETHEUIL M J and DELAHAYE-POLOUVIER G : A propos de deux comas oxycarbonés. (Apropos of 2 carbon monoxide comas). Anesth Analg (Paris) 24: 439-41, 1967. A 652
- BOZEK J, PAJOR Z and WASOWICZ Z : Przypadek ciężkiego zatrucia tlenkiem węgla dziecka 6-letniego. (Case of severe carbon monoxide poisoning in a 6-year old child). Pediat Pol 40: 1271-3, 1965. A 653
- BOUR H : L'intoxication oxycarbonée. (Carbon monoxide poisoning). Gaz Med France 71: 2793-814, 1964. A 654
- BURG F D and DOUGLASS J M : In cold gas. Safeguards against carbon monoxide poisoning. Clin Pediat (Phila) 8: 590-3, 1969. A 655
- CALLIGARI G : Intossicazione da ossido di carbonio, cianuri e sostanze metaemoglobinizzanti. (Carbon monoxide, cyanide and methemoglobinizing substance poisoning). Minerva Anest 34: 1109-16, 1968. A 656
- CHERKAVSKIR N B : (Carbon monoxide poisoning). Voen Med Zh (Russia) 11: 34-5, 1970. A 657
- CROSETTI L, PETTINATI L and RUBINO G F : Alcune considerazioni in tema di prevenzione tecnica e biologica dell'intossicazione da ossido di carbonio. (Considerations on the technical and biological prevention of carbon monoxide poisoning). Med Lavoro 56: 604-12, 1965. A 658
- CUCHE M M and BERNARD C : A propos de quelques formes particulières d'intoxication oxycarbonée aiguë. (Apropos of various special forms of acute carbon monoxide poisoning). Rev Lyon Med 18: 217-22, 1969. A 659
- DALGAARD J B : 550 Kohlenoxyd-todesfälle. (550 deaths from carbon monoxide poisoning). Acta Med Leg Soc (Liege) 18: 25-37, 1965. A 660
- DIMIZIO M, FELICI F and BIETTI C : Su un caso di avvelenamento da ossido di carbonio: contributo clinico. (On a case of carbon monoxide poisoning: clinical case). Riv Neurol 39: 425-34, 1969. A 661
- DOUZE J M : Koolmonoxydevergiftiging door aardgas. Douchen, een gevaar? (Carbon monoxide poisoning from natural gas. Showering, a danger?). Ned Tijdschr Geneesk 115: 1487-93, 1971. A 662

1005051282

- DOUZE J M C, HEYST A N P, KREUKNIET J, LEEUW R J M and HAMELINK M L : Koolmonoxi-devergiftigen. Is gebruik van aardgas wel zo ongevaarlijk? Pharm Weekblad 102: 351-60, 1967. A 663
- FELDMAN I G and LAMPERT F F : (Air pollution by highway photooxidants on various floors of apartment houses). Gin Sanit (Russia) 33: 89-91, 1968. A 664
- FISHER T L : Carbon monoxide poisoning: Legal responsibility for failure to diagnose. Canad Med Ass J 99: 235-6, 1968. A 665
- FRYZE C, GRUSZKA E and ZAWADZKI W : Zatrucie gazem swietlnym (tlenkiem wegla) o bardzo ciezkim przebiegu. (Case of very severe carbon monoxide poisoning). Przeg Lek (Pol) 26: 723-4, 1970. A 666
- GILL G S : The Jiko and carbon-monoxide poisoning. East Afr Med J 48: 85-7, 1971. A 667
- GLASS F, MALLACH H J and WOJAHN H : Über den CO-Gehalt des blutes bei akuter Kohlenmonoxyd-Vergiftung. (On the CO-content of the blood in acute carbon monoxide poisoning). Arzneimittelforschung (Germany) 16: 1553-5, 1966. A 668
- GOLACKA K, JANIK-KURYLICIO S and ROZEK S : Zatrucia tlenkiem wegla. (Carbon monoxide poisoning). Pol Tyg Lek 24: 1813-5, 1969. A 669
- GOLDSMITH J R : Carbon monoxide. Science 157: 842-4, 1967. A 670
- GRÉMY F, SALMON D, FRANCOIS R C and BERTIN M : Asphyxies accidentelles et volontaires par le gaz d'éclairage survenues a Paris de 1949 a 1962. (Accidental and voluntary asphyxiations by coal gas occurring in Paris from 1946 to 1962. Statistical study. Particular influence of the concentration of carbon monoxide and atmospheric conditions). Presse Med (France) 76: 1099-102, 1968. A 671
- GRIFFITHS J C : North Sea gas. Lancet 1: 562, 1970. A 672
- GUEYE S N, BA H and DIOP I : L'intoxication oxycarbonée à Dakar. (Carbon monoxide poisoning in Dakar). Bull Soc Med Afr Noire Lang Fr 10: 422-5, 1965. A 673
- HUGHES E R and FISHER D A : Carbon monoxide poisoning. J Arkansas Med Soc 62: 255-60, 1965. A 674
- KASHIMA T, FUKUI M, MASUDA Y, WAKASUGI C and HAYAMA R : Report of five cases where ordinary vinyl bag was used for suicidal purpose. Suffocation, CO-poisoning and "thinner"-poisoning. Jap J Leg Med 23: 248-52, 1969. A 675
- KITTREDGE R D : Pulmonary edema in acute carbon monoxide poisoning. Am J Roentgenol Radium Ther Nucl Med 113: 680-1, 1971. A 676
- KLINGHOFFER M : Carbon monoxide poisoning. Illinois Med J 128: 310-2, 1965. A 677
- KRISHNAN S, KUPPUSWAMY G, MANI S and MAJID M A : Carbon monoxide poisoning. J Assoc Physicians India 19: 409-11, 1971. A 678
- KUROIWA Y : Carbon monoxide poisoning. Jap J Clin Med 25: 1647-52, 1967. A 679
- LAMY M and NOIRFALISE A : Intoxication oxycarbonée et gaz naturel. (Carbon monoxide and natural gas poisoning). Rev Med Liege 26: 128-33, 1971. A 680
- LeBRETON R and GARAT J : Intoxications oxycarbonées aiguës provoquées par fautes d'utilisateurs de poêles à combustibles solides. (Acute carbon monoxide poisoning caused by negligence in using a solid-fuel stove). Ann Med Leg (Paris) 44: 430-5, 1964. A 681

1005051283

## Additional Bibliography List No. 15

- LeBRETON R and GARAT J: Intoxications oxycarbonées aiguës provoquées par fautes de'utilisation de appareils à combustibles gazeux ou liquides. (Acute monoxide poisoning caused by negligence in using gaseous or liquid fuel appliances). Ann Med Leg (Paris) 41: 435-43, 1964. A 682
- LEHR E L: Carbon monoxide poisoning: A preventable environmental hazard. Am J Public Health 60: 289-93, 1970. A 683
- MACHATA G: Die differenzierung der Kohlenoxidvergiftung. (The distinction of carbon monoxide poisoning). Arch Toxik (Germany) 23: 136-40, 1968. A 684
- MARI E and RIZZATTI E M: Ulteriori osservazioni in tema di intossicazione da ossido di carbonio e sindrome generale dell'adattamento. (Latest observations on the subject of carbon monoxide poisoning and general adaptation syndrome). Minerva Medicoleg 84: 162-7, 1964. A 685
- McDAY A J: Law-Medicine notes. Carbon monoxide poisoning. New Eng J Med 272: 252-3, 1965. A 686
- McNALLY W D: Carbon monoxide poisoning. Illinois Med J 59: 383-88, 1931. A 687
- MENZ M J: Ein pseudogift. (A pseudo-poison). Int Arch Gewerbepath (Germany) 22: 230-5, 1966. A 688
- MÜLLER G M and HUNG N: Einige daten zu fragen der tödlichen CO-vergiftung. (Some data on the problem of lethal CO-poisoning). Z Aerztl Fortbild (Jena) 62: 761-6, 1968. A 689
- MUROFUSHI K and MINAGAWA M: (Case of carbon monoxide poisoning with aging phenomena). Advances Neurol Sci (Tokyo) 13: 39-48, 1969. A 690
- OBERSTEG J I and DELAY F: 6 Jahre erfahrungen mit entgiftetem kochgas. (6 years of experience with detoxicated cooking gas.) Deutsch Z Ges Gerichtl Med 58: 122-6, 1966. A 691
- PETIT A G, PETIT G and GEILLE A: Contribution a l'étude étiologique de l'intoxication oxycarbonée. (Etiology of carbon monoxide poisoning). Med Leg Domm Corpor (Paris) 3: 278-82, 1970. A 692
- POLIARD J A: Burns and carbon-monoxide poisoning. Brit J Surg 57: 635-7, 1970. A 693
- RITUCCI A and LUVONI R: Rilievi medico-legali su casi plurimi mortali di acuta intossicazione ossi-carbonica con riferimento alla cronologia della morte. (Medico-legal findings on plural fatal cases of acute carbon monoxide intoxication with reference to chronology of death). Minerva Medicoleg 85: 45-51, 1965. A 694
- ROSE E F: Carbon monoxide intoxication and poisoning. J Iowa Med Soc 59: 909-17, 1969. A 695
- ROSENTHAL S G: Lethality of carbon monoxide in fires. New Eng J Med 279: 111-2, 1968. A 696
- SATOH K, KIYOTANI T, MINAGI Y and KONDO M: Isolation and identification of narcotics by thin layer chromatography. 4. Isolation and identification of narcotics in carbon monoxide poisoning. Jap J Leg Med 20: 6-10, 1966. A 697
- SIASEV A N: (A severe case of carbon monoxide poisoning with a favorable outcome). Vrach Delo 11: 126-7, 1966. A 698
- SMITH J S and BRANDON S: Acute carbon monoxide poisoning - 3 years experience in a defined population. Postgrad Med J 46: 65-70, 1970. A 699
- STICHINOTH E and ZUMBANSEN H: Über farbveränderungen von leichenorganen nach CO-vergiftung bei spätexhumierung. (Changes in color of cadaver organs following CO-intoxication in late exhumation). Beitr Gerichtl Med 27: 237-41, 1970. A 670

1005051284

Additional Bibliography, List no. 15

- SUCHCICKI L : Uwagi medyka sądowego dotyczące wczesnego rozpoznania zatrucia tlenkiem węgla. (Remarks of a forensic medicine specialist concerning the early diagnosis of carbon monoxide). Wiad Lek 23: 1563-5, 1970. A 671
- THOMSON W : Carbon monoxide poisoning. Nurs Times 67: 668-9, 1971. A 672
- TOTSUKA S, MORO K, HORIE T and YAZAKI M : (Continuous and prolonged symptoms in 2 cases of gas poisoning). Adv Neurol Sci (Tok) 15: 592-605, 1971. A 673
- TOYA G : A clinical study on carbon monoxide poisoning. J Kumamoto Med Soc 41: 377-420, 1967. A 674
- TRAKHTENBERG S A : (The clinical picture of acute carbon monoxide poisoning). Klin Med (Moskva) 44: 112-5, 1966. A 675
- TUTT J B : Carbon monoxide poisoning. Vet Rec 87: 107-8, 1970. A 676
- VICH V : (Problems of carbon monoxide poisoning). Cas Lek Cesk 108: 1485, 1969. A 677
- VIOLET M M and PERROT E : Intoxications par l'oxyde de carbone. (Carbon monoxide poisoning). J Med Lyon 48: 81-6, 1967. A 678
- VIVOLI G and PREITE E : Sulla corruzione dell'aria degli ambienti riscaldati da stufe a gas liquido. (On air pollution in rooms heated with liquid gas stoves). Riv Ital Ig 26: 113-26, 1966. A 679
- VORONCHUCK G Z : (A rare case of severe carbon monoxide poisoning). Klin Med (Moskva) 44: 115-6, 1966. A 680
- WATANABE S, KITAGUCHI T, KIYOFUJI T, MORISAKI Y, MASUDA T, NOGUCHI K and MATSUMOTO S : An autopsy case of CO gas poisoning by incomplete combustion of fuel propane gas. J Kumamoto Med Soc 44: 354-8, 1970. A 681
- WATSON E S : Four cases of carbon monoxide poisoning in one family. S Dakota J Med 21: 15-8, 1968. A 682
- WEIGT H : Ungewöhnlicher verlauf einer Kohlenoxydgasvergiftung bei einem Ehepaar. (Unusual course of carbon monoxide poisoning in a married couple. Psychiat Neurol Med Psychol (Leipzig) 19: 147-52, 1967. A 683
- WILSON E F, RICH T H and MESSMAN H C : The hazardous hibachi. Carbon monoxide poisoning following use of charcoal. JAMA 221: 405-6, 1972. A 684
- ZORN H : Zur diagnostik der chronischen kohlenoxydvergiftung. (On the diagnosis of chronic carbon monoxide poisoning). Russ Med Industr 33: 325-9, 1964. A 685

1005051285

## Additional Bibliography List No. 16

## INDUSTRIAL POISONING INVOLVING CARBON MONOXIDE

- ALEKSANDROV V N, BESCHICHLOW D I, DAVIDOV O V : (Pathogenesis of poisoning by gun-powder gases). Voen Med Zh 7: 28-30, 1970. Reprint A 686
- ANON : Hazards in gasworks. Brit Med J 5439: 876-7, 1965. A 687
- BASMADZHEVA K, KURCHATOWA G, DAVYDKOVA E and TSWETANOV J : (The character of the combined action of sulfurous gas and carbon monoxide in their simultaneous presence in the atmosphere). Gig Sanit (Russia) 33: 73-7, 1968. A 688
- BRIGATTI L, PARIGI A and VARETTO L : Sul rischio di ossicarbonismo in una industria metal-meccanica. (On the risk of carbon monoxide poisoning in a metal machine factory). Rass Med Industr (Italy) 33: 417-9, 1964. A 689
- BUCKLEY A R and FEAR E C : Precautions taken for the protection of workers in the British gas industry against carbon monoxide poisoning. Rass Med Industr 33: 296-308, 1964. A 690
- CAPELLARO F : Aspetti di prevenzione medica in una moderna officina di produzione di gas. (Aspects of medical prevention in a modern gas production plant). Rass Med Industr 33: 398-400, 1964. A 691
- CAPELLARO F and BRAGUZZI E : Su alcuni fattori limitanti l'idoneita' lavorativa in ambienti con rischio di intossicazione da co. (On some factors limiting work ability in areas with risk of carbon monoxide poisoning). Folia Med (Napoli) 47: 1068-73, 1964. A 692
- CASCINI F and GAZZERRO F : Saldatura elettrica ed ossicarbonismo. (Electrical welding and carbon monoxide poisoning). Folia Med (Napoli) 49: 433-46, 1966. A 693
- COSIC V, ARSENIJEVIC M, KANDIC B and GRBESA B : Akutno trovanje ugljen-monoksidom u rudniku Banovici. (Acute carbon monoxide poisoning in the mine). Vojnosanit Pregl 21: 157-64, 1964. A 694
- DATSENKO I I : (The carbon monoxide content of the air in the foundry shop of the "Avtopogruzchik" factory). Gig Tr Prof Zabol 9: 55, 1965. A 695
- DEANE M, GOLDSMITH J R and TUMA D : Respiratory conditions in outside workers. Arch Environ Health (Chicago) 10: 323-331, 1965. A 696
- DE KRETZER A J, EVANS W D and WALDRON H A : Carbon monoxide hazard in the CO<sub>2</sub> arc-welding process. Ann Occup Hyg 7: 253-9, 1964. A 697
- DESBAUMES P : Intoxications mortelles par les gaz de fermentation de silos agricoles (oxyde de carbone et oxydes d'azote). (Fatal poisoning by silo gases (carbon monoxide and oxides of nitrogen). Arch Toxic 23: 160-4, 1968. A 698
- DETTORI G and SCANSETTI G : Importanza del controllo dell'efficienza degli impianti industriali ai fini della prevenzione dell'ossicarbonismo. (Importance of the inspection of the efficiency of industrial installations for the purpose of prevention of carbon monoxide poisoning). Rass Med Industr 33: 394-7, 1965. A 699
- FIRST M W and MURPHY R L H : Carbon monoxide exposures from snow melting machines. Am Ind Hyg Assoc J 31: 754-7, 1970. A 700
- GERITSEN W B : Quelques aspects de la prevention de l'oxycarbonisme des travailleurs. (Some aspects of the prevention of carbon monoxide poisoning in workers). Rass Med Industr 33: 356-9, 1964. A 701

1005051286

## Additional Bibliography List No. 16

- GUEPIN : A propos de'un cas d'oxycarbonisme présumé en cale. (A case of poisoning by carbon monoxide presumably from a ship's hold). Arch Mal Prof 30: 355-7, 1969. A 702
- JORDI A : Gewerbliche toxiologie und soziale medizin. (Industrial toxicology and social medicine). Praxis 57: 785-93, 1968. A 703
- KEMKES B : Berufskrankheiten durch gasförmige gifte. (Occupational gaseous poisonings). Med Klin 37: 109-11, 1941. A 704
- LUNDEVALL J : Carbon monoxide intoxication from insulation of electric cable. Med Sci Law 12: 203-4, 1972. A 705
- MARANZANA P, TRONZANO L and COSCIA G : Su due successivi episodi di intossicazione da ossido di carbonio in uno stesso gruppo di operai friggitori. (On 2 successive episodes of carbon monoxide poisoning in the same group of workers engaged in potato-chip frying). Rass Med Industr 33: 407-10, 1964. A 706
- MARCHIARO G, MARGARIA E, GAIDO P C and AQUARO G : Equilibrio acido base nell'intossicazione acuta sperimentale da ossido di carbonio. (Acid-base equilibrium in acute experimental carbon monoxide poisoning). Rass Med Industr 33: 452-3, 1964. A 707
- MONACO B : L'ossicarbonismo e la determinazione del CO negli ambienti di lavoro ricerche in una grande officina per la produzione del gas di città. (Carbon monoxide poisoning and carbon monoxide determination in work areas. Research in a large workshop for city gas production). Rass Med Industr 33: 495-527, 1964. A 708
- MOSS C J : Machinery hazards. Ann Occup Hyg 12: 69-75, 1969. A 709
- REPLOH H, KLOSTERKÖTTER W and EINCK-ROSSKAMP P : Zur frage der toxizität von kunststoff-verschmelungsprodukten. (On the problem of the toxicity of products of the carbonization of plastics). Arch Hyg Bakt 150: 393-405, 1966. A 710
- RODKEY F L, COLLISON H A and ENGEL R R : Release of carbon monoxide from acrylic and polycarbonate plastics. J Appl Physiol 27: 554-5, 1969. A 711
- ROSSO M and DUGHERA L : Problemi d'inquinamento da CO in ambiente metalmeccanico pesante. (Problems of air pollution with CO in heavy metal machine factories). Rass Med Industr 33: 354-5, 1964. A 712
- SADOKIERSKI W : Obraz krwi obwodowej u pracowników zakładów suchej destylacji drewna. (Peripheral blood picture in workers employed in dry wood distillation). Med Pracy 16: 61-5, 1965. A 713
- SCHAFFERNICHT H, ZIEGLER V and REINHARD H : Untersuchungen über die gefährdung durch Kohlenmonoxid in Räuchereien. (Studies on carbon monoxide hazards in smoke-houses). Z Gesamte Hyg 16: 503-8, 1970. A 714
- SCHIECHE M, KE LER M and KOBER S : Erhebungen zum Krankenstand eines hochofenbetriebes. (Data on sickness in a blast furnace steel plant). Z Aerztl Fortbild (Jena) 64: 617-21, 1970. A 715
- SCHÜTTMANN W : Berufliche intoxicationen unter dem bilde vegetativer dysregulationen. (Occupational poisoning under the clinical picture of vegetative dysregulations). Z Aerztl Fortbild (Jena) 62: 1-8, 1968. A 716
- SHIRABE T, MAWATARI S and KUROIWA Y : (Autopsy case of carbon monoxide poisoning at the Miike coal mine explosion. A case of lung cancer with the fatal outcome in 3 years and 4 months). Advances Neurol Sci (Tokyo) 321-6, 1970. A 717

1005051287



## Additional Bibliography List No. 16

- SIEGRIST H: Die gefährlichkeit der Flüssiggase. (The danger of liquid gases). Deutch Z Ges Gerichtl Med 57: 158-61, 1966. A 718
- STEWART R D, FISHER T N, HOSKO M J, PETERSON J E, BARETTA E D and DODD H C: Carboxyhemoglobin elevation after exposure to dichloromethane. Science 176: 295-6, 1972. A 719
- SUPFLE: Zur frage der chronischen Kohlenoxydvergiftung. (Chronic carbon monoxide poisoning). Med Klin 29: 1727, 1933. A 720
- TATETSU S, TOYA G, MIMURA K, HARADA M and TSUKAYAMA T: (A four-year clinical follow-up study of mass carbon monoxide poisoning due to a coal-mine explosion.) Advances Neurol Sci (Tok) 13: 11-19, 1969. A 721
- TEBBENS B D and SPEAR R C: Quality control of work environments. Am Ind Hyg Assoc J 32: 546-51, 1971. A 722
- THURSTON G: The lethal boiler flue. Medicoleg J 36: 191-6, 1968. A 723
- VENGERSKAIA K, NAZYROV G N, BOPROVA L S, DUBROVSKY S J and DUMKO V P: (Sanitation-chemical assessment of the use of new synthetic materials under hot climatic conditions). Gig Sanit 33: 23-4, 1968. A 724
- VUOPALA U, HUHTI E, TAKKUNEN J and HUIKKO M: Nickel carbonyl poisoning. Report of 25 cases. Ann Clin Res 2: 214-22, 1970. A 725
- WAGNER F J: Reihenuntersuchungen bei Kohlenoxydexponierten. (Mass examinations of persons exposed to carbon monoxide). Z Ges Hyg 10: 621-7, 1964. A 726
- WOHLERS H C, NEWSTEIN H and DAUNIS D: Carbon monoxide and sulfur dioxide adsorption on - and desorption from glass, plastic and metal tubings. J Air Pollut Contr Ass 17: 753-6, 1967. A 727
- ZARIVAISKALA K: (Pollution of the air of gas-supplied dwellings with various ventilation systems). Gig Sanit 31: 100-2, 1966. A 728

1005051288

## Additional Bibliography List No. 17

## THERAPY OF CARBON MONOXIDE POISONING

- AQUINAS M : Coal gas poisoning: a nursing care study. Nurs Times 60: 1208-10, 1964. Reprint A 729
- BEGHE<sup>1</sup> RAOUL : Esperienze in tema di trattamento dell'ossicarbonismo acuto. (Experience in the treatment of acute carbon monoxide poisoning). Rass Med Industr 33: 337-48, 1964. A 730
- BELAISCH J : La plasmathérapie. (Plasmatherapy.) Vie Med Paris 35: 999-1002, 1954. A 731
- BORBÉLY F : Die behandlung der kohlenoxydvergiftungen. (Treatment of carbon monoxide poisoning). Deutsch Med Wschr 90: 1963-4, 1965. A 732
- BOULETREAU P and MOTIN J : Etude d'une série homogène de 50 comas oxycarbonés traités par l'oxygénothérapie hyperbare. (Study of a homogeneous series of 50 cases of coma due to carbon monoxide poisoning and treated with hyperbaric oxygenation). Moroc Med 50: 386, 1970. A 733
- BOUR H, PASQUIER P and BERTRAND-HARDY J M : Le coma oxycarboné. Etude générale, clinique, biologique et thérapeutique de 290 cas. (Carbon monoxide coma. General study, clinical, biological and therapeutics of 290 cases). Sem Hop Paris 42: 1839-61, 1966. A 734
- BRANDON S : Treatment of carbon monoxide poisoning. Lancet 1: 626, 1970. A 735
- BURMEISTER H, BARCKOW D, HUMPERT U, IBE K and LERCHE D : Die künstliche beatmung. Ein erfahrungsbericht. (Artificial respiration. Report based on personal experience). Deutsch Med Wschr 93: 517-22, 1968. A 736
- BURMEISTER H and HEUHAU G A : Die behandlung der schweren subakuten leuchtgasvergiftung beim menschen. (Treatment of severe, subacute natural gas poisoning in humans). Arch Toxikol 26: 277-92, 1970. A 737
- CARUSO G and BARNABA A : Trattamento con diazepam e remissione della sintomatologia extrapiramidale in un caso di parkinsonismo da intossicazione con ossido di carbonio. (Treatment with diazepam and remission of the extrapyramidal symptoms in a case of parkinsonism caused by carbon monoxide poisoning). Acta Neurol Napoli 23: 103-10, 1968. A 738
- CENTI R and ZAFFIRI O : Considerazioni su di un caso gravissimo di ossicarbonismo acuto favorevolmente risolto con acido ascorbico ad alte dosi e con l-dopa. (On a very grave case of acute carbon monoxide poisoning favorably resolved with high doses of ascorbic acid with L-dopa). Minerva Anesthesiol 37: 406-14, 1971. A 739
- CHUKHRIENKO D P and LULKO A V : (Extrarenal hemodialysis in carbon monoxide poisoning). Vrach Delo 12: 18-21, 1968. A 740
- CIOCATTO E and PATTONO R : Organizzazione di un centro di rianimazione respiratoria terapia dell'intossicazione acuta da ossido di carbonio. (Organization of a respiratory resuscitation center. Therapy of acute carbon monoxide poisoning). Rass Med Industr 33: 330-3, 1964. A 741
- COLLINS J V and GOULDING R : Treatment of acute poisoning at Guy's Hospital: October 1969 to September 1970. Guys Hosp Rep 120: 31-46, 1971. A 742
- DAMIA G, SIBILIA E and DAMIA G : Considerazioni sulla prognosi e trattamento di alcuni casi di avvelenamento da barbiturici e da CO. (Considerations on the prognosis and treatment of some cases of poisoning due to barbiturates and carbon monoxide). Minerva Anest 31: 564-8, 1965. A 743

1005051289

## Additional Bibliography List No. 17

- ESTLER C J : Über Veränderungen des Hirnstoffwechsels nach akuter Kohlenmonoxydvergiftung und über den Einfluss von Natriumsuccinat auf die Kohlenmonoxydvergiftung. (On changes in brain metabolism following acute carbon monoxide poisoning and about the effect of sodium succinate on carbon monoxide poisoning). Arch Int Pharmacodyn 158: 415-28, 1965. A 744
- FOURNIER E : Traitement de l'intoxication oxycarbonée aiguë. (Treatment of acute carbon monoxide poisoning). Cah Coll Med Hop Paris 8: 1227-30, 1967. A 745
- GAULTIER M, FOURNIER E, GERVAIS P, EFTHYMIU M L, SRAER C, BISMUTH C, BODIN F, CHRISTOFOROV B, SICOT C and FREJAVILLE J P : Travail du secteur de réanimation de la clinique toxicologique pendant l'année 1967. (Work of the resuscitation center of the Toxicology Clinic during the year 1967). Bull Soc Med Hop Paris 119: 75-86, 1968. A 746
- GERBER H U and SEEWALD H : Pathophysiologie und Therapie der akuten Kohlenmonoxidvergiftung. (Pathophysiology and therapy of acute carbon monoxide poisoning). Z Aerztl Fortbild (Jena) 60: 1085-92, 1966. A 747
- GLÄSEL E : Häufigkeit und organisierte Bereitschaft bei der Behandlung exogener Intoxikationen. (Incidence and organized readiness in the treatment of exogenous intoxications). Z Gesamte Inn Med 7: 195-8, 1970. A 748
- GOTO H : (First aid in gas poisoning in large cities). Naika 22: 251-6, 1968. A 749
- GOULDING R : The treatment of acute poisoning. Trans Med Soc London 83: 9-18, 1967. A 750
- GOULON M, BAROIS A, RAPIN M, NOUAILHAT F, AUGUSTIN P, HENNETIER G, BAGUET J C, KUNTZIGER H and BRETEAU M : Traitement de l'intoxication oxycarbonée par l'oxygène hyperbare (à propos de 20 observations). (Treatment of carbon monoxide poisoning with hyperbaric oxygenation. Apropos of 20 cases). Bull Soc Med Hop Paris 116: 649-71, 1965. A 751
- GOULON M, BAROIS A, GAJDOS P, LABROUSSE J, SCHORTGEN G, AMERONGEN G and ROBERT J F : Problèmes physiopathologiques et thérapeutiques posés par l'œdème aigu pulmonaire toxique. (Physiopathologic and therapeutic problems raised by acute toxic pulmonary edema). Poumon Coeur 26: 1039-65, 1970. A 752
- HEYNDRICKX A, SCHEIRIS C, VERCROYSE A and OKKERSE E : Gas chromatographic determination of carbon monoxide in blood and the hyperbaric oxygen treatment in carbon monoxide poisoning cases. J Pharm Belg 25: 247-58, 1970. A 753
- KILLICK E M and MARCHANT J V : The effect of barbiturates on the resuscitation of dogs from severe acute CO poisoning. J Physiol London 180: 80-95, 1965. A 754
- LAVERNE A A : Carbon dioxide therapy, healing, and air pollution. A more effective rapid coma technic for psychiatric disorders. Behav Neuropsychiat 2: 10-25, 1970. A 755
- LEVIN M : Smoke poisoning treated by antacids. J Med Soc New Jersey 62: 215-6, 1965. A 756
- LIZANETS M N and ZARKEVICH U P : (Experience with intensive therapy in cases of acute poisoning by automobile exhaust and carbon monoxide. Voen Med Zh 11: 72-4, 1971. A 757
- MATTHEW H : Treatment of carbon monoxide poisoning. Lancet 1: 518, 1970. A 758
- NIELSEN B : Thermoregulation during work in carbon monoxide poisoning. Acta Physiol Scand 82: 98-106, 1971. A 759
- NORMAN J N and LEDINGHAM I M : Carbon monoxide poisoning: Investigations and treatment. Progr Brain Res 24: 101-22, 1967. A 760
- NOUAILHAT F : Conduite à tenir en présence d'une intoxication oxycarbonée aiguë. (Procedure to follow in the presence of acute carbon monoxide poisoning). Progr Med Paris 92: 537-45, 1964. A 761

1005051290

## Additional Bibliography List No. 17

- OKULOVSKI V K and HACHATUROV G S : (Resuscitation of patients in severe forms of carbon monoxide poisoning). Voen Zh 11: 66-7, 1968. A 762
- OTSUKI S, NINOMIYA K, YAMAMOTO M, NAKASHIMA Y, SHUTARA M, URAKAMI S and ITAMI Z : (Effect of cytidine monophosphate on the organic syndrome in the brain.) Brain Nerve (Tokyo) 18: 745-9, 1966. A 763
- PAPAVASILIOU P S, COTZIAS G C, DUBY S E, STECK A J, FEHLING C and BELL M A : Levodopa in parkinsonism: Potentiation of central effects with a peripheral inhibitor. N Engl J Med 285: 8-14, 1972. A 764
- PAULET G and CHEVRIER R: De l'action du cobalt sur la détoxication oxycarbonée. (On the action of cobalt on carbon monoxide detoxication). C R Soc Biol Paris 163: 1199-202, 1969. A 765
- PECORA L : La terapia ferrosa nell'ossicarbonismo acuto. (Iron therapy in acute carbon monoxide poisoning). Russ Med Industr 33: 352-3, 1964. A 766
- PLACE M : For each his own: the self-rescuer apparatus against CO. Occup Health London 22: 107-11, 1970. A 767
- PUKHOV V A : (Pathogenesis and complex treatment of CO poisoning). Biul Eksp Biol Med 60: 39-42, 1965. A 768
- RADUSHVICH V P : (Auxiliary extracorporeal circulation for resuscitation). Khirurgia Moskva 44: 3-8, 1968. A 769
- RAPOPORT K M : (On the feasibility of using methylene blue in carbon monoxide poisoning). Voen Zh 12: 34-5, 1967. A 770
- REJSEK K : (News in the therapy of poisoning by industrial agents). Cas Lek Cesk 110: 133-4, 1971. A 771
- RIZZI R : La rianimazione nell'avvelenamento da CO. (Resuscitation in CO poisoning). Minerva Anest 34: 1454-6, 1968. A 772
- SALNIS A N and HACHATUROV G S : (Resuscitation measures in the treatment of acute carbon monoxide poisoning). Voen Med Zh 11: 32-4, 1970. A 773
- SNASHALL P D : Treatment of carbon monoxide poisoning. Lancet 1: 357-8, 1970. A 774
- TATEGAMI T : Treatment of acute carbon monoxide poisoning from the point of view of pathophysiology. J Kumamoto Med Soc 42: 318-38, 1968. A 775
- THURSTON J : Treatment of carbon monoxide poisoning. Lancet 1: 468-9, 1970. A 776
- TORELLI G : Meccanismo dell'azione tossica da CO e principi terapeutici conseguenti. (Mechanism of the toxic action of CO and consequent therapeutic principles). Rass Med Industr 33: 458-64, 1964. A 777
- TOWNSEND G L and STETSON J B : Treatment of carbon monoxide poisoning by mechanical ventilation: Case report. Canad Anaesth Soc J 15: 184-96, 1968. A 778
- WAGNER K D and RICHTER D : Exogene Ursachen, Behandlung und klinische Spätfunde bei Vergiftungen im Kindesalter. (Exogenous causes, treatment and late clinical findings in poisoning during childhood). Z Aerztl Fortbild (Jena) 62: 492-7, 1968. A 779
- WATABE Y : (Emergency treatment in respiratory management) Sanfujinka Jissai 18: 496-506, 1969. A 780

1005051291

## V. CIRCULATORY SYSTEM

The investigation of circulatory effects of carbon monoxide has been more extensive than that of its effects on the respiratory system discussed in the preceding section. There has been increasing concern that chronic exposure to carbon monoxide present in cigarette smoke would lead to diseases of the heart and blood vessels and abnormalities in the composition of the blood. However, the problem has not been solved by direct experimentation relating to the carbon monoxide in cigarette smoke. There are numerous observations regarding the effects of sublethal concentrations of carbon monoxide in man and animals. These are reviewed in the following paragraphs, although they are only remotely related to the small amount of carbon monoxide contained in cigarette smoke.

### V A. Heart Rate

The acceleration of heart rate known to occur during inhalation of cigarette smoke is explained by the nicotine content. The amount of carbon monoxide in the smoke does not influence heart rate, since experiments consisting of administering carbon monoxide alone in amounts even exceeding that produced by cigarettes failed to alter the electrocardiogram.

The electrocardiograms of patients suffering from acute carbon monoxide poisoning or chronic exposure to carbon monoxide show the following alterations: depression of S-T segment (Steinmann, 1937; Störmer, 1938; Wendt, 1941; Graybiel, 1942; Breu, 1943; Patz, 1949; Saracoglu, 1951); sinus arrhythmia (Breu, 1942); premature systole (Parade and Franke, 1939); atrial flutter

1005051115

Additional Bibliography List No. 18

THERAPY OF POISONING BY USE OF NORMAL AND HYPERBARIC OXYGEN

- |  |                  |
|--|------------------|
| ALVIS H J : What hyperbaric medicine has to offer the industrial physician. <u>J Occup Med</u> 9: 304-7, 1967.   | Reprint<br>A 784 |
| ANON : Hyperbaric medicine. <u>Brit Med J</u> 1: 312-3, 1969.  | A 788            |
| ANON : Hyperbaric oxygen therapy. <u>Med Lett Drugs Ther</u> 13: 29-32, 1971.  | A 789            |
| ASCORBE DOMINGUEZ A : Cinco casos de intoxicacion por monoxido de carbono tratados con oxigeno hiperbarico. (5 cases of carbon monoxide poisoning treated with hyperbaric oxygenation). <u>Rev Esp Anest</u> 16:542-4, 1969.   | A 790            |
| BARTHÉLEMY L, PARC J, MICHAUD A and MATHÉ P : L'oxygène hyperbare dans le cadre de la marine nationale expérience de 10 ans. (Hyperbaric oxygen in the national Navy. Ten years' experience). <u>Anesth Analg Paris</u> 24: 375-86, 1967.  | A 791            |
| BEREZIN I P and PUZARES V A : (The therapeutic effect of oxygen under high pressure). <u>Klin Med Moskva</u> 47: 32-8, 1969.   | A 792            |
| BERNHARD W F and FILLER R M : Hyperbaric oxygenation: Current concepts. <u>Am J Surg</u> 115: 661-8, 1968.   | A 793            |
| BESZNYÁK I : Die hyperbare Sauerstoffbehandlung in der Medizin. (Hyperbaric oxygenation therapy in medicine). <u>Z Aerztl Fortbild (Jena)</u> 61: 985-91, 1967.  | A 794            |
| BIMONTE D, PORTOLANO F and TUFANO R : L'ossigenoterapia iperbarica nel trattamento della intossicazione acuta da ossido di carbonio. (Hyperbaric oxygen therapy in the treatment of acute carbon monoxide poisoning). <u>Rass Int Clin Ter</u> 51: 364-9, 1971.  | A 795            |
| BOCCALETTI E, NOFRINI U, JOSI G and MAGGIO G : L'ossigenoterapia iperbarica e le sue applicazioni cliniche. (Hyperbaric oxygen therapy and its clinical uses). <u>Policlinico (Prat)</u> 73: 1157-70, 1966.  | A 796            |
| BRUSADELLI S, PALMA G and DiPRETORO L : Il problema del monitoraggio durante il trattamento terapeutico in camera iperbarica. (The problem of monitoring during therapeutic treatment in the hyperbaric chamber). <u>Acta Anaesth (Padova)</u> 8: 363-81, 1968.  | A 797            |
| BULTERIJS A : Application of oxygen under high atmospheric pressure. <u>Int Nurs Rev</u> 12: 20-2, 1965.   | A 798            |
| BURSTON G R : Self-poisoning in elderly patients. <u>Geront Clin (Basel)</u> 11: 279-89, 1969.   | A 799            |
| CAILAR J, SERRE L, ROQUEFFEUIL B, LEFEBVRE F and MALZAC P : Résultats du traitement de l'intoxication oxycarbonée par l'oxygène hyperbare. A propos de 43 observations. (Results of the treatment with hyperbaric oxygen of carbon monoxide poisoning. Apropos of 43 cases). <u>Sem Hop Paris</u> 44: 3155-60, 1968. | A 800            |
| CHEW H E R, HANSON G C and SLACK W K : Hyperbaric oxygenation. <u>Brit J Dis Chest</u> 63: 113-39, 1969.   | A 801            |
| CHICHET-GOSSET C, CHASSON J, OHRESSER P, ARNAUD A and DUBOULOZ F : Les comas toxiques dans une unité de réanimation respiratoire. Bilan de trois années. (Toxic comas in a respiratory resuscitation unit. A 3-year observation). <u>Poumon Coeur</u> 23: 721-7, 1967.   | A 802            |
| CIOCATTO E, PATTONO R, QUERCI M and ZAFFIRI O : L'ossigenazione iperbarica in rianimazione. (Hyperbaric oxygenation in resuscitation). <u>Minerva Anest</u> 32: 97-9, 1966.  | A 803            |

1005051293

## Additional Bibliography List No. 18

- DEEN L : Hyperbare oxygenierung. (Hyperbaric oxygenation) Anaesthetist 18: 205-7, 1969 A 804
- DuCAILLAR J, LEFEBVRE F, ROQUEFEUIL B, MALZAC P and PROST T J : Oxygénothérapie hyperbare et intoxication oxycarbonée. A propos de 100 cas. (Hyperbaric oxygenation and carbon monoxide poisoning. Apropos of 100 cases). Moroc Med 50: 383-5, 1970. A 805
- GERHARDT T, GÖTHERT M, MALORNY G and WILKE H : Zur Frage der Toxizität von Kohlenoxid bei Atmung von CO-Luftgemischen unter erhöhtem Druck. (Carbon monoxide toxicity during the breathing of CO and air mixtures under hyperbaric pressure). Int Arch Arbeitsmed 28: 127-40, 1971. A 806
- GOULON M and HENNETIER G : Traitement de l'intoxication oxycarbonée par l'oxygène hyperbare. (Treatment of carbon monoxide poisoning by hyperbaric oxygen). Ann Chir Thorac Cardio 5: 619-23, 1966. A 807
- GOULON M, LEVY-ALCOVER M A, NOUAILHAT F and DORDAIN G : Étude de l'E. E. G. au cours de l'oxygénothérapie hyperbare. (Study of the EEG in the course of hyperbaric oxygen therapy). Rev Neurol Paris 117: 521, 1967. A 808
- GOULON M, BAROIS A, RAPIN M, NOUAILHAT F, GROSBUIS S and LABROUSSE J : Intoxication oxycarbonée et anoxie aiguë par inhalation de gaz de charbon et de hydrocarbures. (Carbon monoxide poisoning and acute anoxia due to inhalation of coal gas and hydrocarbons). Ann Med Interne Paris 5: 335-49, 1969. A 809
- GOULON M, LÉVY-ALCOVER M A, NOUAILHAT F and DORDAIN G : The EEG in hyperbaric oxygen therapy. Clin Neurophysiol 25: 90, 1968. A 810
- GRAMER L and BECKENKAMP H : Über die Abatmung von Kohlenmonoxyd bei Vergifteten unter Sauerstoffbeatmung. (On the expiration of carbon monoxide in poisoned persons treated with oxygen inhalation). Int Arch Gewerbepath 22: 282-96, 1966. A 811
- GRAZIANI G and PAGGIE : Trattamento dell'ossicarbonismo acuto sperimentale con ossigeno iperbarico. (Treatment of experimental acute carbon monoxide poisoning by hyperbaric oxygen). Folla Med Napoli 48: 1219-29, 1965. A 812
- HANQUET M and LAMY M : L'oxygénothérapie hyperbare en caisson monoplace à oxygène pur. (Hyperbaric oxygen therapy in a caisson using pure oxygen). Laval Med 42: 647-67, 1971. A 813
- HIRANO H, INOUE Y and TANAMI J : Studies on the mode of actions of carbon monoxide. 3rd report: The central action of carbon monoxide. Jap J Hyg 25: 488-93, 1971. A 814
- IWANOWSKA H : O możliwościach leczenia tlenem stosowanym pod zwiększonym ciśnieniem. (On the possibilities of treatment with hyperbaric oxygenation). Pol Tyg Lek 22: 1981-3, 1967. A 815
- JACOBSEN E : The theory and indications of hyperbaric oxygen. A review. Laval Med 42: 291-300, 1971. A 816
- KOKAME G M and SHULER S E : Carbon monoxide poisoning. Arch Surg 96: 211-5, 1968. A 817
- MAEROVICH I M :  
KONDRASHENKO V T, GLANTS B R, and / (Hyperbaric oxygen therapy of hypoxic states in acute cranio-cerebral trauma and acute exogenous psychoses). Zh Nevropatol Psikhiatr 71: 271-7, 1971. A 818
- KUCHER R and RIEDEL W : Die Behandlung des Gasbrandes in der Sauerstoffüberdruckkammer. (Treatment of gas gangrene in hyperbaric oxygenation chamber). Wien Klin Wochenschr 81: 308-10, 1969. A 819
- LAMY M and HANQUET M : L'oxygénothérapie hyperbare. Premières applications. (Hyperbaric oxygenotherapy. First applications) Acta Anaesth Belg 19: 46-82, 1968. A 820

1005051294

## Additional Bibliography List No. 18

- LAMY M and HANQUET M : Cinquante cas d'intoxication oxycarbonée traités par l'oxygénothérapie hyperbare. (Fifty cases of carbon monoxide poisoning treated by hyperbaric oxygenation). Acta Anaesthesiol Belg 20: 49-64, 1969. A 821
- LARCAN A, ROBERT J, CALAMAI M and FREJAVILLE J P : Le traitement de l'intoxication oxycarbonée par le caisson mobile de l'oxygénation hyperbare. Premiers résultats. (Treatment of carbon monoxide poisoning using a mobile hyperbaric oxygenation unit. 1st results) Presse Med 75: 1325-9, 1967. A 822
- LEDINGHAM I M: New approaches to poisoning. Hyperbaric oxygenation. Proc Roy Soc Med 57: 807-9, 1964. A 823
- LEE K : The use of portable oxygen sets in carbon monoxide poisoning. Trans Soc Occup Med 16: 85-6, 1966. A 824
- MANTZ J M and TEMPE J D : L'ossigenoterapia iperbarica. (Hyperbaric oxygen therapy). Minerva Med 59: 3137-49, 1968. A 825
- MANTZ J M and TEMPE J D : Die Sauerstoff-Uberdruckbehandlung. (Treatment with hyperbaric oxygen). Munch Med Wochensh 110: 2186-97, 1968. A 826
- MATTHEW H : Mobile pressure chamber. Brit Med J 600, 1970. A 827  
and BAYLISS G J A
- MILLER J N A: Clinical applications of hyperbaric oxygen therapy in Sydney - a review of recent cases. Med J Aust 1: 835-8, 1966. A 828
- MOLFINO F and ZANNINI D: Sulla cura dell'ossicarbonismo acuto mediante ossigenoterapia in camera pressurizzata. (On the treatment of acute carbon monoxide poisoning by oxygen therapy in a pressure chamber. Rass Med Industr 33: 334-6, 1964. A 829
- NORMAN J N, MacINTYRE J, SHEARER J R and SMITH G : Use of a one-man mobile pressure chamber in the treatment of carbon monoxide poisoning. Brit Med J 2: 333-4, 1970. A 830
- OHRESSER P, CHASSON J, JOUGLARD J, GOUIN F, DUBOULOZ F and TASSY J : Traitement hyperbare des intoxications oxycarbonées. (Hyperbaric treatment of carbon monoxide poisoning). Marseille Med 105: 555-7, 1968. A 831
- PETTY T L : Oxygen therapy. Ann Intern Med 71: 666-7, 1969. A 832
- ROCHE L, BERTOYE A, VINCENT P, MOTIN J, GARIN J P, BOLOT J F and CHADENSON O : Comparaison de deux groupes de vingt intoxications oxycarbonées traitées par oxygène normobare et hyperbare. (Comparison of 2 groups of 20 cases of carbon monoxide poisoning treated with normobaric and hyperbaric oxygen. Lyon Med 220: 1483-99, 1968. A 833
- SAVATEEV N V, TONKOPIJ V D and FROLOV S F : (Hyperbaric oxygenation in certain types of acute poisoning. Review of the literature). Voenn Med Zh 2: 23-8, 1970. A 834
- SCHULTE J H : The use of hyperbaric oxygen in clinical medicine. J Occup Med 11: 462-5, 1965. A 835
- SIMPSON B R and RITCHIE H D : Expérience personnelle du traitement à l'oxygène à haute pression. (Personal experience with high pressure oxygen therapy). Lyon Chir 64: 5-12, 1968. A 836
- SLUIJTER M E : The treatment of carbon monoxide poisoning by administration of oxygen at high atmospheric pressure. Progr Brain Res 24: 123-82, 1967. A 837
- SMITH G : Carbon monoxide poisoning. Ann NY Acad Sci 117: 684-7, 1965. A 838
- SUGIMOTO T and YASUMITSU T : (Hyperbaric oxygen therapy of carbon monoxide poisoning). Jap J Clin Med 27: 2157-64, 1969. A 839

1005051295



## Additional Bibliography List No. 18

- TAKEYA H, TAKANO M, TAMURA A, HOJO Y, YASUDA K, YOSIDA T and HURUKAWA Y :  
(Carbon monoxide poisoning). Jap J Anesth 19: 172-9, 1970. A 840
- TEMMERMAN P and ETIENNE A : Résultats de l'oxygénothérapie hyperbare dans l'intoxication  
oxycarbonée. (Results of hyperbaric oxygen therapy in carbon monoxide poisoning).  
Acta Anaesthesiol Belg 20: 119-41, 1969. A 841
- THURSTON J : Hyperbaric oxygen in carbon monoxide poisoning. Brit Med J 4: 386, 1968. A 842
- THURSTON J G B : Place of hyperbaric oxygen in intensive care. Proc R Soc Med 64: 1287-8,  
1971. A 843
- WERNITSCH W : Zur Technik und Anwendung der hyperbaren Oxygenation. (Technic and ad-  
ministration of hyperbaric oxygenation). Med Welt 49: 2668- , 1969. A 844
- WINTER A and SHATIN L : Hyperbaric oxygen in reversing carbon monoxide coma. Neurolo-  
gic and psychologic study. NY State J Med 70: 880-4, 1970. A 845
- ZORN H : Carbogen oder reiner Sauerstoff zur Beatmung bei Kohlenmonoxidvergiftung?  
(Carbogen or pure oxygen for artificial respiration in carbon monoxide poisoning?)  
Dtsch Med Wochenschr 93: 1536, 1968. A 846

1005051296

(Reprint number) followed by pages in the text.

- ABELSON 1967 (640) 114, 141  
ABELSON 1968 (A 240) 192  
ACHTEN, LEDOUX, CORBUSIER and  
THYS 1971 (615) 112, 111  
AFANSEV 1967 (392) 76, 74  
AINSWORTH, SCHLOEGEL, DOMANSKI and  
GOLDBAUM 1967 (A 1) 176  
AINSWORTH 1968 (A 55) 180  
AITKEN, BUGLASS and KREITMANN 1969  
(A 617) 216  
AITKEN, DALY, KREITMAN and  
PROUDFOOT 1966 (A 460) 206  
AJEMIAN and WHITMAN 1970 (A 318) 197  
ALBEN and CAUGHEY 1968 (A 56) 180  
ALEXANDRE 1967 (A 646) 218  
ALEKSANDROV, BESCHOCHLOW and  
DAVIDOV 1970 (A 686) 222  
ALFSEN, CHIANCONE, ANTONINI, WAKS and  
WYMAN 1970 (A 57) 180  
ALIVISATOS, BAZAS, ALEXOPOULOS and  
VERYKOKAKIS 1967 (168) 40, 38  
ALPERT and BANERJEE 1971 (A 58) 180  
ALVAPES, SCHILLING, LEVIN and  
KUNTZMAN 1967 (A 544) 211  
ALVARES, SCHILLING, LEVIN and  
KUNTZMAN 1971 (A 545) 211  
ALVIS 1967 (A 784) 229  
AMENDT and REDDEMANN 1970 (A 618) 216  
AMERICAN THORACIC SOCIETY 1966  
(A 241) 192  
AMYOT, GLARD and ROBERT 1967 (A 461)  
206  
ANDERHUB, HOFER and SCHERRER 1970  
(A 2) 176  
ANDERSEN and GIBSON 1971 (A 59) 180  
ANDERSON 1971 (A 270) 194  
ANDERSON 1967 (A 242) 192  
ANDERSON, REED and CHANCE 1970  
(A 60) 180  
ANDERSON, ALLENSWORTH and DeGROOT  
1967 (309) 64, 63  
ANDERSON and ANTONINI 1968 (A 3) 176  
ANDERSON 1970 (A 647) 218  
ANDERSON 1971 (271) 59, 56  
ANDERSSON 1972 (310) 64, 62  
ANDO 1966 (A 462) 206  
ANDO, SENO and HAGIWARA 1969 (498)  
94, 93  
ANON 1964 (A 463) 206  
ANON 1965 (641) 114, 142  
ANON 1966 (A 243) 192  
ANON 1968 (A 465) 206  
ANON 1968 (A 464) 206  
ANON 1968 (643) 114, 144  
ANON 1968 (642) 114, 143  
ANON 1969 (A 619) 216  
ANON 1969 (A 788) 229  
ANON 1970 (A 649) 218  
ANON 1970 (A 648) 218  
ANON 1970 (A 271) 194  
ANON 1971 (644) 114, 145  
ANON 1971 (645) 114, 146  
ANON 1971 (A 789) 229  
ANON 1972 (A 319) 197  
ANON 1972 (A 187) 189  
ANTONINI, ANDERSON and MAURIZIO  
1972 (A 65) 180  
ANTONINI, BRUNORI, WYMAN and Noble  
1966 (A 63) 180  
ANTONINI, BUCCI, FRONTICELLI, WYMAN  
and ROSSI-PANELLA 1965 (A 62) 180  
ANTONINI, CHIANCONE and BRUNORI 1967 (A 64) 180  
ANTONINI, SCHUSTER, BRUNORI and  
WYMAN 1965 (A 61) 180  
ANTOS and SEVCIK 1971 (393) 76, 74  
AOYAMA 1970 (A 272) 194  
APPLEBY 1969 (A 66) 180  
AQUINAS 1964 (A 729) 223  
ARAI 1969 (A 466) 206  
ARIN and WARNECK 1972 (A 151) 186  
ARMSTRONG 1922 (109) 30, 29  
ARNOTT, PETIT and CHABRIER 1964  
(A 467) 206  
ARONDEL, GAUBERTI and ROC'ET  
1964 (A 650) 218  
ARONOW, DENDINGER and ROKAW 1971  
(311) 64, 63; (647) 114, 147  
ARONOW, HARRIS, ISBELL, ROKAW and  
IMPARATO 1972 (312) 64, 62;  
(649) 114, 147  
ARONOW, KAPLAN and JACOB 1938 (646)  
114, 147  
ARONOW and ROKAW 1971 (313) 64, 64;  
(648) 114, 147  
ARTURSON, CARBY, ROBERT and ZAAR  
1972 (35) 17, 15; (142) 34, 34  
ASAI and TORU 1969 (A 468) 206  
ASORBE DOMINGUEZ 1969 (A 790) 229  
ASMUSSEN and VINTHER-PAULSEN 1953  
(272) 59, 57  
ASMUSSEN and VINTHER-PAULSEN 1949  
(394) 76, 75  
ASTRUP 1964 (360) 72, 69  
ASTRUP 1966A (361) 72, 69  
ASTRUP 1966B (362) 72, 69  
ASTRUP 1967 (363) 72, 70  
ASTRUP 1969 (364) 72, 70  
ASTRUP 1970 (A 67) 180  
ASTRUP 1970 (365) 72, 70  
ASTRUP 1972 (366) 72, 70  
ASTRUP 1966 (367) 72, 66  
ASTRUP 1967 (368) 70, 70  
ASTRUP, KJELDSSEN and WANSTRUP 1970A  
(369) 72, 70  
ASTRUP, KJELDSSEN and WANSTRUP 1970B  
(370) 72, 70

1005051297

- ASTRUP, KJELDSEN and SIGAARD-ANDERSEN 1971 (479) 91, 89  
 ASTRUP, PAULI, KJELDSEN and PETERSEN 1968 (200) 46, 45  
 ASTRUP, TROMME, OLSEN and KJELDSEN 1972 (563) 105, 103, 104  
 AUBEAU, LEROY and CHAMPEIX 1965 (A 320) 197  
 AYRES 1972 (652) 114, 149  
 AYRES and BUEHLER 1970 (169) 40, 38  
 AYRES, BUEHLER and ARMSTRONG 1964 (A 36) 200  
 AYRES, CRISCITIELLO and GIANNELLI 1956 (A 4) 176  
 AYRES, EVANS and BUCHLER 1972 (A 245) 192  
 AYRES, GIANNELLI and ARMSTRONG 1965 (A 36) 17, 15; (143) 34, 33; (193) 44, 42  
 AYRES, GIANNELLI and MUELLER 1970 (A 314) 64, 63; (650) 114, 148  
 AYRES, MUELLER, GREGORY, GIANNELLI and PENNY 1969 (315) 64, 63; (651) 114, 148  
 BACK 1959 (499) 94, 93  
 BADAL 1964 (A 469) 206  
 BADEN 1970 (616) 112, 111  
 BAKER and TUMASONIS 1971 (A 582) 214  
 BAKER and TUMASONIS 1972 (A 583) 214  
 BAKER, FISHER, MASEMORE and SOPHER 1972 (A 273) 194  
 BALBO, MARUCCI and RONCHI 1966 (37) 17, 13, 15; (144) 34, 33  
 BANERJEE, DOUZOU and LOMBARD 1968 (A 68) 180  
 BANKL and JELLINGER 1967 (564) 105, 102  
 BANYAI 1970 (653) 115, 151  
 BARACH, ECKMAN and MOLOMUT 1941 (66) 22, 20, 21  
 BARBOSA and PETERS 1971 (A 584) 214  
 BARIBAUD, YACOB, FAURE, MALINAS and CAU 1970 (565) 105, 103  
 BARRETT, BENNETT and BUCKMASTER (A 321) 197  
 BARRIOS, KOLL and MALORN 1969 (500) 94, 93  
 BARTEK, GAUME and ROSTAMI 1970 (A 274) 194  
 BARTH, 1970 (A 275) 194  
 BARTHE, PARIS, DUCHEMIN and THOMAS 1953 (145) 34, 33; (38) 17, 14  
 BARTHELEMY, PARC, MICHAUD and MATHE 1967 (A 791) 229  
 BARTLETT 1968 (654) 115, 153  
 BASMADZHIEVA, KURCHATOWA, DAVYDKOVA and TSWETANOV 1968 (A 688) 222  
 BATES, CHRISTIE and VARVIS 1960 (A 362) 200  
 BAUER 1965 (A 5) 176  
 BAUMBERGER 1923 (110) 30, 29  
 BAXTER and HOBBS 1967 (111) 30, 28  
 BEAN 1968 (A 620) 216  
 BEARD 1969 (1) 9, 6  
 BEARD and GRANDSTAFF 1970 (480) 91, 88  
 BEARD and WERTHEIM 1967 (501) 94, 93  
 BEARON 1965 (A 470) 206  
 BEAUDOING, GACHON, BUTIN and BOST 1969 (566) 105, 102  
 BECK and SUTER 1938 (316) 64, 63  
 BEDELL and OSTIGUY 1967 (A 363) 200  
 BECKMANS 1967 (A 6) 176  
 BENDEL and VANCE 1965 (A 364) 200  
 BEGHE 1964 (A 730) 225  
 BEHRMAN, FISHER and PATON 1971 (567) 105  
 BELAISCH 1954 (A 731) 225  
 BELLI and GUILLANO 1955 (67) 22, 20  
 BELYAEV 1967 (a 546) 211  
 BENDER, GOTHERT, MALORNEY and SEBBESSE 1971 (481) 91, 89  
 BENESCH, GIBSON and BENESCH 1964 (A 69) 180  
 BENESCH, MAEDAN and BENESCH 1972 (A 70) 180  
 BENSON and GREENBERG 1969 (438) 83, 81  
 BEREZIN and PUZARES 1969 (A 792) 229  
 BERNHARD and FILLER 1968 (A 793) 229  
 BERTIN, FRANCOIS, PEQUIGNOT and SOULAIRAC 1970 (A 651) 218  
 BERTONE 1965 (A 471) 206  
 BESZNYAK 1967 (A 794) 229  
 BETHUILL and DELAHAYE-POLOUVIER 1967 (A 652) 218  
 BETHLENFALVAY 1971a (395) 76, 74  
 BETHLENFALVAY 1971b (396) 76, 74  
 BETKE 1968 (A 72) 181  
 BETKE and SHEPARD 1968 (A 71) 180  
 BEUMER 1964 (A 365) 200  
 BEUMER 1965 (A 366) 200  
 BHATNAGAR 1970 (A 547) 211  
 BROWN 1969 (39) 17, 15  
 BIDE and COLLIER 1964 (A 73) 181  
 BILCHIK, MULLER-BERGH and FRESHMAN 1971 (439) 83, 81  
 BILS and ROMANOVSKY 1967 (207) 49, 48  
 BIMONTE, PORTOLANO and TUFANO 1971 (A 795) 229  
 BINET and BURSTEIN 1948 (333) 67, 66  
 BINET and BETOURNE 1951 (273) 59, 57  
 BINNENBRUCK, HAUSEN, RUNOW and WERHEIT 1970 (A 591) 214  
 BIRNSTINGL, COLE and HAWKINS 1966 (371) 72, 69  
 BIRNSTINGL, COLE and HAWKINS 1967 (398) 76, 75  
 BIRNSINGL, BRINSON and CHAKRABARTI 1971 (397) 76, 74  
 BJURE 1965 (A 367) 200  
 BJURE and FALLSTROM 1963 (568) 105  
 BJURE and NILSSON 1965 (A 7) 176  
 BLACKMORE 1970 (399) 76  
 BLACKMORE 1970 (A 8) 176  
 BLACKMORE 1970 (A 165) 187  
 BLANC, HUYNH and ESPAGNO 1967 (A 322) 197  
 BLOOM 1972 (A 166) 187  
 BLUMER 1970 (A 276) 194

1005051298

BOCCADORO and INGIULLA  
1968 (A 277) 194  
BOCCALETTI, NOFRINI, JOSI and MAGGIO  
1966 (A 796) 229  
BOECK 1958 (170) 40, 38  
BOGDAN and JUCHAU 1970 (A 548) 211  
BOKHOVEN and NIESSEN 1961 (112) 30, 28  
BOLLINELLI, ROUCH, PUJOL, CARRIERE  
and CARLES 1971 (A 368) 200  
BONNET, GRATADOU, BONNET and LECIAK  
1967 (A 472) 206  
BOOZ 1969 (A 369) 200  
BORBELY 1965 (A 732) 225  
BORST 1967 (275) 59, 57  
BOTTEAU and MOUSSION 1967 (A 323) 197  
BOUHUYS, GEORG, JONSSON, LUNDIN and  
LINDELL 1960 (A 370) 200  
BOULETTEAU and MOTIN 1970 (A 733) 225  
BOULEY, GODIN, ROUSSEL and GIRARD  
1971 (A 585) 214  
BOUR 1964 (A 654) 218  
BOUR, GUY-GRAND, TUTIN and TAMINIAUX  
1967 (A 473) 206  
BOUR, GUY-GRAND, ROGER, TUTIN and DORF  
1968 (599) 110, 108  
BOUR, PASQUIER and BERTRAND-HARDY  
1966 (A 734) 225  
BOUR, TUTIN and PASQUIER 1967 (A 474) 206  
BOVE and SIEBENBERG 1970 (A 278) 194  
BOWDEN and WOODHALL 1964 (66) 22, 21;  
(146) 34, 33  
BOZEK, PAJOR and WASOWICZ 1965 (A 653) 218  
BRAJA and TROMPEO 1964 (201) 46, 45  
BRANDENBERGER 1967 (A 621) 216  
BRANDON 1970 (A 735) 225  
BRANDT 1965 (A 279) 194  
BRAUSER, VERSMOLD and BUCHER 1968 (A 549)  
211  
BREITNECKER 1938 (429) 80, 79  
BRETON, CAROFF, MARTIN, DEHOUE and  
DEHOUE 1969 (430) 80, 79  
BRETON, GARAT and DEROBERT 1969  
(A 34)  
BREU 1942 (230) 53, 51  
BREU 1942 (274) 59, 57  
BREU 1943 (231) 53, 51  
BREWER, EATON, GROVER and WEIL  
1971 (41) 17, 15  
BREWER, EATON, WEIL and GROVER 1970  
(40) 17, 15  
BREWER 1937 (334) 67, 66  
BREYSSE, BOVEF and GABAY 1966 (A 9) 176  
BREYSSE and BOVEE 1969 (171) 33, 39  
BRICE and ROESLER 1966 (A 280) 194  
BRIDGE and CORN 1972 (113) 30, 28  
BRIGATTI, PARIGI and VARETTO 1964 (A 689)  
222  
BROBERG and SMITH 1967 (A 586) 214  
BRODY and COBURN 1969 (276) 59, 56;  
(400) 76, 74  
BRODY and COBURN 1970 (277) 59, 56  
BRUNNER 1939 (232) 53, 52  
BRUNORI 1966 (A 74) 181  
BRUNORI, ANTONINI, WYMAN, TENTORI  
VIVALDI and CARTA 1968 (A 75) 181  
BRUNORI, BONAVENTURA, BONAVENTURA  
and WYMAN 1972 (A 76) 181  
BRUSADELLI, PALMA and DIPRETORO 1968  
(A 797) 229  
BUCHWALD 1969 (172) 40, 38, 39  
BUCHWALD 1969 (A 10) 176  
BUCKLEY and FEAR 1964 (A 690) 222  
BULTERLIS 1965 (A 798) 229  
BUNCHER 1969 (569) 105, 103  
BURCK and PORTWICH 1964 (551) 101, 99  
BURG and DOUGLASS 1969 (A 655) 218  
BURGESS, GILLESPIE, GRAF and NADEL  
1968 (A 371) 200  
BURMEISTER, BARCKOW, HUMPERT, IBE  
and LERCHE 1968 (A 736) 225  
BURMEISTER and HEUHAS 1970 (A 737) 225  
BURSTON 1969 (A 622) 216 (A 799) 229  
BURVILL 1970 (623) 216  
BYSTROM 1970 (281) 194  
CACCURI 1955 (278)  
CAILAR, SERRE, ROQUEFFEUIL, LEFEBVRE  
and MALZAC 1968 (A 800) 229  
CAIRNS and DENHARDT 1968 (A 587) 214  
CALLIGARI 1968 (A 656) 218  
CAMMA 1967 (655) 115, 154  
CAMPBELL 1936 (208) 49, 48  
CAMPBELL 1968 (A 282) 194  
CANDURA and CRAVERIA 1964 (401) 76, 74  
CANEPÀ, CAVALLO and MUZIO 1968  
(A 372) 200  
CAPELLARO 1964 (A 691) 222  
CAPELLARO and BRAGUZZI 1964 (A 692)  
222  
CAPELLARO and GANDOLFO 1964 (233)  
53, 52  
CARDING 1968 (502) 94, 93  
CARNOW 1971 (A 246) 192  
CAROFF, DEHOUE and DEROBERT 1970  
(279) 59, 57  
CARUSO and BARNABA 1968 (A 738) 225  
CASARETT 1971 (2) 9  
CASCINI and GAZZERIO 1966 (A 693) 222  
CASTELLINO 1955 (69) 22, 20  
CASULA, NISSARDI, SANNA-RANDACCIO  
and FRAU 1969 (373) 200  
CASULA, NISSARDI, SANNA-RANDACCIO  
and FRAU 1969 (A 374) 200  
CAUGHEY 1970 (A 78) 181  
CAUGHEY, ALBEN, MCCOY, BOYER,  
CHARACHE and HATHAWAY 1969  
(A 77) 181  
CELEGIN, HANSSON and SUNDSTROM  
1971 (A 324) 197  
CENTI and ZAFFIRI 1971 (A 73) 225  
CHANCE, ERECINSKA and WAGNER 1970  
(A 550) 211  
CHEAH 1970 (A 79) 181  
CHERKAVSKIIR 1970 (A 657) 218  
CHEVALIER, KRUMHOLZ and ROSS 1963  
(280) 59, 56

1005051299

CHEVAMER, KRUMHOLZ and ROSS 1963 (280) 59, 56  
 CHEVALIER, KRUMHOLZ and ROSS 1966 (191) 44, 43  
 CHEW, HANSON and SLACK 1969 (801) 229  
 CHICHET-GOSSET, CHASSON, OIRESSER, ARNAUD and DUBOULOZ 1967 (A 802) 229  
 CHICHKALDO, BENEVELSKY and MINSKY 1966 (A 283) 194  
 CHINET, MICHEL and HAAB 1971 (A 375) 201  
 CHIODI, DILL, CONSOLAZIO and HORVATH 1941 (195) 44, 42  
 CHOSY, GEE and RANKIN 1963 (196) 44, 43  
 CHOVIN 1967 (173) 40, 38, 39  
 CHRISTIANSEN and MAGID 1970 (A 80) 181  
 CHUDZIKIEWICZ 1957 (335) 67, 66  
 CHUKHRIENKO and LULKO 1968 (A 740) 225  
 CINKOTAL and THOMSON 1966 (A 376) 201  
 CIOCATTO and PATTONO 1964 (A 741) 225  
 CIOCATTO, PATTONO, QUERCI and ZAFFIRI 1966 (A 803) 229  
 CIS and PERANI 1964 (458) 86, 85  
 CIUHANDU, DIACONOVICI, KISS and RUSU 1964 (325) 197  
 CIUHANDU, DIACONOVICI, KISS and RUSU 1968 (377) 201  
 CIUHANDU and RUSU 1968 (A 326) 197  
 CIUHANDU, RUSU, DIACONOVICI and KISS 1966 (A 11) 176  
 CLARK and BUCKINGHAM 1971 (A 327) 197  
 CLAUZEL, TRINQUET, CARRE and MEYER 1966 (197) 44, 43  
 CLAYTON 1969 (A 247) 192  
 COBURN 1967 (A 206) 190  
 COBURN 1970 (3) 9, 7= (A 154) 186  
 COBURN 1970 (A 208) 190  
 COBURN 1970 (A 209) 190  
 COBURN, FORSTER and KANE 1965 (A 152) 186  
 COBURN and MAYERS 1971 (617) 112  
 COBURN, SWERDLOW, LUGMANMAKI, FORSTER and POWELL 1968 (A 153) 186  
 COBURN, WALLACE and ABOUD 1971 (210) 190  
 COBURN, WILLIAMS and KAHN 1966 (A 205) 190  
 COBURN, WILLIAMS, WHITE and KAHN 1967 (A 207) 190  
 COHEN, DEANE and GOLDSMITH 1969 (317) 64, 61  
 COHEN, DORION, GOLDSMITH and PERMUTT 1971 (174) 40, 38, 39  
 COHEN, PERKINS, URY and GOLDSMITH 1971 (70) 22, 19  
 COLE, HAWKINS and ROBERTS 1972 (570) 105, 104  
 COLLINS and GOULDING 1971 (A 742) 225  
 COLLISON, RODKEY and O'NEAL 1968 (A 12) 176  
 COLTMAN and DUDLEY 1969 (A 212) 190  
 COLTMAN, DUDLEY and LEVERETT 1969 (211) 190  
 COMMINS and LAWTHOR 1965 (A 13) 176  
 COMSTOCK, SHAH, MAYER and ABBEY 1971 (571) 105, 102  
 CONKLE, MABSON, ADAMS, ZEFT and WELCH 1967 (A 167) 187  
 CONNEY, LEVIN, IKEDA and KUNTZMAN 1968 (A 551) 211  
 CONROY 1969 (656) 115, 155  
 COOPER 1966 (4) 9, 8  
 COOPER, SCHLEYER and ROSENTHAL 1970 (A 552) 211  
 CORET and HUGHES 1964 (336) 67, 66  
 COSBY and BERGERON 1963 (234) 53, 52  
 COSCIA, PERRELLI, GAIDO and CAPELLARO 1964 (402) 76, 74  
 COSIC, ARSENIJEVIC, KANDIC and GRBESA 1964 (A 694)  
 COTES, DABBS, EVANS and HOLLAND 1972 (A 378) 201  
 CRAMOND 1968 (A 475) 206  
 CROSETTI, PETTINATI and RUBINO 1965 (A 658) 218  
 CROSETTI, RUBINO and PETTINATI 1966 (71) 22, 20; (A 81) 181  
 CUCHE and BERNARD 1969 (A 659) 218  
 CULVERWELL 1915 (114) 30, 27  
 CURPHEY, HOOD and PERKINS 1965 (42) 17, 13, 15; (147) 34, 33  
 CURPHEY 1968 (657) 115, 156  
 DAHMSTROM, NORDSTROM-OHRBERG and ROTHCHILD 1958 (43) 17, 14; (148) 34, 33  
 DALGAARD 1965 (A 660) 218  
 DALHAMN, EDFORD and RYLANDER 1968 (11t) 30, 28  
 DALLE, TOURNAIRE, BRUDIEUX and DELOST 1971 (A 381) 201  
 DALY 1969 (A 380) 201  
 DALY and WALDHIAUSEN 1967 (A 379) 201  
 DAMIA, SIBILLA and DAMIA 1965 (A 743) 225  
 D'AMORE, GIORDANO and PENATI 1968 (A 476) 206  
 DANTO 1964 (618) 112, 111  
 D'ARCA, GUALDI and ARCIERI 1964 (202) 46, 45  
 DATSENKO 1964 (A 588) 214  
 DATSENKO 1965 (A 14) 176  
 DATSENKO 1965 (A 695) 222  
 DATSENKO 1966 (235) 53, 52  
 DATSENKO, DOTSENKO, MARTYNIUK and PALCHEVSKY 1965 (A 553) 211  
 DATSENKO, DOTSENKO 1967 (A 554) 211  
 DAVID 1971 (624) 216  
 DAVIES, JONES and WARNER 1965 (A 328) 197  
 DEANE, GOLDSMITH and TUMA 1965 (A 696) 222

1005051300

DeBIAS, BIRKHEAD, BANERJEE, KAZAL,  
HOLBURN, GREENE, HARPER, ROSENFELD  
MENDUKE, WILLIAMS and FRIEDMAN  
1972 (318) 64, 62

DeBRUIN 1967 (175) 40, 38

DeBRUIN and HAERINGEN 1965 (176)  
40, 38, 39

DeBRUIN, VROEGE and VAN HAERINGEN  
1965 (177) 40, 38

DEEN 1969 (804) 230

DeGRAFF, TAYLOR, ORD, CHUANG and  
JOHNSON 1965 (A 382) 201

DELIVORIA-PAPADOPOULOS and COBURN  
1972 (572) 105, 102

DELWICHE 1970 (A 188) 189

DEMANGE and AUZAS 1969 (482) 91, 90

DESBAUMES 1968 (A 698) 222

DESOILLE 1967 (178) 40, 38

DESOILLE, CASTILLON, du PERRON, CREMER  
and LEBBE 1963 (A 213) 190

DESOILLE, CREMER and GIRARD 1965  
(A 214) 190

DETTORI and SCANSETTI 1965 (A 699) 222

DeVALOIS and SCHADE 1967 (529) 97, 96

DEVIATKA 1956 (337) 67, 66

DIAMANT-BERGER, GAJDOS, RAPIN and  
GOULON 1970 (281) 59, 56

DILLE and MOHLER 1969 (A 168) 187

DIMIZIC, FELICI and BIETTI 1969 (A 477) 206;  
(A 661) 218

DINMAN 1968 (5) 9, 7

DINMAN 1969 (319) 64, 61

DINMAN 1970 (658) 115, 160

DINMAN, EATON and BREWER 1970 (A 82)  
181

DIXON 1927a (116) 30, 27

DIXON 1927b (117) 30, 27

DOBOSZ and LUCZYWEK 1971 (A 478) 206

DOMINGUEZ, HALSTEAD and DOMANSKI  
1964 (A 15) 176

DONATELLI 1940 (236) 53, 52

DONTENWILL 1967 (210) 49, 48

DONTENWILL 1970 (211) 49, 48

DONTENWILL, RECKZEH and STADLER  
1966 (72) 22, 20; (209) 49, 48

DONTENWILL, RECKZEH and STADLER 1967  
(73) 22, 20

DORSCH and KOSTER 1965 (a 16) 176

DOUGLAS 1967 (A 189) 189

DOUMER and MERLEN 1946 (237) 53, 52

DOUZE 1971 (A 662) 218

DOUZE, HEYST, KREUKHET, LEEUW and  
HAMELINK 1967 (A 663) 219

DOYLE 1969 (659) 115, 162

DRABKIN 1970 (A 215) 190

DRISCOLL, DEUBER, BAETTIG and  
GRANDJEAN 1972 (74) 22, 20

DROGICHIINA and RYZHIKOVA 1967 (A 479)  
207

DUBLIN 1972 (94) 26, 25

DUBOIS 1970 (6) 9, 7

DUBOIS and MONKMAN 1972 (A 330) 197

DUBOIS, ZDROJEWSKI and MONKMAN  
1966 (A 329) 197

DuCAILLAR, LEFEBVRE, ROQUEFEUIL, MALZAC  
and PROST 1970 (A 805) 230

DUCROS 1968 (A 331) 197

DUGNAT 1965 (503) 94, 93

DUKE, GREEN and NEIL 1952 (212) 49, 47

DUKE and KILLICK 1952 (213) 49, 47

DUNLAP 1961 (A 284) 194

DUPLAY, ZIEGLER, GARDO and PINTO 1967  
(403) 76, 74

DVORAK, PROKSAN and ZITKA 1951 (238) 53, 52

EBERSOLD 1958 (A 169) 187

ECKARDT, MacFARLAND, ALARIE and

BUSEY 1972 (214) 49, 48

EFFENBERGER 1967 (A 17) 176

EHRICH, BELLET and LEWEY 1944 (239)  
53, 52

EHRISMANN and ABEL 1934 (118) 30, 29

EISEN and HAMMOND 1956 (404) 76, 75

EISENBUD and EHRLICH 1972 (A 248) 192

EL-ATTAR 1968A (405) 76, 74

EL-ATTAR 1968B (406) 76, 74

EL-EBRASHY, EL-ASHMAWY and ALY  
1967 (407) 76, 74

ELFIMOVA and KHUCHIATURYAN 1968  
(A 249) 192

ELIOT and BRA TT 1969 (408) 77, 75

ELLIS and SEATONBERRY 1966 (A 170) 187

ENGEL, RODKEY, O'NEAL and COLLISON  
1969 (A 83) 181

ENGEL, RODKEY and KRILL 1971 (A 18)  
177

ENVIRONMENTAL HEALTH SERVICE 1970  
(7) 9, 7

ENVIRONMENTAL PROTECTION AGENCY  
1971 (A 250) 192

EPSTEIN 1969 (A 251) 192

ERBEN 1967 (A 19) 177

ESTABROOK, FRANKLIN and HILDEBRANDT  
1970 (A 555) 211

ESTLER 1965 (A 744) 226

ESTLER, AMMON and HEIM 1971A (605)  
94, 93

ESTLER, AMMON and ZIMMERMANN  
1969 (504) 94, 93

ESTLER, HEIM, AMMON and ZIMMERMANN  
1971B (506) 94, 93

FABRE, TRUIAUT and BERROD 1951  
(75) 22, 21

FAIVRE, DUREAX, VINCENT and MULLER  
1954 (240) 53, 52

FAIVRE, GILGENKRANTZ and HUEBER 1959  
(241) 53, 52

FALCONER and MOLLER 1971 (A 20) 177

FALLSTROM 1968 (A 218) 190

FALLSTROM 1968 (A 219) 190

FALLSTROM 1969 (A 220) 190

FALLSTROM and BJURE 1967 (A 216) 190

FALLSTROM and BJURE 1968 (A 217) 190

FARBEROW and SIMON 1969 (A 625) 216

FATINI and GALLENGA 1968 (A 626) 216

1005051301

FAIRE, VINCENT, ESCHAPASSE, CASTAING,  
LOISEAU and CHEVAIS 1965 (530) 97, 96  
FAURE, VINCENT, ESCHAPASSE, LOISEAU  
and CASTAING 1965 (A 480) 207  
FAZEKAS 1967 (507) 94  
FELDMAN and LAMPERT 1968 (A 664) 219  
FELDSTEIN 1965 (A 21) 177  
FELDSTEIN 1965 (A 332) 197  
FELDSTEIN 1967 (A333) 198  
FELDSTEIN 1969 (A 252) 192  
FENN 1970 (A 155) 186  
FENN 1971 (A22) 177  
FIANDACA and VERCELLOTTI 1964 (179)  
40, 37  
FINCK 1966 (8) 9, 6  
FIRST and MURPHY 1970 (A 700) 222  
FISHER, HYDE, BAUE, REIF and KELLY 1969  
(198) 44, 42; (215) 49, 48  
FISHER 1968 (A 665) 219  
FLAXMAN 1939 (282) 59, 56  
FLEIG 1908 (A 589) 214  
FLETCHER 1972 (A 383) 201  
FODOR 1969 (431) 80, 79  
FODOR, MALORNY and COLMANT 1964  
(531) 97, 96  
FORBES 1970 (A 385) 201  
FORBES, SARGENT and ROUGHTON 1945  
(A 384) 201  
FORMAN and FEIGELSON 1971 (A 84) 181  
FORSTER 1970 (A 156) 186  
FORSTER, ROUGHTON, CANDER, BRISCOE  
and KREUZER 1957 (A 386) 201  
FORTUNATO and CATALANO 1970 (459) 86, 85  
FOURNIER 1967 (A 745) 226  
FRANCHINI, CANALE and CELESTRI  
1967 (A 23) 177  
FRANCISCO and SILVEY 1971 (A 590) 214  
FRANCOIS and BERTIN 1964 (A 627) 216  
FREIGANG, SEIDEL and FLACH 1968 (460)  
86, 85  
FREIREICH and LANDAU 1971 (A 24) 177  
FREYSSCHUSS and HOLMGREN 1965 (A 387)  
201  
FRIBERG, NYSTROM and SWANBERG 1959  
(573) 105, 102  
FRITSCH 1969 (461) 86, 85  
FRONING, MATHER, DADDARIO and  
HARTUNG 1969 (A 592) 214  
FRYZE, GRUSZKA and ZAWADZKI 1970 (A 666)  
219  
FUKUI and KAKIUCHI 1970 (A 85) 181  
GABRIEL 1969 (A 388) 201  
GAENSLE, CADICAN, ELLICOTT, JONES  
and MARKS 1957 (44) 17, 14; (149) 34, 33  
GARLAND and PEARCE 1967 (A 481) 207  
GARREL, PERRET, PELLAT and ARNOULD  
1970 (A 483) 207  
GARREL, PERRET, PELLAT and ARNOULD  
1970 (A 482) 207  
GASSMAN and WHANNE 1967 (A 25) 177  
GATTO, CANEPA, CAVALLO and MASSIMILIA  
1967 (A 389) 201

GAULTIER, FOURNIER, GERVAIS and  
BODIN 1964 (A 484) 207  
GAULTIER, FOURNIER, GERVAIS,  
EFTHYMIU, SRAER, DISMUTH,  
BODIN, CHRISTOFOROV, SICOT and  
FREJAVILLE 1968 (A 746) 226  
GAULTIER, FREJAVILLE, BISMUTH and  
PEBAY-PEYROULA 1970 (A 628) 216  
GAUME, BARTEK and ROSTAMI 1971 (508)  
94, 93  
GEDDES and STEINHARDT 1968 (A 86)  
181  
GEIER, TUTTIN, PASQUIER, NAJMAN  
and DOOR 1966 (532) 97, 96  
GEMZELL, ROBBE and STROM 1958  
(574) 105, 102  
GEORGE and SCHEJTER 1964 (A 87) 181  
GERBER and SEEWALD 1966 (A 747) 226  
GERHARDT, GOTHERT, MALORNY and  
WILKE 1970 (A 390) 201  
GERHARDT, GOTHERT, MALORNY and  
WILKE 1971 (A 806) 230  
GERITSEN 1964 (A 701) 222  
GETTLER and MATTICE 1933 (45) 17  
11, 14  
GIBBONS and MITROPOULOS 1972 (372)  
72, 70  
GIBSON, HELLER and YAKULIS 1966  
(A 89) 181  
GIBSON and KAMEN 1966 (A 90) 181  
GIBSON, PALMER and WHARTON 1965  
(A 88) 181  
GIBSON and PARKHURST 1968 (A 91) 182  
GIEL 1965 (660) 115, 163  
GIEVER 1967 (9) 9, 6  
GIEVER and RUCH 1971 (A2 53) 192  
GILL 1971 (A 667) 219  
GIRARDI, CIS and PLATTI 1967 (A 485) 207  
GIROUD 1967 (575) 105, 102  
GLADYSHEVSKAYA, DOLOSHITSKY and  
SOBCHUK 1966 (619) 112, 111  
GLASEL 1970 (A 748) 226  
GLASS, MALLACH and WOJAHN 1966  
(A668) 219  
GLASS, EDWARDS, DeGARRETA and  
CLARK 1969 (A 92) 182  
GLASS, GARRETA, LEWIS, GRAMMATICOS  
and SZUR 1968 (409) 77, 74  
GLASS, JACOBY, WESTERMAN, CLARK,  
ARNOT and DIXON 1968 (576) 105, 102  
GLOWACKI, GRUDZINSKA and WACLAWIK 1958  
(432) 80, 79  
GOKIN 1971 (283) 59, 57  
GOLACKA, JANIK-KURYLCIO and  
ROZEK 1969 (A 669) 219  
GOLDBERG and CHAPPELL 1967 (509) 94, 93  
GOLDSMITH 1964 (10) 9, 6  
GOLDSMITH 1967 (A 254) 192  
GOLDSMITH 1967 (A 670) 219  
GOLDSMITH 1969 (661) 115, 164  
GOLDSMITH 1970 (320) 64, 61  
GOLDSMITH 1972 (662) 115, 165

1005051302

- GOLDSMITH and COHEN 1969 (11) 9, 7  
 GOLDSMITH and DEANE 1965 (A 286) 194  
 GOLDSMITH and LANDAW 1968 (12) 9, 6  
 GOLDSMITH and ROGERS 1959 (285) 194  
 GOLDSMITH, SCHUETTE and NOVICK 1963  
 (46) 17, 13, 14; (150) 34, 33  
 GOLDSMITH and TERZAGHI 1963 (76) 22, 21;  
 (151) 34, 33  
 GOLDSMITH, TERZAGHI and HACKNEY 1963  
 (180) 40, 38  
 GOLDSTEIN 1965 (577) 105, 102  
 GOLDSTEIN and EPSTEIN 1972 (663) 115, 166  
 GORALSKI and JANUSZKO 1968 (A 486) 207  
 GORDON 1965 (A 487) 207  
 GORODINSKY, LEVINSKY and SCHERBAKOV  
 1967 (A 334) 198  
 GORSKI 1962 (284) 57, 59  
 GOTHE, FRISTEDT, SUNDELL, KOLMODIN  
 EHKNER-SAMUEL and GOTHE 1969  
 (181) 40, 38, 39  
 GOTHERT, LUTZ and MALORNY 1970 (338)  
 67, 66  
 GOTO 1968 (A 749) 226  
 GOULDING 1965 (A 629) 216  
 GOULDING 1967 (A 750) 226  
 GOULON, BAROIS, RAPIN, NOUAILHAT,  
 AUGUSTIN, HENNETIER, BAGUET,  
 KUNTZIGER and BRETEAU 1965 (A 751) 226  
 GOULON, BAROIS, RAPIN, NOUAILHAT,  
 GROSBUIS and LARROUSSE 1969 (A 809) 230  
 GOULON, BAROIS, GAJDOS, LABROUSSE,  
 SCHORTGEN, AMERONGEN and ROBERT  
 1970 (A 752) 226  
 GOULON and HENNETIER 1966 (A 807) 230  
 GOULON, LEVY-ALCOVER, NOUAILHAT  
 and DORDAIN 1967 (A 808) 230  
 GOULON, LEVY-ALCOVER, NOUAILHAT and  
 DORDAIN 1968 (A 810) 230  
 GRAHAM and HITCHENS 1968 (A 630) 216  
 GRAMER and BECKENKAMP 1966 (A 811) 230  
 GRAMER and ROUF 1968 (410) 77, 74  
 GRAZIANI and PAGGI 1965 (A 812) 230  
 GRAZIANI, ROSSI, CASTELLINO and  
 SILVERSTRON 1957 (243) 53, 52  
 GRAY and GIBSON 1970 (A 93) 182  
 GRAY and GIBSON 1971 (A 94) 182  
 GRAY and GIBSON 1971 (A 95) 182  
 GRAYBIEL 1942 (242) 53, 51  
 GREGORY, MALINOSKI and SHARP 1969  
 (A 631) 216  
 GREMY, SALMON, FRANCOIS and BERTIN  
 1968 (A 671) 219  
 GRIFFIN and HOLLOCHER 1967 (A 96) 182  
 GRIFFITHS 1970 (A 672) 219  
 GROB 1968 (119) 30, 29  
 GROHME, SCHNEIDER and MASSHOFF 1969  
 (533) 97, 96; (A 488) 207  
 GROSSE and NEUHAUS 1970 (552) 101, 99  
 GRUNDY 1969 (A 255) 192  
 GRUT, ASTRUP, CHALLEN and GERHARDSSON  
 1970 (13) 9, 7  
 GUELIN 1969 (A 702) 223  
 GUEST, DUNCAN and LAWTHIER 1970  
 (462) 86, 85  
 GUEYE, BA and DIOP 1965 (A 673)  
 219  
 GUIDOTTI and KONIGSBERG 1964 (A 97) 182  
 GUILLERM, BADRE and GAUTIER 1967  
 (285) 59, 56  
 GULERIA, PANDE, SEITHI and ROY 1971  
 (A 391) 201  
 GULZOW 1957 (A 392) 201  
 GUNTHER 1971 (A 489) 207  
 GUTENKAUF, BRATT and ELIOT 1967  
 77, 75  
 GUY, SALHANY and ELIOT 1971 (412)  
 77, 75  
 GUYATT, NEWMAN, CINKOTAL, PALMER  
 and THOMSON 1965 (A 393) 202  
 GYDELL 1966 (A 26) 177  
 HAAB and PIPER 1968 (A 394) 202  
 HAAGEN-SMIT 1966 (A 287) 194  
 HADDON, NESBITT and GARCIA 1961  
 (578) 105, 104  
 HAEBISCH 1970 (120) 30, 28  
 HAGGARD and GREENBERG 1934 (601) 110,  
 109  
 HAHN and COPELAND 1966 (A 593) 214  
 HAKIM 1970 (580) 106, 102  
 HALL 1970 (463) 86, 85  
 HALL 1972 (433) 80, 79  
 HALPERIN, McFARLAND, NIVEN and  
 ROUGHTON 1959 (440) 83, 81  
 HAMEL-PUSKARIC, BERITIC, JUSIC and  
 FRANJIC 1970 (A 490) 207  
 HAMILL and O'NEILL 1969 (77) 22, 21  
 HAMILTON and KERSTING 1970 (A 395)  
 202  
 HAMM 1966 (A 396) 202  
 HANISCH 1969 (A 98) 182  
 HANKE and KIERES 1967 (602) 110, 108  
 HANKS 1970 (483) 91, 89  
 HANQUET and LAMY 1971 (A 813) 230  
 HANSEN 1970 (A 491) 207  
 HANSEN, WILKE, MALORNY and GOTHERT  
 1972 (47) 17, 15; (153) 34  
 HANSON and HASTINGS 1933 (78) 22,  
 34, 33  
 HANSSON and SUNDSTROM 1969 (A 27)  
 177  
 HANSZ and STYPEREK 1968 (464) 86, 85  
 HARADA and KOZUMA 1968 (534) 97, 96  
 (A 492) 207  
 HARADA, TSUKAYAMA, MIMURA, MINAMI  
 and TATETSU 1971 (535) 97, 96  
 HARDING, WONG and NELSON 1964  
 (603) 110, 108  
 HARKE 1970 (95) 26, 24, 25  
 HARKE 1971 (100) 26, 25  
 HARKE and DREWS 1968 (121) 30, 29  
 HARTRIDGE 1920 (48) 17, 11, 14  
 HATZFELD, WIENER and BRISCOE 1967  
 (A 397) 202  
 HAYASINI, MOTOKAWA and KIKUCHI  
 1966 (A 99) 182

1005051303



- HAYES and HALL 1964 (244) 54, 52  
 HAYWOOD, WALBERG, KERR, MOHSEININ  
 and MOHLER 1972 (321) 64, 61  
 HEGGLIN 1944 (245) 54, 52  
 HEIDRICH, BARCKOW and FRISIUS 1970 (339)  
 67, 66  
 HEIDRICH and KLEMS 1969 (373) 72, 69  
 HEISTAD and WHEELER 1972 (340) 67, 66  
 HELLUNG-LARSEN, KJELDSSEN, MELLEMGAARD  
 AND ASTRUP 1966 (A 28) 177  
 HELLUNG-LARSEN, LAURSEN, KJELDSSEN and  
 ASTRUP 1968 (374) 72, 70  
 HELMCHEN and KUNKEL 1964 (441) 83, 81  
 HERNANDEZ, MAZEL and GILLETTE 1967  
 (A 556) 211  
 HERON 1962 (581) 106, 102, 104  
 HESS 1971 (101) 26, 25  
 HEXTER and GOLDSMITH 1971 (A 256) 192  
 HEYDENREICH 1970 (442) 83, 81  
 HEYNDRIKX, SCHEIRIS, VERCRUYSE and  
 OKKERSE 1970 (A 753) 226  
 HILDEBRANDT, FRANKLIN, ROOTS and  
 ESTABROOK 1971 (A 100) 182  
 HILPERT 1971 (A 398) 202  
 HIRANO, INOUE and TANAMI 1967 (604)  
 110, 109  
 HIRANO, INOUE and TANAMI 1968 (605)  
 110, 108  
 HIRANO, INOUE and TANAMI 1971 (A 814)  
 230  
 HIRATA, HIOKI and HASHIMOTO 1969 (510)  
 94, 93  
 HIRSCH 1968 (A 594) 214  
 HLAVICA, KIESE, LANGE and MOR 1969  
 (A 557) 211  
 HOCHSTRATE and OBERDISSE 1970 (A 558)  
 211  
 HODY and BAILEY 1968 (A 171) 187  
 HOFREUTER, CATCOTT and XINTARAS  
 1962 (49) 17, 15; (154) 34, 33  
 HOLCZABEK 1971 (286) 59, 58  
 HOLETON 1971 (A 595) 214  
 HOLETON 1971 (A 596) 214  
 HOLLAND 1965 (A 101) 182  
 HOLLAND 1967 (A 102) 182  
 HOLLAND 1969 (A 104) 182  
 HOLLAND 1969 (A 103) 182  
 HOLLAND 1970 (A 105) 182  
 HOLM 1950 (287) 59, 57  
 HOLMGREN 1965 (A 402) 202  
 HOLMGREN 1965 (A 399) 202  
 HOLMGREN 1965 (A 400) 202  
 HOLMGREN 1965 (A 401) 202  
 HORIE 1964 (A 106) 182  
 HORIE 1965 (A 107) 182  
 HORVATH, DAHMS and O'HANLON 1971  
 (484) 91, 89  
 HOSKO 1970 (443) 83, 81  
 HOWSE and SEDDON 1966 (620) 112, 111  
 HSIEH, ROSS, SMALL and THOMPSON 1968  
 (A 403) 202  
 HSI-PU and LI-MING 1910 (79) 22, 21  
 HUBER, EPP and FORMANEK 1970  
 (A 108) 182  
 HUBERT 1943 (322) 64, 63  
 HUEPER 1944 (375) 72, 70  
 HUGHES and FISHER 1965 (A 674) 219  
 HUNDT and GRUNBERG 1960 (246)  
 54, 52  
 HYDE, MARIN, RYNE, KARREMAN  
 1971 (A 404) 202  
 IABLOCHKIN 1966 (A 335) 198  
 ICHIKAWA, HAGIHARA and YAMANO  
 1967 (A 559) 211  
 IKEDA 1969 (444) 83, 81  
 IKUTA 1969 (A 494) 207  
 INANAGA 1966 (A 497) 208  
 INANAGA 1968 (A 498) 208  
 INANAGA, KUHARA, KUWAHARA,  
 TORISU and SUZUKI 1966A (536)  
 97, 96  
 INANAGA, KUHARA, KUWAHARA and  
 OGATA 1966B (537) 97, 96  
 INGIULLA, GRASSO and MARIOTTINI  
 1968 (A 288) 194  
 IPPEN and GOERZ 1969 (621) 112, 111  
 ISHIKAWA 1969 (A 499) 208  
 IWANOWSKA 1967 (A 815) 230  
 JACOBSEN 1971 (A 816) 230  
 JAFFE 1968 (A 190) 189  
 JAFFE 1970 (A 191) 189  
 JAFFE 1965 (288) 59, 57  
 JAFFE 1968 (323) 64, 61  
 JAGI and ZIMMERMANN 1934 (289)  
 59, 56  
 JAMES and RUMBLE 1967 (A 405) 202  
 JARRELL 1965 (122) 30, 28  
 JAVANOVIC and POLOVINA 1964 (A 29)  
 177  
 JECH 1972 (14) 9, 6  
 JEDRYCHOWSKI, KUS, PIOTROWSKI  
 and SAWICKI 1965 (A 560) 212  
 JEDRYCHOWSKI, KUS, PIOTROWSKI and  
 SAWICKI 1965 (A 561) 212  
 JEFFCOATE and GAYLOR 1969 (A 109) 182  
 JENEY and MEDVE 1967 (A 597) 214  
 JESCHECK 1967 (A 500) 208  
 JOELS and NEIL 1962 (216) 49, 47  
 JOHNSON, DWORETZKY and HELLER 1968  
 (A 289) 195  
 JOHNSON and MILLER 1968 (A 408) 202  
 JOHNSON, TAYLOR and De GRAFF 1965  
 (A 407) 202  
 JOHNSON, TAYLOR and LAWSON 1965  
 (A 406) 202  
 JOHNSTON and BURGER 1971 (A 172) 187  
 JONES, YANT and BERGER 1923 (102) 26, 24  
 JONES, STRICKLAND, STUNKARD and  
 SIEGEL 1971 (413) 77, 75  
 JONGBLOED 1939 (80) 22, 19  
 JOPKIEWICZ, KONECKI and WENTKOWSKI  
 1965 (622) 112, 111  
 JORDI 1967 (A 501) 208  
 JORDI 1968 (A 703) 223  
 JUDD 1971 (A 173) 187

1005051304

- JUNGE, SEILER, BROCK, GREESE and  
RADLER 1971 (a 192) 189
- KALIAEVA 1951 (290) 60, 56
- KAMATAKI and KITAGAWA 1971 (A 562) 212
- KAMPFFMEYER and KIESE 1965 (A 563) 212
- KAMRAJ-MAZURKIEWICZ 1967 (581) 106, 102
- KANAZIRSKY 1965 (A 409) 202
- KARACAN, BARNARD and WILLIAMS 1971  
(538) 97, 96
- KASHIMA, FUKUI, MASUDA, WAKASUGI and  
HAYAMA 1969 (675) 219
- KATSUKI 1965 (A 502) 208
- KATSUKI 1966 (606) 110, 108
- KATSURA 1971 (445) 83, 82
- KATZSCHMANN 1970 (291) 60, 56
- KAWAMOTO 1966 (A 410) 202
- KAWAMURA 1971 (465) 86, 85
- KAYES 1965 (14a) 8, 6
- KAYES 1970 (A 632) 216
- KAYSER 1939 (341) 67, 66
- KEHL and KEHL 1967 (A 503) 208
- KEITH and TESH 1965 (123) 30, 29
- KELLS 1968 (582) 106, 102
- KEMKES 1941 (A 704) 223
- KENT 1970 (A 174) 187
- KERTESZ, ANTONINI, BRUNORI, WYMAN and  
ZITO 1965 (A 175) 187
- KEYES, MIZUKAMI and LUMRY 1967 (A 110)  
83
- KHACHATURYAN, MITAREVSKAYA and  
EGORENKOVA 1969 (511) 94, 93
- KHROLENKO 1969 (A 504) 208
- KILLICK 1940 (15) 9, 6
- KILLICK and MARCHANT 1965 (A 754) 226
- KIM and PARK 1968 (A 633) 216
- KIM and RYO 1966 (A 336) 198
- KIRIACHKO 1966 (217) 49, 48
- KITTEL and THEISSING-ERLANGEN 1968  
(466) 86, 85
- KITTREDGE 1971 (A 676) 219
- KJELDSSEN 1969 (376) 73, 69
- KJELDSSEN 1970a (377) 73, 70
- KJELDSSEN 1970b (378) 73, 70
- KJELDSSEN, ASTRUP and WANSTRUP 1972  
(379) 73, 70
- KJELDSSEN and DAMGAARD 1968 (380) 73, 70
- KJELDSSEN and DAMGAARD 1968 (414) 77, 74
- KJELDSSEN and MOZES 1969 (381) 73, 69
- KLAUSEN, RASMUSSEN, GJELLEROD, MADSEN  
and PETERSEN 1968 (203) 46, 45,  
(292) 60, 56
- KLAUIS and SCHULZ 1966 (293) 60, 57
- KLEDECKI and WINIARSKI 1963 (247) 54, 52
- KLINGENMAIER, BEHAR and SMITH 1969  
(A 30) 177
- KLINGHOFFER 1965 (A 677) 219
- KOBUTNICKY 1966 (553) 101, 99
- KOCII 1965 (A 31) 177
- KOELSCH 1936 (294) 60, 57
- KOHN-ARREST 1949 (124) 30, 29
- KOKAME and SIHLER 1968 (A 817) 230
- KOJLB 1968 (623) 112, 111
- KOJLB 1968 (624) 112, 111; (A 505) 208
- KOMURA 1967 (539) 97, 96
- KONDRASHENKO, GLANTS and MAEROVICH  
1971 (A 818) 230
- KORNER 1965 (218) 49, 47
- KOSMIDER, ZURKOWSKI and WEGIEL 1965  
(A 111) 183
- KOSTLER, OTTO, RITTIG and POLSTER  
1967 (A 32) 177
- KOSTYUKOVA 1951 (248) 54, 52
- KOTTER, HUICH, STOTZ and PIPER 1969  
(A 411) 203
- KRAL, CERNOCHOVA and TUSL 1966  
(A 412) 203
- KRATOCHVIL, WILKS and GERRARD 1957  
(485) 91, 89
- KRATZ 1968 (A 564) 212
- KRETZER 1964 (A 697) 222
- KREUKNIET 1964 (A 413) 203
- KREUZER and CAMPAGNE 1965 (413a) 203
- KRISHMAN, KUPPUSWAMY, MANI and MAJID  
1971 (A 678) 219
- KROBER, LANGE, MATHIES and MOR  
1968 (A 565) 212
- KROBER, LANGE, MATHES and MOR  
1970 (A 566) 212
- KROETZ 1936 (295) 60, 56
- KROETZ 1936a (324) 64, 63
- KROETZ 1936b (325) 64, 63
- KRUG 1965 (A 506) 208
- KRUSMANN, SCHRODER and SCHRODER  
1971 (A 33) 177
- KRUSZYNSKI and HENRIKSEN 1969 (125)  
30, 28
- KUCHER and RIEDEL 1969 (A 819) 230
- KUNZ, DONDES and HARTECK 1970 (A 337)  
198
- KUNTZMAN, LEVIN, JACOBSON and  
CONNEY 1968 (A 567) 212
- KUPFER and WUNSCHER 1968 (512) 94, 93
- KUROIWA 1967 (A 679) 219
- KUROIWA, MURAO, HARUMI, KATAYAMA,  
YAMAMOTO, CHEN and UEDA  
1968 (249) 54, 52
- KUROIWA, KATO and UMEZAKI 1968  
(540) 97, 96
- KUROIWA, SHIDA and KATO 1969 (A 507)  
208
- KUROIWA, SHIDA, NAGAMATSU, KATO and  
SANTA T 1967 (446) 83, 81
- KUTTNER 1968 (467) 86, 85
- LACHNIT 1964 (A 634) 216
- LACOSTE 1971 (A 415) 203
- LACOSTE and ROUCH 1966 (A 414) 203
- LAMY and HANQUET 1968 (A 820) 230
- LAMY and HANQUET 1969 (A 821) 231
- LAMY and NOIRFALISE 1971 (A 680) 219
- LANDAU, SMITH and LYNN 1969 (A 290)  
195
- LANDAW 1969 (A 222) 190
- LANDAW 1970 (A 223) 191
- LANDAW, CALLAHAN and SCHMID 1970  
(A 112) 183

- LANDAW and WINCHELL 1966 (A 221) 190  
 LANDAW and WINCHELL 1970 (A 224) 191  
 LANG, SCHUSTER, UNGERN-STERBERG, HAUM  
 and KNOLLE 1969 (250) 54, 52  
 LANGAUER-LEWICKA 1966 (A 508) 208  
 LANGE, KASTNER and JUNG 1970 (A 568) 212  
 LANGMANN 1964 (A 257) 192  
 LANGMANN and KETTNER 1968 (A 291) 195  
 LAPRESLE and FARDEAU 1966 (A 509) 208  
 LAPRESLE and FARDEAU 1967 (A 510) 208  
 LAPRLE and FARDEAU 1971 (A 511) 208  
 LARCAN, LANDES and VERT 1970 (583)  
 106  
 LARCAN, ROBERT, CALAMAI and FREJAVILLE  
 1967 (A 822) 231  
 LARSON, HAAG and SILVETTE 1961 (16)  
 9, 7  
 LARSON and SILVETTE 1968 (17) 9, 7  
 LARSON and SILVETTE 1971 (18) 9, 7  
 LATALSKI and PAWLOWSKA 1969 (A 569) 212  
 LATALSKI and PAWLOWSKA 1970 (A 570) 212  
 LAVERNE 1970 (486) 91, 88  
 LAVERNE 1970 (A 755) 226  
 LAWSON 1970 (A 416) 203  
 LAWSON 1971 (A 113) 183  
 LAWSON 1972 (A 417) 203  
 LAWTHER and COMMINS 1970 (18a) 9, 7;  
 (155) 34, 33  
 LEAVELL and McINTYRE 1969 (625) 112, 111  
 LEBRETON and GARAT 1964 (A 292) 195  
 LeBRETON and GARAT 1964 (A 681) 219  
 LeBRETON and GARAT 1964 (A 682) 220  
 LECLERQ 1970 (19) 10, 6  
 LEDINGHAM 1964 (A 823) 231  
 LEE 1966 (A 824) 231  
 LEE and SCHRAUZER 1968 (A 598) 214  
 LEE 1908 (126) 30, 29  
 LEHMANN 1968 (A 35) 178  
 LEHMANN 1908 (127) 30, 27  
 LEHMANN 1909 (128) 30, 28  
 LEHR 1970 (A 683) 220  
 LEONOWICZ 1967 (554) 101, 99  
 LEUCHTENBERGER, LEUCHTENBERGER and  
 WEISS 1965 (81) 22, 20  
 LEVIN 1965 (A 756) 226  
 LEVIN, ALVARES and KUNTZMAN 1970 (A 114)  
 183  
 LEVIN and KUNTZMAN 1969 (A 572) 212  
 LEVIN and KUNTZMAN 1969 (A 571) 212  
 LEWEY and DRABKIN 1944 (251) 54, 52  
 LEWIS and BRINK 1966 (A 418) 203  
 LEWIS 1967 (A 599) 214  
 LIGHTFOOT 1972 (A 176) 187  
 LIKOFF, SEGAL and KASPIAN 1967 (415)  
 77, 75  
 LILIENTHAL 1950 (20) 10, 6  
 LILIENTHAL and FUGITT 1946 (447) 83, 81  
 LINCII and PFAFF 1971 (A 338) 198  
 LINDERHOLM 1965 (A 36) 178  
 LINDERHOLM 1969 (A 225) 191  
 LINDERHOLM, SJOSTRAND and SODERSTROM  
 1966 (A 37) 178  
 LINDQUIST 1970 (664) 115, 167  
 LINTON, ADAMS and LAWSON 1968  
 (555) 101, 99  
 LITMAN 1968 (A 635) 217  
 LITZNER 1936 (342) 67, 66  
 LIZANETS and ZARKEVICH 1971 (A 757) 226  
 LOCKSHIN and BURRIS 1965 (A 600) 215  
 LO COCO 1970 (A 419) 203  
 LOEYER, VARAY and COTTET 1942  
 (252) 54, 42  
 LOGUE, ROSSE, SMITH, SALTZMAN and  
 GUTTERMAN 1971 (A 226) 191  
 LOMONACO 1971 (A 177) 187  
 LONG 1969 (626) 112, 111  
 LONGO 1970 (584) 106, 102  
 LONGO, POWER and FORSTER 1969 (586)  
 106, 102  
 LONGO, POWER and FORSTER 1967 (585)  
 106, 102  
 LOPEZ-MAJANO 1971 (A 420) 203  
 LORENTE, VARELA and SEIJAS 1953  
 (253) 54, 52  
 LUDERITZ 1971 (182) 40, 38; (A 293) 195  
 LUNDEVALL 1972 (A 705) 223  
 LUOMANMAKI 1966 (A 157) 186  
 LUOMANMAKI and COBURN 1969 (A 158)  
 186  
 LUSTMAN and GEERTS 1971 (254) 54, 52  
 LYNCH and MOEDE 1972 (A 227) 191  
 Mac FARLAND, ROUGHTON, HALPERIN and  
 NIVEN 1944 (82) 22, 21  
 MACHATA 1968 (A 684) 220  
 MACKINTOSH 1965 (A 512) 208  
 MacQUARRIE and GIBSON 1971 (A 115)  
 183  
 MAENO and FEIGELSON 1968 (A 601)  
 215  
 MAGDALENO 1968 (A 178) 187  
 MAHRLEIN 1967 (A 422) 203  
 MAINARDI 1964 (255) 54, 52  
 MAISELS, PATHAK, NELSON, NATHAN  
 and SMITH 1971 (A 228) 191  
 MALIK 1971 (A 636) 217  
 MAMATSASHVILI 1970 (607) 110, 108  
 MANN 1965 (541) 97, 96  
 MANSLEY, STANBURY and LEMBERG 1966  
 (A 116) 183  
 MANTELL 1964 (587) 106  
 MANTZ and TEMPE 1968 (A 825) 231  
 MANTZ and TEMPE 1968 (A 826) 231  
 MARANZANA 1964 (A 706) 223  
 MARCELET 1907 (129) 30, 29  
 MARCHIARO, MARGARIA, CAIDO and  
 AQUARO 1964 (A 707) 223  
 MARI and RIZZATTI 1964 (A 685) 220  
 MARKIEWICZ 1966 (A 38) 178  
 MARKIEWICZ 1967 (A 229) 191  
 MARKIEWICZ 1970 (103) 26, 24  
 MARKS and SWIECICKI 1971 (513)  
 95, 93  
 MARLAND and BERSAY 1972 (326) 64, 61  
 MATSUYAMA 1969 (588) 106, 102  
 MATTHEW 1970 (A 758) 226  
 MATTHEW 1970 (A 827) 231

1005051306

- MATTHEW 1971 (A 637) 217  
 MATTHEW and PROUDFOOT 1965 (A 513) 208  
 MAUGH 1972 (A 193) 189  
 MAUNDERLY 1972 (A 121) 203  
 MAURER 1941 (343) 67, 66  
 MAURER 1941 (134) 80, 79  
 MAUTNER 1955 (382) 73, 69  
 MAWATARI 1970 (511) 95, 93  
 MAZALESKI, COLEMAN, DUNCAN and NAU 1970 (A 573) 212  
 McBay 1965 (A 686) 220  
 Mc CONNELL, DEAL and OGATA 1969 (A 117) 183  
 McCREMIE and JOSE 1967 (A 39) 178  
 McDOWELL 1971 (339) 198  
 Mc FARLAND 1952 (187) 91, 88  
 McFARLAND 1970 (148) 83, 81  
 McFARLAND 1971 (A 179) 187  
 McFARLAND, ROUGHTON, HALPERIN and NIVEN 1944 (149) 83, 81  
 McFEE, LAVINE and SULLIVAN 1970 (A 340) 198  
 McGRATH and JAEGER 1971 (A 602) 215  
 Mc GRATH and MOFFA 1972 (A 603) 215  
 McILVAINE, NELSON and BARTLETT 1969 (50) 18, 15, (156) 34, 33  
 McMILLAN and COPE 1969 (A 194) 189  
 McNALLY 1931 (A 687) 220  
 MEDA 1964 (A 118) 183  
 MEDVEDOWSKY, SACCO and BELZUNCE 1965 (256) 54, 52  
 MEIGS 1948 (52) 13, 14; (157) 34, 33  
 MEIGS and RYAN 1971 (607) 110, 108  
 MELANOWSKI 1963 (150) 83, 82  
 MEL'NICHENKO 1968 (A 119) 183  
 MENKES, SERA, ROGERS, HYDE, FORRESTER and DuBOIS 1970 (A 423) 203  
 MENZ 1966 (A 688) 220  
 MERLI 1969 (A 180) 187  
 MESOLELLA, PERRELLA, TESTA and MORELLI 1970 (468) 86, 85  
 METCALFE, MALL, BARTELS, HILPERT and PAPER 1965 (589) 106, 102  
 MEYER, GROVER and WEIL 1972 (219) 49, 47  
 MIHAJ and WEBER 1964 (344) 67, 66  
 MIKULPA 1970 (488) 91, 89  
 MILLER 1966 (A 828) 231  
 MILLS and EDWARDS 1968 (220) 49, 47  
 MIRANDA, KONOPINSKI and LARSEN 1967 (204) 46, 45  
 MIRKIN 1966 (A 294) 195  
 MISCHENKO and FRENKEL 1966 (515) 95, 93  
 MITCHELL and RENZETTI 1968 (A 424) 203  
 MITTMAN 1967 (A 425) 203  
 MIYAGISHI and HAYASHI 1968 (517) 95, 93  
 MIYAGISHI and SUWA 1969 (516) 95, 93  
 MIYAHARA and TAKAHASHI 1971 (A 230) 191  
 MOC, RICKARD and MOSS 1969 (A 601) 215  
 MOKHOV 1967 (A 341) 198  
 MOLFINO and ZANNINI 1964 (A 829) 231  
 MONACO 1964 (A 708) 223  
 MONAUI 1940 (296) 60, 57  
 MONTGOMERY and RUBIN 1971 (A 574) 213  
 MOON and RICHARDS 1972 (A 120) 183  
 MOORE and FINESTONE 1968 (A 514) 208  
 MORANDO and ROVIDA 1965 (183) 40, 38  
 MORGANSTERN, ASH and LYNCH 1970 (A 342) 198  
 MOROVIC 1968 (A 515) 208  
 MORRIS 1969 (469) 86, 85  
 MORRISON and HORIE 1965 (A 121) 183  
 MORROW 1967 (A 258) 192  
 MOSINGER, BISSCHOP and LUCCIONI 1969 (257) 54, 52  
 MOSS 1969 (A 709) 223  
 MOTLEY 1971 (52) 18, 15; (158) 35, 34  
 MOTTA 1940 (258) 54, 52  
 MOUNIER-KUHN, ROCHE, MORGAN and BERNAR 1968 (170) 86, 85  
 MOUNTAIN, CASSELL, WOLTER MOUNTAIN, DIAMOND and McCARROLL 1968 (184) 4, 37  
 MOUREN, POINSO, JOUGLARD, GIUDICELLIS, FRESCO and D'OMEZON 1972 (627) 112, 111  
 MOUREU 1964 (185) 4, 38  
 MULHAUSEN, ASTRUP and KJELDSSEN 1967 (382A) 73, 69  
 MULHAUSEN, ASTRUP and MELLEMGAAARD 1968 (A 159) 186  
 MULLER and HUNG 1968 (A 689) 220  
 MULLER and VOIGT 1968 (416) 77, 74  
 MUMPOWER, LEWIS and TOUEY 1962 (130) 31, 29  
 MUROFUSCHI and MINAGAWA 1969 (A 690) 220, (A 516) 209  
 MURPHY and MULCAHY 1971 (590) 106, 103  
 MURPHY, LENG, ULRICH and DAVIS 1963 (A 295) 195  
 MURRAY 1971 (A 122) 183

1005051307

- NAGANO 1967 (A 123) 183  
 NAGEL 1937 (297) 60, 57  
 NAGEL and GIBSON 1966 (A 124) 183  
 NAGEL, GIBSON and HAMILTON 1971 (A 126) 183  
 NAGEL, GIBSON and JENKINS 1971 (A 125) 183  
 NAHUM 1965 (665) 115, 168  
 NAHUM 1968 (666) 115, 169  
 NAHUM 1969 (667) 115, 170  
 NAIRN, POWERM HYDE, FORSTER, LAMBERTSEN and DICKSON 1965 (A 426) 203  
 NAKAO 1969 (608) 110, 109  
 NASH and BEEBE 1969 (628) 112, 111  
 NATIONAL ACADEMY OF SCIENCES AND NATIONAL ACADEMY OF ENGINEERING 1969 (21) 10, 7  
 NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS 1964 (22) 10, 8  
 NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS 1967 (23) 10, 8; (635) 114, 117  
 NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS 1968 (24) 10, 8; (636) 114, 119  
 NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS 1969 (25) 10, 8; (637) 114, 121  
 NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS 1971 (26) 10, 8; (638) 114, 123  
 NATIONAL CLEARINGHOUSE FOR SMOKING AND HEALTH USPHS 1972 (27) 10, 8; (639) 114, 128  
 NATIONAL INSTITUTE FOR OCCUPATIONAL SAFETY AND HEALTH 1972 (28) 10, 7  
 NAVRATIL 1956 (345) 67, 66  
 NECAS and NEUWIRT 1971 (518) 95, 93  
 NEWSOME and KEITH 1965 (131) 31, 29  
 NICOLAS and NICOLAS 1964 (556) 101, 99  
 NIDEN and SCHULZ 1965 (221) 49, 47  
 NIEBROJ 1969 (451) 83, 82  
 NIELSEN 1971 (489) 91, 88, (A 759) 226  
 NIELSEN 1971 (346) 67, 66  
 NISHIGORI 1932 (347) 67, 66  
 NISHII 1968 (A 296) 195  
 NISSARDI, SANNA-RANDACCIO and SARNA 1968 (A 430) 204  
 NISSARDI, SANNA-RANDACCIO, TORRAZZA and CASCIU 1965 (A 427) 204  
 NISSARDI, SANNA-RANDACCIO, TORRAZZA and GARIEL 1965 (A 428) 204  
 NISSARDI, TORRAZZA and ANEDDA 1967 (A 429) 204  
 NIZHEGORODOV and MARCHOZKY 1969 (A 575) 213  
 NOBEL and RICKER 1967 (A 40) 178  
 NOBLE, BRUNORI, WYMAN and ANTONINI 1967 (A 127) 183  
 NOBLE, PARKHURST and GIBSON 1970 (A 128) 184  
 NORMAN, DOUGLAS and SMITH 1966 (222) 49, 48  
 NORMAN and LEDINGHAM 1967 (A 760) 226  
 NORMAN, MacINTYRE, SHEARER and SMITH 1970 (A 830) 231  
 NOUILHAT 1964 (A 761) 226  
 OBERSTEG and DELAY 1966 (691) 220  
 O'DONNELL, CHIKOS and THEODORE 1971a (490) 91, 89  
 O'DONNELL and MIKULKA 1970 (492) 91, 89  
 O'DONNELL, MIKULKA, HEINIG and THEODORE 1971b (491) 91, 89  
 OETTEL 1967 (104) 26, 24  
 OGATA 1968 (A 495) 207  
 OHARA 1968 (452) 83, 82  
 OIRESSER, CHASSON, JOUGLARD, GOUIN, DUBOULOZ and TASSY 1968 (A 831) 231  
 OKALYI 1969 (A 181) 187  
 OKESON and DIVERTIE 1970 (223) 49, 47  
 OKULOVSKI and HACHATUROV 1968 (A 762) 227  
 OKUMA, ISHINO, SUNAMI and MOTOIKE 1968 (A 517) 209  
 OKUNYEV and PROKHORENKO 1966 (519) 95, 93  
 OMURA and SATO 1964 (A 576) 213  
 OMURA, SATO, COOPER, ROSENTHAL and ESTABROOK 1965 (A 129) 184  
 ORINIUS 1968 (259) 54  
 ORIOLI and CATTANIA 1965 (542) 97, 96  
 ORIZAGA and DUCHARME 1967 (383) 73, 69  
 OSBORNE, ADAMEK and HOBBS 1956 (132) 31, 29  
 OSKI, GOTTLIEB, MILLER and DELIVORIA-PAPADOPOULOS 1970 (A 130) 184  
 OSTROWSKA 1967 (A 518) 209  
 OSWALD 1966 (A519) 209  
 OTIS 1970 (417) 77, 75  
 OTSUKA, FUJIWARA, IKAWA and HIRAYAMA 1970 (133) 31, 29  
 OTSUKI, NINOMIYA, YAMAMOTO, NAKASHIMA, SHUTARA, URAKAMI and ITAMI 1966 (A 763) 227  
 OWEN and REYNOLDS 1967 (134) 31, 28  
 PAEZ 1970 (A 431) 204  
 PALADE, MIHAI, GOILAV and SOVAREL 1969 (A 520) 209  
 PALMA-CARLOS, PALMA-CARLOS and SOARES 1964 (A 41) 178  
 PALMA-CARLOS, PALMA-CARLOS and SOARES 1966 (A 231) 191  
 PAPAVASLIOU, COTZIAS, DUBY, STECK, FEHLING and BELL 1972 (A 764) 227  
 PARADE and FRANKE 1939 (260) 54, 51  
 PARE and YEUNG 1969 (520) 95, 93  
 PARIS 1964 (629) 112, 111  
 PARKHURST, GERACI and GIBSON 1970 (A 132) 184

1005051308

- PARKHURST and GIBSON 1967 (A 131) 184  
 PARMEGGIANI and GILARDI 1952 (53) 18, 13, 14;  
 (83) 22, 21; (159) 35, 33  
 PARROT, STUPFEL, ROMARY and MORDELET-  
 DAMBRINE 1971 (224) 50, 47  
 PASECHNIK, SHTUMM, VLADISLAVLEV and  
 ZAMAYATNIN 1971 (A 133) 184  
 PATTONO, MARCHIARO, CAPELLARO and  
 ORIONE 1964 (A 160) 186  
 PATZ 1949 (261) 55, 51  
 PAULEIKOFF, MULLER-FAHBUSCH, MESTER  
 and MEIBNER 1971 (A 521) 209  
 PAULET and CHEVRIER 1966 (A 134) 184  
 PAULET and CHEVRIER 1969 (A 765) 227  
 PAULI, TRUNIGER, LARSEN and MULHAUSEN  
 1968a (557) 101, 99  
 PAULI, TRUNIGER, LARSEN and MULHAUSEN  
 1968b (558) 101, 100  
 PEARCE 1968 (A 522) 209  
 PECORA 1964 (A 766) 227  
 PEDRERO and RODRIGO 1964 (A 347) 198  
 PERRELLI, PREVOT and SULOTTO 1970 (418)  
 77, 74  
 PERRELLI and ROSETTANI 1964 (A 297) 195  
 PERRELLI, ROSETTANI and BRAGUZZI 1965  
 (A 278) 195  
 PETERSON, SIGGAARD-ANDERSEN,  
 KRISTENSEN and KJELDSSEN 1968 (348) 67,  
 66  
 PETERSON and STEWART 1970 (A 161) 186  
 PETIT, PETIT and GEILLE 1970 (A 692) 220  
 PETRILLI and KANITZ 1970 (186) 41, 38, 39  
 PETROVA, DALAKMANSKI and BAKALOV  
 1966 (A 299) 195  
 PETROVIC 1970 (419) 77, 75  
 PETTER, BOURBON, MALTIER and JOST  
 1971 (A 135) 184  
 PETTY 1969 (A 832) 231  
 PHELPS and ANTONINI 1969 (A 136) 184  
 PHILIPPE and HOBBS 1956 (135) 31, 29  
 PLACE 1970 (A 767) 227  
 PICKWELL 1970 (A 195) 189  
 PIEDELIEVRE, BRETON and DEROBERT  
 1969 (591) 106, 102  
 PIERCE and COLLINS 1971 (A 343) 198  
 PINCHERLE and SHANKS 1967 (420) 77, 75  
 PIPER, PFEIFER and SCHEID 1969 (A 433) 204  
 PIPER and SIKAND 1966 (A 432) 204  
 PIRNAY, DEROANNE, DUJARDIN and PETIT  
 1971b (299) 60, 56  
 PIRNAY, DUJARDIN, DEROANNE and PETIT  
 1971a (298) 60, 56  
 PIRNAY, FASSOTTE, DEROANNE and PETIT  
 1968 (A 137) 184  
 PIRNAY, FASSOTTE, GAZON, DEROANNE  
 and PETIT 1969 (A 434) 204  
 PIRNAY, PETIT and ROBERTS 1970 (A 435)  
 204  
 PODLESCH and STEVANOVIC 1966 (A 436) 204  
 POGRUND 1969 (609) 110, 108  
 POLITZER 1968 (A 138) 184  
 POLLARD 1970 (A 693) 220  
 PORTHEINE 1971 (105) 26, 25  
 POWER 1968 (A 162) 186  
 POWER, AOKI, LAWSON and GREGG 1971  
 (A 438) 204  
 POWER, HYDE, SEVER, HOPPIN and NAIRN  
 1965 (A 437) 204  
 PRELLWITZ, SCHUSTER, SCHYLLA, BAUM,  
 SCHONBORN, UNGERN-STERMBERG,  
 BRODERSEN and POEPLAU 1970 (421) 77, 74;  
 (A 638) 217  
 PREROVSKA and DRDKOVA 1967a (384) 73, 70  
 PREROVSKA and DRDKOVA 1967b (385) 73, 70  
 PREROVSKA and DRDKOVA 1971 (386) 73, 71  
 PREZIOSI, LINDENBERG, LEVY and  
 CHRISTENSON 1970 (521) 95, 93  
 PROKOP and WABNITZ 1970 (453) 83, 81  
 PUKHOV 1964 (610) 110, 108  
 PUKHOV 1965 (A 768) 227  
 PUREC and KRASNA 1967 (A 605) 215  
 QUINTANA, MIRETE and GARCIA 1969 (559)  
 101, 99  
 RADUSHVICH 1968 (A 769) 227  
 RAMSEY 1972 (84) 22, 20  
 RAMSEY 1966 (A 300) 195  
 RAMSEY 1967 (A 344) 198  
 RAMSEY 1969 (422) 77, 75  
 RAMSEY 1970 (188) 41, 37  
 RAMSEY 1972 (493) 91, 89  
 RAMSEY 1967 (187) 41, 38, 39  
 RANDOWA 1967 (A 440) 204  
 RANDOWA and SIERAWSKI 1964 (A 439) 204  
 RAPOPORT 1967 (A 770) 227  
 RAUSA, DIANA and PERIN 1968 (A 139) 184  
 RAUSA, PERIN and DIANA 1967 (A 300) 195  
 RAY and ROCKWELL 1970 (494) 91, 89  
 RAY 1967 (A 606) 215  
 RAYFIELD 1967 (A 301) 195  
 RECKZEH and DONTENWILL 1970 (423) 77, 75  
 RECKZEH, RUCKER, HARKE and DONTENWILL  
 1969 (106) 26, 25  
 REDDEMANN, AMENDT and JAHRIG 1970  
 (A 639) 217  
 REED and TROTT 1971 (A 302) 195  
 REED 1970 (A 140) 184  
 RIECHEL, WOBITH and ULMER 1970 (189) 41, 37  
 REJSEK 1971 (A771) 227  
 REMMERS and MITHOEFER 1969 (A 441) 204  
 REPLOH, KLOSTERKOTTER and  
 EINCK-ROSSKAMP 1966 (A 710) 223  
 REVOL, MONIER, COURJON, FOURMET and  
 GERIN 1966 (543) 97, 96  
 REVSIN and BRODIE 1969 (A 607) 215  
 RHODES 1971 (225) 50, 48  
 RICCI, CAPELLARO and GAIDO 1964 (424) 77, 74  
 RI 1966 (630) 112, 111  
 RIKANS and VAN DYKE 1971 (A 577) 213  
 RINGEL and KLAUANS 1972 (A 523) 209  
 RINGOLD, GOLDSMITH, HELLWIG, FINN and  
 SCHUETTE 1962 (85) 23, 19

1005051309

- RISPLER and ROSS 1965 (A 303) 195  
 RITTER 1956 (300) 60, 57  
 RITUCCI and LUVONI 1965 (A 694) 220  
 RIZZI 1968 (A 772) 227  
 ROBBINS, BORG and ROBINSON 1968 (A 345) 198  
 ROBIN, RAVENS and BING 1969 (327) 65, 61  
 ROBINSON and ROBBINS 1970 (A 196) 189  
 ROCHE, BERTOYE, VINCENT, MOTIN, GARIN, BOLOT and CHADENSON 1968 (A 833) 231  
 RODKEY 1970 (A 346) 198  
 RODKEY and COLLISON 1970 (A 42) 178  
 RODKEY, COLLISON and ENGEL 1969 (A 711) 223  
 RODKEY, COLLISON and O'NEAL 1971 (A 43) 178; (A 182) 187  
 RONDIA 1970 (A 578) 213  
 RONDIA, GUYAUX and HEUSGHEM 1966 (A 304) 195  
 ROOT 1962 (29) 10, 6  
 ROPSHITZ and OVENSTONE 1968 (A 640) 217  
 ROSE, JONES, JENKINS and SIEGEL 1970 (522) 95, 93  
 ROSE 1969 (A 695) 220  
 ROSE and ROSE 1971 (668) 115, 172  
 ROSENBERG 1968 (86) 23, 20  
 ROSENBERG 1971 (87) 23, 20  
 ROSENBERG 1972 (88) 23, 20  
 ROSENBLUTH 1968 (A 524) 209  
 ROSENTHAL 1968 (A 696) 220  
 ROSKAMM 1964 (301) 60, 56  
 ROSSI-FANELLI and ANTONINI 1958 (631) 112, 111  
 ROSSIN and ROBERTS 1972 (30) 10, 7  
 ROSSO and DUGHERA 1964 (A 712) 223  
 ROUCH, RIOUFOL and BOURBON 1971 (54) 18, 13, 15; (160) 35, 33a  
 ROUGHTON 1970 (A 141) 184  
 RUBINO 1964 (226) 50, 47  
 RUDOLPH, BOYLE, DRESDEN and GILL 1972 (632) 112, 11  
 RUEL and BARTHE 1954 (55) 18, 14; (161) 35, 33  
 RUHL and LIN 1936 (56) 18, 14; (89) 23, 20, 21  
 RUMEN and CHANCE 1970 (A 142) 184  
 SADOKIERSKI 1965 (A 713) 223  
 SAITA and LUSSANA 1971 (435) 80, 79  
 SALNIS and HACHIATUROV 1970 (A 773) 227  
 SANDERS and WARRINGTON 1971 (A 525) 209  
 SANNA-RANDACCIO and NISSARDI 1969 (A 442) 205  
 SANZHEVA 1970 (A 608) 215  
 SANZHEVA and ZAVARZIN 1971 (A 609) 215  
 SARACOGU 1951 (262) 55, 51  
 SARTORELLI 1967 (A 403) 205  
 SARUTA 1937 (136) 31, 29  
 SASAKI, HIRANO, NAGAHAMA and USUI 1966 (544) 98, 96  
 SATAKE, HIDA, TATSUBANA, YAMAZAKI and MATSUOKA 1968 (A 444) 205  
 SATO 1966 (471) 86, 85  
 SATOH, KIYOTANI, MINAFI and KONDO 1966 (A 697) 220  
 SAVATEEV, TONKOPII and FROLOV 1970 (A 834) 231  
 SAYERS and DAVENPORT 1930 (31) 10, 6  
 SAYERS, YANT, LEVY and FULTON 1929 (495) 91, 88  
 SCHAEFER 1964 (A 183) 187  
 SCHAFFERNICHT, AIEGLER and REINHARD 1970 (A 714) 223  
 SCHIECHE, KEBLER and KOBER 1970 (A 715) 223  
 SCHIEVELBEIN 1968 (32) 10, 7  
 SCHIEVELBEIN and EBERHARDT 1972 (328) 65, 61  
 SCHLECHT 1971 (A 610) 215  
 SCHMELZER, STEINER, MAYER, NEDETZKA and FASOLD 1972 (A 143) 184  
 SCHMIDT 1970 (454) 83, 82  
 SCHMIDT 1971 (107) 26, 25  
 SCHMIDT 1939 (57) 18, 13, 14; (90) 23, 21; (162) 35, 33  
 SCHMIDT 1940 (58) 18, 14; (91) 23, 21  
 SCHOTT, TOMMASI, BOURRAT and MICHEL 1967 (A 526) 209  
 SCHRAUZER and LEE 1970 (A 611) 215  
 SCHRENK 1942 (59) 18, 13, 14  
 SCHULTE 1963 (496) 92, 88  
 SCHULTE 1965 (A 835) 231  
 SCHUTTMANN 1968 (A 716) 223  
 SCOPPETTA 1968 (592) 106  
 SCORER 1971 (A 259) 192  
 SEGAL 1970 (523) 95, 93  
 SELING 1966 (263) 55, 52  
 SELTZER 1970 (569) 115, 174  
 SEMAR, TRESER and LANGE 1967 (A 348) 198  
 SESSA and SANNA 1966 (A 527) 209  
 SCASSELLATTI SFORZOLINI and SAVINO 1968 (137) 31, 29  
 SFORZOLINI and SAVINO 1970 (A 305) 196  
 SHAFER, SMILAY and MacMILLAN 1965 (264) 55, 52  
 SHAW, CINKOTAI and THOMSON 1966 (A 445) 205  
 SHIDA and KUROIWA 1969 (A 528) 209  
 SHIELDS 1971 (92) 23, 20  
 SHIMOJIMA 1970 (A 529) 209  
 SHINTANI 1968 (227) 50, 48  
 SHIRABE, MAWATARI and KUROIWA 1970 (A 717) 223  
 SHIRAKI 1969 (A 530) 209  
 SHIRUKI and TATETSU 1967 (A 531) 209  
 SHOJI, YAMAMOTO, NISHIDA, ISHIKAWA, TAKADA and INOUE 1967 (A 306) 196  
 SIASEV 1966 (A 698) 220  
 SIEGEL and MOHLER 1969 (A 184) 188  
 SIEGENTHALER 1965 (A 232) 191  
 SIEGRIST 1966 (A 718) 224  
 SIEVERS, EDWARDS, MURRAY and SCHRENK 1942 (60) 13, 14  
 SIGGAARD-ANDERSEN, KJELDSSEN, PETERSEN and ASTRUP 1967 (387) 73, 70

1005051310

SIGGARD- ANDERSON, PETERSEN, HANSEN  
and MELLEMGAAARD 1968 (349)  
67, 66

SIGGAARD-ANDERSEN, PETERSEN, HANSEN  
and MELLEMGAAARD 1969 (350) 68, 66

SIKAND and PIPER 1966 (A446) 205

SILVER 1971 (A 307) 196

SILVERMAN and GARDNER 1965 (A 349) 198

SIMONE, REGGIANI and BET 1965 (A 447) 205

SIMPSON and RITCHIE 1968 (A 836) 231

SIRS 1964 (A 144) 184

SJOSTRAND 1948 (436) 80, 79

SJOSTRAND 1970 (A 233) 191

SLATER 1950 (302) 60, 58

SLATER 1967 (A 350) 198

SLUIJTER 1967 (A 837) 231

SMALL, RADFORD, FRAZIER, RODKEY and  
COLLISON 1971 (A 44) 178

SMITH 1965 (A 838) 231

SMITH and BRANDON 1970 (A 699) 220

SMITH, BRIERLEY and BRANDON 1971 (A 532)  
209

SMITH, BRYAN, FELESTEIN, LEVADIE,  
MILLER, STEPHENS and WHITE 1970  
(A 352) 199

SMITH, BRYAN, FELDSTEIN, LEVADIE,  
MILLER, STEPHENS and WHITE 1970  
(A 353) 199

SMITH, BRYAN, FELDSTEIN, LEVADIE,  
MILLER, STEPHENS and WHITE 1970  
(A 354) 199

SMITH, BRYAN, FELDSTEIN, LEVADIE,  
STEPHENS and WHITE 1970 (A 351) 198

SMITH, BRYAN, FELDSTEIN, LEVADIE, MILLER  
and WHITE 1972 (A 355) 199

SMITH 1966 (A 356) 199

SMOKING and HEALTH 1964 (634) 114, 116

SNASHALL 1970 (A 774) 227

SNYDER 1970 (A 533) 209

SOBOTKA and SOBOTKA 1969 (303) 60, 57

SOKOLOVSKY, PINCHUK and PRAVOVEROV  
1967 (A 260) 192

SOLVSTEN and KRISTJANSEN 1968 (388)  
73, 69

SPINAZZOLA, MARRACCINI, DEVOTO and  
ZEDDA 1966 (A 302) 196

SRCH 1967 (190) 41, 36, 38

STAMM 1967 (A 357) 199

STANKOVIC, KANTA, FOCO and ALJINOVIC  
1964 (472) 86, 85

STATHERS, HAEGER-ARONSEN, JONSSON  
and MARCIC 1968 (234) 191

STEINER, FRAYSER and ROSS 1965 (A 448)  
205

STEINER, LARSEN, DONATH and PAULI  
1971 (560) 101, 100

STEINMAN 1937 (265) 55, 51

STENIAGEN 1959 (138) 31, 29

STEVENS 1968 (A 185) 188

STEWART, FISHER, HOSKO, PETERSON,  
BARETTA and DODD 1972 (A 719) 224

STEWART, PETERSON, BARETTA,  
BACHAND, JOSKO and HERRMANN  
1970 (497) 92, 89

STICHNOTH and ZUMBANSEN 1970 (A 670) 220

STOKINGER 1969 (A 261) 192

STOKOWSKI and KESIAK 1969 (593) 106, 102

STORMER 1938 (266) 55, 51

STRZELCZYK and ZENK 1964 (473) 87, 85

STUPFEL 1970 (A 236) 191

STUPFEL and BOULEY 1970 (525) 95, 93

STUPFEL, BOULEY, DEKOV, BOURGEOIS  
and ROUSSEL 1968 (524) 95, 93

STUPFEL, BOULEY and POLIANSKI 1970 (611)  
110, 103

STUPFEL and GODIN 1969 (A 197) 189

STUPFEL and ROUSSEL 1968 (A 235) 191

STURNER 1971 (A 641) 217

SUCHCICKI 1970 (A 671) 221

SUGIMOTO and YASUMITSU 1969 (A 839) 231

SULOTTO, BONZANINO, MEO and RUBINO  
1969 b (352) 68, 66

SULOTTO, MEO, POLI and RUBINO 1969a  
(351) 68, 66

SUNDSTROM 1970 (A 45) 178

SUNDSTROM 1972 (A 46) 178

SUPFLE 1933 (A 72C) 224

SUPFLE 1934 (353) 68, 66

SUZUKI 1969 (304) 60, 57

SWINNERTON 1971 (A 200) 189

SWINNERTON, LINNENBOM and LAMONTAGNE  
1970 (A 199) 189

SWINNERTON, LINNENBOM and LAMONTAGNE  
1970 (A 198) 189

SZADKOWSKI, MASTALL, SCHALLER and  
LEHNERT 1970 (191) 41, 38

SZILAGYI 1967 (594) 106, 102

SZLIWOWSKI and KLEES-DELANGE  
1970 (455) 83, 81

SZOLLOSI, MEDVE and JENEY 1970 (329)  
65, 61

SZUCHOVSKY, KENYERES and HARSANYI  
1969 (A 642) 217

TABER and MORRISON 1964 (A 612) 215

TACCOLA, JEDRYCHOWSKI and CAVALLERI  
1965 (456) 84, 81

TAKAHATA and MIYAGISHI 1969 (527)  
95, 98

TAKAMATSU, TAKEICHI and YUKITAKE  
1969 (A 534) 209

TAKEYA, TAKANO, TAMURA, HOJO,

YOSIDA and HURUKAWA 1970 (A 840) 232

TANAKA 1967 (595) 106, 102

TANIEWSKI and KUGLER 1964 A (474)  
87, 85

TANIEWSKI and KUGLER 1964 B (475) 87, 85

TARTULIER, TOURNAIRE, DEYRIEUX and  
BLUM 1967 (A 449) 205

TATEGAMI 1968 (A 775) 227

TATETSU, HARADA, NAKAMURA,  
KASAGI, ISIKAWA and KAMANO 1968  
(547) 98, 96

1005051311



- TATETSU, KIYOTA, TOYA, TEROKA,  
FUJITA, INQUE, MIMURA, HARADA,  
TAKAGI, YAMAGATA, KOZUMA,  
MIYAGAWA, TOMONARI, TERAOKA,  
MURAYAMA, YASUOKA, MIYOSHI and  
KASAGI 1967 A (545) 98, 96
- TATETSU, TOYA, FUJITA, INQUE,  
HARADA, TOMONARI, MURAYAMA  
YASUOKA, MIYOSHI and KASAGI 1967b  
-98, 96
- TATETSU, TOYA, MIMURA, HARADA and  
TSUKAYAMA 1969 (A 721) 224
- TAYLOR and MILLER 1965 (A 48) 178
- TEBBENS and SPEAR 1971 (A 722) 224
- TEICHNER 1967 (528) 95, 93
- TEMMERMAN and ETIENNE 1969 (A 841)  
232
- THEODORE, O'DONNELL and BACK 1971  
(425) 78, 75; (33) 10, 6
- THIELS, DURME, VERMEIRE and  
PANNIER 1972 (267) 55, 52
- THOMAS and PEARSE 1964 (528a) 95, 93
- THOMSON 1971 (A 672) 221
- THURSTON 1968 (A 723) 224
- THURSTON 1968 (A 842) 232
- THURSTON 1970 (A 776) 227
- THURSTON 1971 (A 843) 232
- TIBBLIN 1971 (330) 65, 61
- TIBBLING 1969 (476) 87, 85
- TIMMONS 1970 (A 613) 215
- TKACHENKO, TISHCHENKO, ZATSEPIN  
and DIMITROVA 1966 (426) 78, 74
- TLUSTY, HLOUSKOVA, KROFTA and DAUM  
1972 (A 540) 205
- TOKANA 1971 (A 47) 178
- TOMASINI 1967 (A 535) 210
- TOMONARI 1968 (548) 98, 96
- TONOMURA, YAMATE and TSUJI 1967  
(A 262) 193
- TORELLI 1964 (A 777) 227
- TOTH 1907 (139) 31, 29
- TOTSUKA, MORO, HORIE and YZAKI 1971  
(549) 98, 96; (A 673) 221
- TOWNSEND and STETSON 1968 (A 778) 227
- TOYA 1967 (A 674) 221
- TOYAMA 1968 (A 263) 193
- TRAKHTENBERG 1966 (A 675) 221
- TRILLET, GIRARD and BOULETREAU 1970  
(A 536) 210
- TRINDER and HARPER 1962 (61) 18, 15;  
(163) 35, 33
- TRINQUET, CLAUZEL, CARRE and  
MEYER 1967 (A 451) 205
- TRINQUET and MEYER 1971 (199) 44, 43
- TROMPEO, TURLETTI and GIARRUSO  
1964 (A 309) 196
- TRONZANO and COSCIA 1964 (A 537) 210
- TROUTON and EYSENCK 1961 (197 A) 92, 88
- TRUDINGER 1970 (A 614) 215
- TRUHAUT, BOUDENE and CLAUDE 1965  
(612) 110, 108
- TRUHAUT, BOUDENE and CLAUDE 1967  
(A 163) 186
- TRUHAUT, BOUDENE and CLAUDE 1968  
(389) 73, 70
- TRUHAUT, BOUDENE and CLAUDE 1968  
(427) 78, 75
- TRUHAUT, BOUDENE, CLAUDE, JACOTOT  
1968 (354) 66, 66
- TUTT 1970 (A 676) 221
- TZAGOLOFF 1965 (A 145) 184
- UBISCH and WESTERLUND 1971 (A 314) 196
- UMEZAWA 1968 (A 310) 196
- VALIC and DURIC 1954 (62) 18, 14; 164,  
35, 33
- VANDENBERGH, BILLIET, WOESTIJNE  
and GYSELEN 1968 (A 452) 205
- VAN LIEW 1968 (356) 68, 66
- VAN LIEW 1968 (355) 68, 66
- VAN LIEW 1970 (357) 68, 66
- VANNESTE 1966 (A 358) 199
- VANROUX and GREGOIRE 1964 (A 453) 205
- VAN VUGT 1968 (428) 78, 74
- VARESE and SORANZO 1968 (A 643) 217
- VAUGHAN, JENNELLE and LEWIS 1969  
(228) 50, 48
- VEIL 1968 (A 311) 196
- VEITH 1940 (305) 60, 58
- VENGERSKAIA, NAZOROV, BOBROVA  
SUBROVSKY and DUMKO 1968 (A724224
- VENNESLAND and JETSMANN 1971  
(A 615) 215
- VICH 1969 (A 677) 221
- VIDAL and PICARD 1967 (A 538) 210
- VIEWEG, GRUNEWALD and ZIEGLER 1970  
(A 644) 217
- VIOLET and PERROT 1967 (A 678) 221
- VIVOLI and PREITE 1966 (A 679) 221
- VOGEL 1968 (A 645) 217
- VOGEL, WHEELER and WHITTEN 1972 (307)  
60, 56
- VOGEL and GLESER 1972 (306) 60, 56
- VON BERGMANN 1934 (268) 55, 52
- VON OETTINGEN, DONAHUE and VALAER  
1941 (358) 68, 66
- VOORHOEVE, REMEIK, FREELAND and  
MATTHIAS 1972 (A 312) 196
- VORONCHUCK 1966 (A 680) 221
- VOROSMARTI, BRADLEY, LINAWEAVER,  
KLECKNER and ARMSTRONG 1970 (A 186)  
188
- VUIA 1967 (A 539) 210
- VUOPALA, HUHTI, TAKKUNEN and HUIKKO  
1970 (A 725) 224
- VYSKOCIL 1956 (359) 68, 66
- WAGNER 1964 (A 726) 224
- WAGNER and RICHTER 1968 (A 779) 227
- WAGNER, MAZZONE and WEST 1971 (A 451)  
205
- WAGNER, LATHAM, BRINKMAN and FILLEY  
1969 (A 455) 205
- WAHL 1899 (110) 31, 27
- WAJGT 1971 (437) 80, 79

1005051312

## Additional Bibliography List No 10

- KOTTER D, HUCH A, STOTZ H and PIPER J : Single breath CO diffusing capacity in anesthetized dogs with increased oxygen consumption. Resp Physiol 6: 202-8, 1969. A 411
- KRAL B, CERNOCHOVA Z and TUSL M : Diffusing capacity of the lungs for CO and its components in healthy men of different age at rest and in physical load. Sborn Ved Prac Lek Fak Karlov Univ 9: 625-9, 1966. A 412
- KREUKNIET J and VISSER B F : The pulmonary CO diffusing capacity according to Bates and according to Filley in patients with unequal ventilation. Pflug Arch Ges Physiol 281: 207-11, 1964. A 413
- KREUZER F and CAMPAGNE P L : Resting pulmonary diffusing capacity for CO and O<sub>2</sub> at high altitude. J App Physiol 20: 519-22, 1965. A 413a
- LACOSTE J and ROUCH Y : Mesures simultanées chez l'Homme de l'efficacité des échanges pulmonaires pour le CO<sub>2</sub> et le CO avec addition d'un espace mort. (Simultaneous measurements in humans of the efficiency of pulmonary exchanges for CO<sub>2</sub> and CO with addition of a dead space.) C R Soc Biol(Paris) 160: 1667-70, 1966. A 414
- LACOSTE J : Exploration fonctionnelle respiratoire. La ductance DuCO (monoxyde de carbone): évaluation globale, non sanglante, de l'échangeur pulmonaire. (Functional respiratory investigations. Ductance of CO (carbon monoxide): global estimation of pulmonary exchanger, omitting blood tests.) Presse Med 79: 1781-4, 1971. A 415
- LAWSON W H Jr : Rebreathing measurements of pulmonary diffusing capacity for CO during exercise. J Appl Physiol 29: 896-900, 1970. A 416
- LAWSON W H Jr : Effect of drugs, hypoxia, and ventilatory maneuvers on lung diffusion for CO in man. J Appl Physiol 32: 788-94, 1972. A 417
- LEWIS C M and BRINK A J : Carbon monoxide diffusion of lungs in assessment of pulmonary blood flow in patients with intracardiac shunts. Brit Heart J 28: 359-65, 1966. A 418
- LO COCO A : Sul comportamento della capacità di diffusione per il monossido di carbonio in due gruppi di soggetti affetti da tubercolosi polmonare e da broncopatia cronica ostruttiva. (Behavior of diffusion capacity for carbon monoxide in 2 groups of subjects with pulmonary tuberculosis and chronic obstructive bronchopathy.) G Ital Mal Torace 24: 27-30, 1970. A 419
- LOPEZ-MAJANO V : Reproducibility of the carbon monoxide diffusion capacity method. Respiration 28: 114-9, 1971. A 420
- MAUDERLY J L : Steady-state carbon monoxide-diffusing capacity of unanesthetized Beagle dogs. Am J Vet Res 33: 1485-91, 1972. A 421
- MÄHRLEIN W, KRAUSE M and MÜLLER H R : Apnoetechnik zur Bestimmung der pulmonalen Diffusionskapazität für Kohlenmonoxid. (Apnea technic for the determination of the pulmonary diffusion capacity of carbon monoxide.) Z Ges Med 22: 373-5, 1967. A 422
- MENKES H A, SERA K, ROGERS R M, HYDE R W, FORSTER R E II and DuBOIS A B : Pulsatile uptake of CO in the human lung. J Clin Invest 49: 335-45, 1970. A 423
- MITCHELL M M and RENZETTI A D Jr : Application of the single-breath method of total lung capacity measurement to the calculation of the carbon monoxide diffusing capacity. Am Rev Resp Dis 97: 581-4, 1968. A 424
- MITTMAN C : Nonuniform pulmonary diffusing capacity measured by sequential CO uptake and washout. J Appl Physiol 23: 131-8, 1967. A 425
- NAIRN J R, POWER G G, HYDE R W, FORSTER R E, LAMBERTSEN C J and DICKSON J : Diffusing capacity and pulmonary capillary blood flow at hyperbaric pressures. J Clin Invest 44: 1591-9, 1965. A 426

1005051267

## VI. NERVOUS SYSTEM

## C. Hearing and Auditory Pathways

- |  | Reprint |
|--|---------|
| CIS G and PERANI G: La funzionalita cocleo-vestibolare nell'ossicarbonismo. (Cochleo-vestibular function in carbon monoxide poisoning.) <u>Arch Ital Otol</u> 75: 635-43, 1964.  | 458     |
| FORTUNATO V and CATALANO G B: Ipoacusie e sordita da intossicazioni esogene. (Bradyacusia and deafness caused by exogenous intoxications.) <u>Minerva Otorinolaringol</u> 20: 193-217, 1970.   | 459     |
| FREIGANG B, SEIDEL P and FLACH M: Zur Frage der Kohlenmonoxydauswirkung auf die Mikrophon- und Summenaktionspotentiale der Meerschweinchencochlea. (On the problem of carbon monoxide effect on the microphonics and sum of action potentials of the guinea pig cochlea.) <u>Arch Klin Exp Ohr Nas Kehlk</u> 190: 24-35, 1968.                               | 460     |
| FRITSCH F: Richtungsaudiometrische Untersuchungen im freien Schallfeld bei U-förmig verlaufenden Horschwellenkurven. (Directional audiometric studies in a free field on patients with U-shaped audiograms.) <u>Z Lar Rhin Otol</u> 48: 291-302, 1969.   | 461     |
| GUEST A D L, DUNCAN C and LAWTHORP P J: Carbon monoxide and phenobarbitone: A comparison of effects on auditory flutter fusion threshold and critical flicker fusion threshold. <u>Ergonomics</u> 13: 587-94, 1970.  | 462     |
| HALL G H: Effects of nicotine and tobacco smoke on the electrical activity of the cerebral cortex and olfactory bulb. <u>Brit J Pharmacol</u> 38: 271-86, 1970.  | 463     |
| HANSZ J and STYPEREK J: Jednostronne uszkodzenie nerwu sluchowego w ostrym zatruciu tlenkiem wegla. (Unilateral damage of the acoustic nerve during acute carbon monoxide poisoning.) <u>Pol Tyg Lek</u> 23: 1441-2, 1968.   | 464     |
| KAWAMURA S: (Clinical observation of tinnitus caused by extrinsic causes - head trauma, acoustic trauma, etc.) <u>J Otol Jap</u> 74: 1007-27, 1971.  | 465     |
| KITTEL G and THEISSING-ERLANGEN G: Histologische Untersuchungen der Cochlea an Häutchen-Präparaten und Treppen-Serienschnitten nach hochgradiger protrahierter Hypoxydose. (Histological studies of the cochlea on cuticle preparations and scalari-form sections following severe protracted hypoxia.) <u>Arch Klin Exp Ohr Nas Kehlk</u> 191: 534-8, 1968. | 466     |
| KÜTTNER K: Zur Pathomorphologie der Veränderungen am peripheren Hörorgan bei wiederholter experimenteller Kohlenmonoxydintoxikation. (On the pathomorphology of changes in the peripheral organ of hearing during repeated experimental carbon monoxide intoxication.) <u>Z Laryng Rhinol Otol</u> 47: 779-85, 1968.   | 467     |
| MESOLELLA C, PERRELLA F, TESTA B and MORELLI G: Rilievi oto-neurologici in un gruppo di operai esposti cronicamente all'azione dell'ossido di carbonio. (Otoneurological observations in a group of workmen chronically exposed to the action of carbon monoxide.) <u>Arch Ital Laring</u> 78: 47-60, 1970.  | 468     |
| MORRIS T M O: Deafness following acute carbon monoxide poisoning. <u>Laryng</u> 83: 1219-25, 1969.   | 469     |
| MOUNIER-KUHN P, ROCHE L, MORGON A and BERNARD P: Les atteintes vestibulaires au décours immédiat des intoxications aiguës par l'oxyde de carbone. (Vestibular involvement immediately following acute carbon monoxide poisoning.) <u>J Franc Otorhinolaryng</u> 17: 512-5, 1968.   | 470     |
| SATO T: (Hearing disturbances in monoxide-gas toxicosis.) <u>Otolaryngology / (Tokyo)</u> 38: 805-16, 1966.  | 471     |
| STANKOVIĆ D, KANTA F, FOCO S and ALJINOVIC M: Oštećenja sluha posle ponovljenih akutnih trovanja ugljen-monoksidom. (Hearing disorders following repeated carbon monoxide poisoning.) <u>Acta Med Jugosl</u> 18: 95-106, 1964.   | 472     |

1005051150